



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

N. 990

Sp. Libri J. L. van Loon

H. N. 990

LIOTHEEK GENT



0000001608

Digitized by Google

FLORA FRIDRICHSDALINA

SIVE

METHODICA DESCRIPTIO
PLANTARUM

IN

AGRO FRIDRICHSDALENSI

SIMULQUE

PER

REGNUM DANIÆ

CRESCENTIUM

CUM

CHARACTERIBUS genericis & specificis; NOMINIBUS
trivialibus, vernaculis, pharmaceuticis; LOCIS
NATALIBUS specialissimis; ICONIBUS optimis
allegatis, ac SPECIEBUS pluribus
in Dania nuper detectis.

Cam Permissu Superiorum.

ARGENTORATI

In Officina Libraria JOH. GODOFR. BAVERI

1767.



Inest in explicatione Naturæ infatiabilis quædam
e cognoscendis rebus voluptas; in qua una con-
fectis rebus necessariis, vacui negotiis, honeste
ac liberaliter possumus vivere.

CICERO de *F. B. Lib. 4.*



CHRISTIANO VII.

DANIÆ NORVEGIÆQUE

R E G I,

VANDALORUM GOTHORUMQUE,

DUCI SLESVICI, HOLSATIÆ,

STORMARIÆ & DITMARSII,

COMITI IN OLDENBURG

& DELMENHORST

&c. &c. &c.

Augustissime & Potentissime
R E X!

*Flores MAJESTATI TUÆ spargit
Danica Flora. Hæc TIBI & sibi
trecentas vindicat stirpes, quas nulli
majorum patria obtulit tellus. Au-
spiciis TUIS reservatum erat inco-
gnita scrutari gramina, minimas
pervestigare plantulas, earumque oc-
cultas nuptias patefacere.*

*Flóreant sic Regna TUA PATRIS
virtutibus, & qui nunc radiis
MAJESTATIS TUÆ hilarantur
Juvenes, Nestorei olim Senes TIBI
gratos adducant Nepotes vota Majo-
rum continuaturos.*

*Floræ Danicæ omnes se Musæ jun-
gant, TUUMQUE Imperium florentis-
simum ac immortale ad omnes posteros
reddant.*

Ita vovet & auguratur

**Sacratissimæ
MAJESTATIS TUÆ**

*Augustæ Taurinæ
d. 26. Julii 1766.*

devotissimus Cultor & Subditus
OTTO FRIDERICUS MULLER.



P R Æ F A T I O.



Plurimam otii mei partem per annos aliquot in Plantarum cryptogamiarum, *Filicum* nempe, *Muscorum*, *Algarum* & *Fungorum* indagatione fuisse absumtam, olim jam in *Præfatione Faunæ Fridrichsdalinæ* monui. Scilicet quæ Botanicorum studia terminare alias solet classis omnium difficillima & maxime intricata, cum & longe serius quam reliquæ plantæ coli cœperit, & specierum exiguitate ac multitudine laboret, & natalis locus florendique tempus scrutantem deterreat, mihi perverso fere ordine cognitionis herbariæ initium fuit, & quæ hucusque scandala Botanices audiverunt stirpes, a quibus exercitati quoque Viri, nedum tyrones abhorruerunt

runt, eæ vixdum delibatæ mihi Scientiæ auspicia, stimuli & præmia extiterunt. Tum demum, ubi fervorem & impetum novi aliquid in patria detegendi minutissimæ, quibus inhiam, plantukæ paulatim deslitterent, in splendidiorem Floræ hortum exspatiatus, majores, quas hæctenus neglexeram, plantas legere cœpi, non majori quidem cum voluptate, at minori vitæ ac sanitatis dispendio.

Illis enim regni vegetabilis minimis incolis habitationes fecit divinum Numen rupes, speluncas, saxa, declivia, præcipitia, proceras arbores, tecta ædium & rudera, paludes, lacus, scaturigines, maria, locaque vitæ nisi periculo plerumque inaccessa, ut sterilissima quoque ac hominibus neglecta quærenti præberent theatrum potentiæ ac sapientiæ summi Conditoris improvisum. His apricos campos, lætantes agros, virentia prata, umbrosas sylvas, salubres montes ac rigidas valles, ut quæ utilissima ac magis necessaria ubique sensibus obvia homines & animalia sensatione dulcissima ad fruendum allicerent. Illis ne globus terraqueus unquam virore ac floribus careret, florendi tempus concessit serum autumnum, tristem hyemem, verisque initia tempestate maxime pluvia: his formosissimum annum, ridens ver, prægnantem æstatem, gratumque autumnum.

Ex utrorumque assidua indagatione tandem exsurgit enumeratio hæc plantarum *Agri Friedrichsdalensis*. Omnes ac singulas ipsemet quæ-

sivi, investigavi, collegi, disposui, opera quidem haud parva at delectatione perquam eximia. *ΑΥΤΟΔΙΔΑΚΤΟ* enim, nullius viva voce institutus, nullo herbario adjutus, nullo socio suffultus, libros Botanicorum noscere, terminos artis subolere, intelligere & discere tandem aliquando post vices reiteratam plantæ ac descriptionis comparisonem primum licuit. In paludosis circumvagari, in declivia repere, a præruptis pendere, hic labor, hoc opus fuit. Quavis vero excursiuncula cognitiones augere, novam plantam detegere, vel novi quid in jam nota observare, hoc operæ pretium, hoc dulce curarum lenimen. Sic crescente indies cognitione ac collectione nascitur Flora, qualem quoad methodum & numerum plantarum nunquam protulit patria; quoad copiam in regione tam exigui ambitus nullus unquam exterorum.

Dedit centum abhinc annis PETRUS KYLLING, Botanicus regius, compendiosam enumerationem plantarum Regni Daniæ indigenarum secundum BAUHINUM.

TYCHO HOLM, herbarum ac Medicinæ Studiosus, posthac Oeconomix Professor Hafniensis, pro gradu Doctoris sub præsidio Illustrissimi a LINNE Upsaliæ 1757 plantas a KYLLINGIO recensitas ad normam methodi sexialis disposuit, solisque nominibus trivialibus expressit, paucas ex herbario BURSERI, paucas a se intra Daniam lectas adjecit. Dissertatio extat in *Amæn. Acad. T. V, No. 78.*

Clarissimi OEDERI, Professoris Botanices, quis ignorat Floram latissimo sensu Danicam, stirpes quippe omnium provinciarum Regi Daniæ subjectarum a Mari Glaciali trans Albim usque, quin & Islandicas complexuram; quam post excursionem plurium annorum in Norvegiæ Daniæque provincias, pictore stipatus impensis publicis peractas, iconibus graphice sculptis ac eleganter pictis, munificentia Regis Optimi publici juris fieri curat. Opus & pulchritudine & impensis regium, at ob insigne pretium, quo opus adeo nitide pictum necessario venire debet, paucorum usui. Exstant Fasciculi quatuor *), plantas ducentas quadraginta variarum regionum & classium continentes.

Quid in reherbaria præstiterint Ol. BORRICHIUS, BARTHOLINUS, C. PAULI, BUCHWALD, haud moror, cum ultra plantas officinales progressi non sint.

Ut, quantum possit, commodo publico inservirem, plantis hujus Floræ concinnos ac concisos Linnæanos characteres genericos compendiosos ex *Syst. Nat. edit. X.* & specificos ex *Spec. Edit. novissima*, Holm. 1762. optimas Auctorum Icones, loca natalia specialis-

*) Quum ob Auctoris Italicum iter & longiorem moram retardata libri impressio fuerit, interea temporis autem quintus Fasciculus *Flora Danica* OEDERIANÆ in lucem prodierit, Editor, ne aliquid Operi deesse videretur, haud ingratam operam & Lectoribus & Auctori se præstiturum esse putavit, si quinti hujus Fasciculi plantas vere Danicas adjiciendas curaret, quarum proinde Catalogum decima quarta pagina Tibi B. L. exhibebit.

fima, nomina trivialia, vernacula & pharmaceutica cum pauciusculis Observationibus adjunxi, & ut Flora Fridrichsdalina simul esset Flora Danica, i. e. una contineret omnes & singulas plantas ad hunc diem per Regnum Daniæ lectas, & in herbationibus per ejus provincias sufficeret, paucas, quæ Agro Fridrichsdalensi exulant, KYLLINGII, HOLMII, Cel. OEDERI & propria auctoritate adposui. Hæ signo crucis †, uti stirpes a me primum intra patriam detectæ Asterisco * notantur.

Numerus cuique generi a dextra adpositus indicat numerum generis in sexta Generum editione. *Gen. Pl.* LINN. Hohniæ 1764.

Phascum ab Illustriss. DE HALLER, Cel. SCHMIDEL, & a me in *Act. Holm.* descriptum, a Perillustri A LINNE' haud tamen in numerum Plantarum relatum suo loco inserui.

Tria nova *Polypodia* Botanicis inquirenda dedi.

Jungermanniarum manifestas nuptias monstravi.

Insecta ducenta sexaginta, quæ intra anni spatium in Agro Fridrichsdalensi legi, *Fauna* inferenda apposui.

Triplicem nominum Genericorum, Vernaculorum & Pharmaceuticorum addidi indicem.

Plantarum nomina vernacula invitus adpono; mallem enim nulla omnino alia existere, quam quæ Systematicis respondent ut

difficultates, confusiones & damna evitarentur, quæ inde oriuntur, quod plures species eodem, eademque species diverso nomine vernaculo dicantur, communi linguarum omnium vivarum vitio; hinc plurima, quæ a KYLLINGIO & aliis recensentur Danica nomina e Latina & Germanica lingua translata lubens omisi, vera nomina vulgo usitata servans.

Optassem quidem, ut in inquirendis & disponendis stirpibus haberem, qui mecum communi studio insudavissent, quibusque gratias publice referrem; at frustra quæsi. Soli mihi & in Entomologia & in Botanica sudor ac pulvis omnino relictus. Haud tamen deerunt in patria tantillæ gloriæ invidi; hos uti contra *Faunam* insurgentes immotus expertus sum, ita in *Floram* tranquillus expecto. Discant, precor, hi rerum haud æqui aestimatores, qui animos ac libros potius quam ipsam naturam horum interpretem consulunt, *res naturales*, uti præclare monuit Illustr. A LINNE', *proprio privilegio munitas persistere, & quemadmodum errores in his commissos a nullo defendi, ita nec veritates Observationibus innixas a toto Eruditorum orbe conculcari posse.*

Argentorati, decimo sexto Aprilis,

1766.

OTTO FRIDERICUS MULLER

Acad. Cæsar. N. C. Elect. Boic. Scient.
& Societ. Helv. Sodal. S.

ADDENDA & CORRIGENDA.

Ad pag. 17. add. No. 77. b. † *POA maritima*, panicula subspicata, spiculis secundis coarctatis, foliis convolutis.
Huds. Angl. 35.

Fl. Dan. t. 251.

Vulgaris in litoribus.

Ad pag. 32. post Flores *incompleti*, *inferi*, ante
GLAUCEM

adde 83. b. ILLECEBRUM. *Cul.* 5-
phyllus cartilagineus. *Cor.* 6.
Stigma simplex. *Caps.* 5-
valvis, monosperma.

& ad pag. 46. ante GLAUCEM adde

83. b. ILLECEBRUM. 290.

198. b. * ILLECEBRUM *verticillatum*, floribus verticillatis nudis, caulibus procumbentibus.

In arenosis aridis Ducatus *Holsatiae*.

pag. 132. Spec. 585. loco stipitulis lege stipulis.

Ad pag. 161. Spec. 719. præfigatur †.

addatur *Fl. Dan.* t. 103.

In agris *Silleröd & Charlottenlund*.

Planta enim a Clar. Oedero pro *OPHY* *anthropophora*
Tab. 103. tradita, est *ORCHIS ustulata*.

Ad pag. 163. Spec. 729. expungatur tota, quam sola *Oederi* auctoritate addideram, quamque *ORCHIDEM ustulatam* esse nunc edoctus sum.

Ibid. add. No. 732. b. † *HELLEBORINE* latifolia, floribus magnis, paucis albisque. *Hall. Helv.* 276.

Fl. Dan. t. 267.

In pratis sylvaticis subuliginosis rarior.

pag. 164. ad Spec. 732. deleantur verba:

In præcipitiis promontorii Insulæ *Monæ*, vulgo
Klint, legi rarissimam plantam.

Ad pag. 192. No. 838. b. † *ASPLENIUM* *Adiantum nigrum*, frondibus subtripinnatis: foliolis alternis: pinnis lanceolatis inciso-ferratis. *Fl. Dan.* t. 250.

In Insula *Bornholm*, in litoris orientalis petrarum rimis,


pag. 193. lin. 14. l. duos vel tres acutissimos.

pag. 214. Spec. 984. loco pyxidatus, lege *pyxidatus*.

Sequentium Plantarum O E D E R I A N Æ Icones

*ex quinto Fasciculo Floræ Danicæ
suppleantur.*

- Spec.* 7. *Circæa alpina* Tab. 256.
 10. *Veronica officinalis*. 248.
 21. ——— *verna*. 252.
 37. *Scirpus palustris*. 273.
 39. ——— *acicularis*. 287.
 48. *Phalaris arundinacea*. 259.
 78. *Briza media*. 258.
 89. *Bromus arvensis*. 293.
 100. *Arundo Calamagrostis*. 280.
 112. *Scabiosa succisa*. 279.
 126. *Plantago maritima*. 243.
 127. ——— *Coronopus*. 272.
 193. *Rhamnus Frangula*. 278.
 216. *Gentiana Pneumonanthe*. 269.
 223. *Sanicula Europæa*. 283.
 229. *Selinum palustre*. 257.
 236. *Sium latifolium*. 246.
 237. ——— *nodiflorum*. 247.
 270. *Allium scorodoprasum*. 290.
 289. *Convallaria bifolia*. 291.
 316. *Epilobium angustifolium*. 289.
 330. *Polygonum amphibium*. 282.
 361. *Silene nutans*. 242.
 363. *Stellaria nemorum*. 271.
 585. *Vicia sylvatica*. 277.
 645. *Serratula tinctoria*. 281.
 776. *Orchis morio*. 253.
 720. ——— *latifolia*. 266.



CLASSIUM

CHARACTERES.

I. MONANDRIA.

Stamen unicum in flore hermaphrodito.

II. DIANDRIA.

Stamina duo in flore hermaphrodito.

III. TRIANDRIA.

Stamina tria in flore hermaphrodito.

IV. TETRANDRIA.

Stamina quatuor in flore hermaphrodito.

(Si Stamina duo breviora sunt, referatur ad *Class. XIV.*)

V. PENTANDRIA.

Stamina quinque in flore hermaphrodito.

VI. HEXANDRIA.

Stamina sex in flore hermaphrodito.

(Si ex his stamina duo opposita breviora, pertinet ad *Class. XV.*)

VII. HEPTANDRIA.

Stamina septem in flore hermaphrodito.

VIII. OCTANDRIA.

Stamina octo in flore hermaphrodito.

IX. ENNEANDRIA.

Stamina novem in flore hermaphrodito.

X. DECANDRIA.

Stamina decem in flore hermaphrodito.

XI. DODECANDRIA.

Stamina duodecim ad novendecim in flore hermaphrodito.

XII. ICOSANDRIA.

Stamina viginti, sæpe plura calyci adnata in flore hermaphrodito.

XIII. POLYANDRIA.

Stamina receptaculo inserta a 20 ad 1000 in flore hermaphrodito.

XIV. DIDYNAMIA.

Stamina quatuor, quorum duo proxima longiora sunt.

XV. TETRADYNAMIA.

Stamina sex, quorum quatuor longiora, duo autem opposita breviora.

XVI. MONADELPHIA.

Stamina filamentis in unum corpus coalita sunt.

XVII. DIADELPHIA.

Stamina filamentis in duo corpora coalita sunt.

XVIII.

XVIII. POLYADELPHIA.

Stamina filamentis in tria vel plura corpora coalita.

XIX. SYNGENESIA.

Stamina antheris (raro filamentis) in cylindrum coalita.

XX. GYNANDRIA.

Stamina pistillis (non receptaculo) infidentia.

XXI. MONOECIA.

Flores masculi & feminei in eadem planta, non in eodem thalamo.

XXII. DIOECIA.

lores masculi in diversa planta a femineis prognati.

XXIII. POLYGAMIA.

lores hermaphroditi & masculi aut feminei in eadem specie.

XXIV. CRYPTOGRAMIA.

lores intra fructum vel singulari modo occultati.

)(

ORDINES a pistillis, ut **Classes** a staminibus desumuntur; in **Classi Syngenesiæ** autem a cæteris differunt ordines; e. gr.

Monogynia, **Digynia**, **Trigynia** &c. i. e. **Pistillum** 1, 2, 3 &c. **Numerus** hic pistillorum desumitur a basi styli; si stylus autem deficiat, a numero stigmatum calculus fit.

POLYGAMIA ÆQUALIS constat multis flosculis, staminibus ac pistillis instructis.

POLYGAMIA SPURIA, ubi flosculi hermaphroditi discum occupant, & marginem cingunt flosculi feminei, staminibus destituti, idque triplici modo:

- a) *Superflua*: cum flores disci hermaphroditi stigmate instruuntur, & semina proferunt; flores quoque feminei radium constituentes, similiter semina ferunt.
- b) *Frustranea*: cum flores disci hermaphroditi stigmate instruuntur, & semina proferunt; flosculi vero radium constituentes, quum stigmate careant, semina proferre nequeunt.
- c) *Necessaria*: cum flores hermaphroditi ob defectum stigmati pistilli semina perficere nequeunt; flores autem feminei in radio semina perfecta proferunt.



Classis



Classis I.

MONANDRIA.

MONOGYNIA.

1. HIPPURIS. *Calyx* 0. *Corolla* 0. *Stigma simplex.*
Semen 1.
2. SALICORNIA. *Cal.* 1-phyllus, ventriculosus, integer. *Cor.* 0. *Sem.* 1. *Stamen* 1. f. 2.

DIGYNIA.

3. CALLITRICHE. *Cal.* 0. *Cor.* 2 - petala. *Capsula*
2 - locularis, 4 - sperma.
4. BLITUM. *Cal.* 3 - fidus, baccatus. *Cor.* 0. *Sem.* 1



MONOGYNIA.

I. HIPPURIS. II.

I. HIPPURIS vulgaris.

Fl. Dan. t. 87.

In amne & piscinis.

2. SALICORNIA. IO.

2. SALICORNIA *herbacea* patula: articulis apice compressis emarginato-bifidis.

P. Fl. Oec. 128.

Danis *Salt-Urt*; *Glas-Urt.*

Pharmac. SAL ALCALI.

In littore Maris Balthici.

DIGYNIA.

3. CALLITRICHE. 13.

3. CALLITRICHE *verna*, foliis superioribus ovalibus; floribus androgynis.

Fl. Dan. t. 129.

In fossis frequens.

4. CALLITRICHE *autumnalis* foliis omnibus linearibus; apice bifidis; floribus hermaphroditis.

In fossis.

4. BLITUM. 14.

5. † BLITUM *capitatum* capitellis spicatis terminalibus.

Dan. *Skov-Melde.*

Classis II.

D I A N D R I A.

MONOGYNIA.

* Flores *inferi*, *monopetali*, *irregulares*.
Fructus *capsulares*.

- 6. VERONICA. *Cor.* 4-fida, rotata: lacinia infima angustiore. *Capsula* bilocularis.
- 7. PINGUICULA. *Cor.* ringens, calcarata. *Cal.* 2-labiatum, 5-fidus. *Capsf.* 1-locularis.
- 8. UTRICULARIA. *Cor.* ringens, calcarata. *Cal.* 2-phyllus, æqualis. *Capsf.* 1-locularis.
† Flores *apetali*. Fructus *filiculosus*.
Lepidium rudemale.

* * Flores *inferi*, *monopetali*, *irregulares*.
Fructus *gymnospermi*.

- 9. VERBENA. *Cor.* infundibuliformis, subæqualis, curva: *Calycis* unico dente truncato. *Sem.* 2. f. 4. nuda. *Stam.* 2. f. 4.
- 10. LYCOPUS. *Cor.* 4-fida: lacinia unica emarginata. *Stam.* distantia. *Sem.* 4. retusa.
* * * Flores *superi*.

- 7. CIRCEA. *Cal.* 2-phyllus, superus. *Cor.* 2-petala, obcordata. *Sem.* 1, 2-loculare.

DIGYNIA.

- 21. ANTHOXANTHUM. *Cal.* Gluma uniflora, 2-valvis. *Cor.* Gluma 2-valvis, acuminata. *Sem.* 1.

MONOGYNIA.

5. CIRCÆA. 24-

6. CIRCÆA *Lutetiana* caule erecto: racemis pluribus.

Fl. Dan. t. 210.

Dan. St. Stephens-Urt.

In vivario & alibi vulgaris.

7. CIRCÆA *Alpina* caule adscendente: racemo unico.

In ripa elatiori paa Næffet, frequens.

6. VERONICA. 25.

* *Spicata.*

8. † VERONICA *maritima* spicis terminalibus: foliis ternis inæqualiter serratis.

9. VERONICA *spicata* spica terminali, foliis oppositis crenatis obtusis: caule adscendente simplicissimo.

Fl. Dan. t. 52.

In apricis extra hortum versus amnem, rarior.

10. VERONICA *officinalis* spicis lateralibus pedunculatis: foliis oppositis: caule procumbente.

P. Fl. Oec. 154.

Dan. *Arenpriis*,

Pharmac. VERONICÆ herba.

In Sylvis.

* * *Corymboso-racemosa.*

11. VERONICA *Serpyllifolia* racemo terminali subspicato: foliis ovatis glabris crenatis.

In pratis.

12. VERONICA *Beccabunga* floribus racemosis lateralibus: foliis ovatis planis, caule repente.

Dan. *Lemmike*, *Ledmye*.

Pharmac. BECCABUNGÆ herba.

In fossis & scaturiginibus.

13. VERONICA *anagallis aquatica*, floribus racemosis lateralibus: foliis lanceolatis serratis: caule erecto.

Dan. *Vand-arfoe*.

In paludosis.

14. VERONICA *scutellata*, floribus racemosis lateralibus alternis: foliis linearibus integerrimis.

Dan. *Vand-arfve*.

In inundatis.

15. * VERONICA *Teucrium* racemis lateralibus longissimis: foliis ovatis rugosis dentatis obtusiusculis: caule erecto.

In Sylvestribus; frequens in lucis prati *Jonstrup*, semper erecta ac major sequenti, num tamen factis distincta.

16. VERONICA *Chamadrys* racemis lateralibus: foliis ovatis sessilibus rugosis dentatis: caule debili.

In pratis.

* * * *Pedunculis unifloris*.

17. VERONICA *agrestis* floribus solitariis: foliis cordatis incis pedunculo brevioribus.

In arvis & cultis.

18. VERONICA *arvensis* floribus solitariis: foliis cordatis incis pedunculo longioribus.

In ruderatis.

19. VERONICA *hederifolia* floribus solitariis: foliis cordatis planis quinquelobis.

In arvis.

20. † VERONICA *triphyllus* floribus solitariis: foliis digitato-partitis pedunculo brevioribus.

21. * VERONICA *verna* floribus solitariis: foliis digitato-partitis, pedunculo longioribus.

In aridis apricis.

22. † VERONICA *peregrina* floribus solitariis sessilibus: foliis lanceolato-linearibus glabris obtusis integerrimis: caule erecto.

In Insula Amagria.

7. PINGUICULA. 30.

23. PINGUICULA *vulgaris* nectario cylindraceo longitudine petali. *Fl. Dan. t. 93. P. Fl. Oec. 107.*

Dan. *Vibefat*; *Haar-vext*.

In prato humido ad lacum *Bagfuærd* ac in uliginosis frequens.

8. UTRICULARIA. 31.

24. UTRICULARIA *vulgaris* nectario conico : scapo paucifloro. *Fl. Dan. t. 138.*

Dan. Vand-röllicke.

In fossis antiquis turfosis, & in paludibus.

25. UTRICULARIA *minor* nectario carinato.

Fl. Dan. t. 128.

In paludoso extra hortum.

9. VERBENA. 32.

26. VERBENA *officinalis* tetrandra spicis filiformibus paniculatis : foliis multifido-laciniatis : caule foliario.

Dan. Jern-urt.

Pharmac. VERBENÆ herba.

Circa maceriem villæ veteris.

10. LYCOPUS. 33.

27. LYCOPUS *Europæus* foliis sinuato-ferratis.

Dan. Vand-marra.

In ripa stagni ; rarior.

DIGYNIA.

11. ANTHOXANTHUM. 42.

28. ANTHOXANTHUM *odoratum* spica ovato-oblonga : flosculis subpedunculatis arista longioribus.

In pratis frequens.

Classis III.

TRIANDRIA.

MONOGYNIA.

* Flores *superi.*

12. VALERIANA. *Cor.* 1 - petala, 5 - fida, basi gibba, supera. *Sem.* 1. *Cal.* 0.
13. IRIS. *Cor.* 6 - petaloidea : petalis alternis reflexis. *Stigmata* petaliformia.

* * Flores *graminei valvulis glumæ calycinae.*

14. SCHOENUS. *Glumæ* paleaceæ, univalves, congestæ. *Cor.* 0. *Sem.* 1. subrotundum.
15. CYPERUS. *Glumæ* paleaceæ, distiche imbricatæ. *Cor.* 0. *Sem.* 1. nudum.
16. SCIRPUS. *Glumæ* paleaceæ undique imbricatæ. *Cor.* 0. *Sem.* 1. imberbe.
17. ERIOPHORUM. *Glumæ* paleaceæ undique imbricatæ. *Cor.* 0. *Sem.* 1. lanigerum.
18. NARDUS. *Cal.* 0. *Cor.* 2 - valvis. *Sem.* tectum.

DIGYNIA.

* Flores *uniflori vagi.*

20. PANICUM. *Cal.* 3 - valvis, tertia dorsali minore.

22. ALOPECURUS. *Cal.* 2 - valvis. *Cor.* 1 - valvis.
21. PHLEUM. *Cal.* 2 - valvis, linearis, truncatus, apice bicuspidato. *Cor.* inclusa.
19. PHALARIS. *Cal.* 2 - valvis : valvis carinatis : longitudine æqualibus, corollam includentibus.
23. MILIUM. *Cal.* 2 - valvis : valvis ventricosus corolla majoribus. *Stigmata* penicilliformia.
24. AGROSTIS. *Cal.* 2 - valvis : valvis acutis corolla paulo majoribus. *Stigm.* longitudinaliter hispida.
29. DACTYLIS. *Cal.* 2 - valvis, obtusissimus : valva altera longiore carinata.

† *Arundo* epigeios; calamagrostis; arenaria.

* * Flores *biflori*, *vagi*.

25. AIRA. *Cal.* 2 - valvis. Flosculi absque rudimento tertii.
26. MELICA. *Cal.* 2 - valvis. Rudimentum tertii inter-flosculos.

† *Avena* elatior.

* * * Flores *multiflori*, *vagi*.

28. BRIZA. *Cal.* 2 - valvis. *Cor.* cordata : valvis ventricosus. Spicula disticha.
27. POA. *Cal.* 2 - valvis. *Cor.* ovata : valvulis ovato-acuminatis.
31. FESTUCA. *Cal.* 2 - valvis. *Cor.* oblonga : valvis mucronatis.



32. BROMUS. *Cal.* 2 - valvis. *Cor.* oblonga: valvis
infra apicem aristatis.
33. AVENA. *Cal.* 2 - valvis. *Cor.* oblonga: valvis
dorso arista contorta.
34. ARUNDO. *Cal.* 2 - valvis. *Cor.* basi lanata, mu-
tica.
- * * * * Flores *multiflori insidentes receptaculi elon-*
gati dentibus.
35. LOLIUM. *Involucrum* uniphyllum, fixum, mul-
tiflorum. *Cal.* 0.
36. ELYMUS. *Involucrum* diphyllum, spiculis ge-
miniis. *Cal.* 0.
37. HORDEUM. *Involucrum* 6 - phyllum, 3 - florum.
38. TRITICUM. *Cal.* 2 - valvis, subtriflorus.
39. CYNOSURUS. *Cal.* 2 - valvis multiflorus. *Re-*
cept. unilaterialia.

TRIGYNIA.

39. MONTIA. *Cal.* 2 - phyllus. *Cor.* 1 - petala, irre-
gularis. *Caps.* 1 - locularis, 3 - valvis, 3 - sperma.



MONOGYNIA.

12. VALERIANA. 44.

29. VALERIANA *officinalis* floribus triandris: foliis omnibus pinnatis. *P. Fl. Oec.* 153.

Dan. *Oyen-rod*; BALDRIAN.

Pharmac. VALERIANÆ radix, herba.

In paludosis nemoris.

30. VALERIANA *Dioica* floribus triandris dioicis: foliis pinnatis.

In pratis uliginosis.

31. VALERIANA *locusta* caule dichotomo: foliis linearibus: floribus triandris. *P. Fl. Oec.* 152.

Dan. *Fettekaas*; *Feldkrop*.

In arvis.

13. IRIS. 59.

* *Barbata nectariis petalorum reflexorum.*

32. † IRIS *Germanica* corollis barbatis: caule foliis longiore multifloro: floribus inferioribus pedunculatis.

In Insula *Saltholm*.

* * *Imberbes petalis deflexis levibus.*

33. IRIS *pseudacorus* corollis imberbibus: petalis interioribus stigmatibus: foliis ensiformibus.

Dan. *Sverd-Lilie*.

Pharmac. ACORI PALUSTRIS radix.

In paludosis.

14. SCHOENUS. 65.

34. * SCHOENUS *Mariscus* culmo tereti, foliis marginis dorsoque aculeatis. *Morif. Hist.* p. 273. t. II. f. 24.

In lacu *Lyngby* copiose.

35. * *SCHOENUS albus* culmo subtriquetro folioso;
floribus fasciculatis: foliis fetaceis. *Morif. Hist.*
t. 9. f. 39.
In paludosis.

15. CYPERUS. 66.

36. † *CYPERUS fuscus* culmo triquetro nudo: umbella
trifida: pedunculis simplicibus inæqualibus: spi-
cis confertis linearibus. *Fl. Dan. t. 179.*
In pratis paludosis.

Flora Danica huic assignat locum natalem *Fridrichsdal*; ibi-
dem hucusque invenire nequeo.

16. SCIRPUS. 67.

* *Spica unica.*

37. *SCIRPUS palustris* culmo tereti nudo: spica subo-
vata terminali. *Fl. Dan. t. 167.*

Dan. Sif. nomen generis.

Obs. Figura, t. 167. quam *Scirpum acicularem* vult
Cl. *Oederus*, potius offert palustrem; acicularis enim duplo
vel triplo minor, tenuior &c. culmo plerumque curvato.

In ripis ac inundatis frequens.

38. *SCIRPUS cespitosus* culmo striato nudo: spica bi-
valvi terminali longitudine calycis: radicibus
squamula interstinctis. *Scheuchz. Agrost. t. 7. f. 18.*
P. Fl. Oec. 132.

Dan. Myre-Sif.

In paludibus turfosis.

39. *SCIRPUS acicularis* culmo tereti nudo fetiformi;
spica ovata terminali bivalvi: seminibus nudis.
Scheuchz. Agrostog. t. 7. f. 19. Morif. Hist. t. 10.
f. 37.

Sub aqua piscinæ in Sylva *Hareskov* copiose.

40. † *SCIRPUS fluitans* culmis teretibus nudis alternis :
caule folioso flaccido. *Scheuchz. Agrost. t. 7. f. 20.*
In inundatis.

* * *Culmo tereti polystachio.*

41. *SCIRPUS lacustris* culmo tereti nudo : spicis ovatis
pluribus pedunculatis terminalibus.

In stagnis ubique.

42. * *SCIRPUS fetaceus* culmo nudo fetaceo : spicis la-
teralibus subsolitariis sessilibus. *Morif. Hist. t. 10.*
f. 23.

Ad piscinas ac in ripa lacus *Lyngby.*

* * * *Culmo triquetro, panicula foliacea.*

43. *SCIRPUS maritimus* culmo triquetro : panicula
conglobata foliacea, spicularum squamis trifidis :
intermedia subulata.

In littore Maris Balthici.

44. *SCIRPUS sylvaticus* culmo triquetro folioso : um-
bella foliosa : pedunculis nudis supradecompositis :
spicis confertis.

Dan. *Cyper-Gras.*

In paludosis ad *Valbulla* vulgaris.

17. ERYOPHORUM. 68.

45. *ERYOPHORUM Polystachion* culmo folioso tereti :
foliis planis : spicis pedunculatis. *Vaill. Bot. Paris.*
t. 16. f. 1. P. Fl. Oec. 67.

Dan. *Enguld; Kier-uld.*

In paludibus uliginosis.

46. *ERYOPHORUM vaginatum* culmis teretibus vagina-
tis : spica paleacea. *Fl. Dan. t. 236. Scheuchz.*
Agrost. t. 7. f. 1.

In paludibus.

18. NARDUS. 69.

47. NARDUS *stricta* spica recta setacea. *Morif. Hist.*
p. 217. t. 7. f. 8.
In sterilibus.

DIGYNIA.

19. PHALARIS. 74.

48. PHALARIS *arundinacea*, panicula oblonga ventricosa.
In ripa amnis, ac in prato humido nemoris copiosissima.
49. PHALARIS *phleoides* panicula cylindrica spiciformi.
In pratis siccis.

20. PANICUM. 76.

50. PANICUM *crus galli* spicis alternis conjugatisque:
spiculis subdivisis; glumis aristatis hispidis, rachis
quinquangulari. *Morif. Hist.* p. 189. t. 4. f. 15.
In cultis; passim.

21. PHLEUM. 77.

51. PHLEUM *pratense* spica cylindracea longissima.
Morif. Hist. p. 188. t. 4. f. 1.
In agris frequens.
52. PHLEUM *nodosum* spica cylindrica basi sterili: culmo
adscendente, radice bulbifera.
In aridis.
53. * PHLEUM *alpinum* spica ovato-cylindracea. *Fl.*
Dan. t. 213.
Rare.

54. † *PHLEUM arenarium* spica ovata ciliata: caule
ramoso.
In arenosis.

22. ALOPECURUS. 78.

55. ALOPECURUS *pratensis* culmo spicato erecto.
P. Fl. Oec. 15.
Dan. *Kasse-Gras*; *Revrumpe-Gras.*
In pratis fertilibus.
56. ALOPECURUS *geniculatus* culmo spicato infracto!
Dan. *Suppe-Gras.*
In sterilioribus.

23. MILIUM. 79.

57. * *MILIUM effusum* floribus paniculatis dispersis
muticis. *Moris. Hist. t. 5. f. 10.*
In Nemore & Sylva frequens.

24. AGROSTIS. 80.

* *Aristata.*

58. AGROSTIS *Spica venti* petalo exteriori aristam re-
ctam, strictam, longissimam exsertente.
Dan. *Hvine-Gras.*
In agris Secalinis.
59. AGROSTIS *rubra* paniculae parte florente patentif-
sima: petalo exteriori glabro, terminato arista tor-
tili-recurva.
In pratis.
60. AGROSTIS *canina* calycibus coloratis: petalorum
arista dorsali recurva: culmis prostratis subca-
mosis.
In pascuis humidiusculis.

* * Mutica.

61. * AGROSTIS *stolonifera* paniculae ramulis divaricatis - muticis: culmo repente: calycibus æqualibus.

In agris.

62. AGROSTIS *capillaris* panicula capillari patente: calycibus subulatis æqualibus hispidiusculis coloratis: flosculis muticis. *Fl. Dan. T. 1. t. 163.*

In pratis.

25. A I R A. 81.

* Mutica.

63. AIRA *cærulea* foliis planis: panicula coarctata: floribus pedunculatis muticis convoluto-subulatis.

Fl. Dan. t. 239. P. Fl. Oec. 10.

Dan. *Blaatoppet - Gras.*

In pascuis aquosis.

64. AIRA *aquatica* foliis planis: panicula patente: floribus muticis lævibus calyce longioribus.

Vaill. Bot. Paris. t. 17. f. 7.

In aquosis pascuis.

* * Aristata.

65. AIRA *cespitosa* foliis planis: panicula patente: petalis basi villosis aristatisque arista recta brevi.

Fl. Dan. t. 240. P. Fl. Oec. 11.

In pratis.

66. AIRA *flexuosa* foliis setaceis: culmis subnudis panicula divaricata: pedunculis flexuosis.

Fl. Dan. t. 157.

In Sylva.

67. AIRA *canescens* foliis setaceis, summo spathaceo paniculam inferne obvolvante.

Dan. *Hvas Sive - Gras.*

In arvis.

68. *AIRA precox* foliis setaceis: vaginis angulatis: floribus paniculato-spicatis: flosculis basi aristatis.
In arenosis.

26. MELICA. 82.

69. *MELICA nutans* petalis imberbibus: panicula nutante simplici.
Dan. *Havre-Gras*.
In Nemore passim.
70. † *MELICA ciliata* flosculi inferioris petalo exteriori ciliato.
In sterilibus.

27. POA. 83.

71. *POA aquatica* panicula diffusa: spiculis sexfloris linearibus. *Morif. Hist. p. 201. t. 6. f. 25.*
Dan. *Vand-Gras*.
Obs. Spiculæ communiter quinquefloræ, raro sexfloræ, sæpe quadrifloræ, interdum trifloræ; culmus humanam superans altitudinem.
In ripa stagni *Lyngby* æ æmnis.
72. *POA trivialis* panicula subdiffusa: spiculis trifloris, quadrifloris pubescentibus: culmo erecto tereti.
In pascuis ubique.
73. * *POA angustifolia* panicula diffusa: spiculis quadrifloris pubescentibus: culmo erecto tereti.
In pascuis & alibi, copiose in horto.
74. *POA pratensis* panicula diffusa: spiculis quinquefloris glabris: culmo erecto tereti.
In pratis.
75. *POA annua* panicula diffusa angulis rectis: spiculis obtusis: culmo obliquo compresso.
In pratis.

76. POA

76. *POA compressa* panicula secunda coarctata: culmo obliquo compresso. *Vaill. Bot. Paris. t. 18. f. 5.*
In sterilibus.

77. * *POA nemoralis* panicula attenuata: spiculis sub-
bifloris, mucronatis scabris: culmo incurvo.
In nemore.

28. **B R I Z A.** 84.

78. *BRIZA media* spiculis ovatis: calyce flosculis brevior. *Morif. Hist. p. 203. t. 6. f. 45.*
Dan. Befver-Gras, Kierligheds-Gras.
In pratis frequens.

29. **DACTYLIS.** 86.

79. *DACTYLIS glomerata*, panicula secunda glomerata.
Morif. Hist. p. 202. t. 6. f. 38.
Dan. Hvas-Gras.
Vulgaris.

30. **CYNOSURUS.** 87.

80. *CYNOSURUS cristatus* bracteis pinnatifidis.
Fl. Dan. t. 238.
Dan. Hane, Korns-Gras.
In pratis frequens.

† *Cynosurus panicus* panicula subspicata: flosculis simplicibus biaristatis.

Gramen hoc in *Fl. Su.* an. 1755. descriptum; altera species in *Linnei* editione an. 1762. exulat.

31. **FESTUCA.** 88.

* *Panicula secunda.*

81. *FESTUCA ovina* panicula secunda coarctata aristata: culmo tetragono nudiusculo: foliis setaceis.
P. Fl. Oec. 71. Morif. Hist. p. 182. t. 3. f. 13.
In collibus aridis frequens.

82. * *FESTUCA rubra* panicula secunda scabra: spiculis septifloris aristatis: flosculo ultimo mutico: culmo femitereti.

In sterilibus:

* * *Panicula aequali.*

83. *FESTUCA decumbens* panicula erecta: spiculis subovatis muticis: calyce flosculis majore: caule decumbente. *Fl. Dan. t. 162.*

Dan. *Heyre-Gras.*

In pascuis.

84. *FESTUCA elatior* panicula secunda erecta: spiculis subaristatis: exterioribus teretibus.

Buxb. Cent. 5. t. 16.

In pratis humidiusculis.

85. *FESTUCA fluitans* panicula ramosa erecta: spiculis subseffilibus teretibus muticis.

Fl. Dan. t. 237. P. Fl. Oec. 70.

Dan. *Maræ-Gras; Suppe-Gras.*

In paludosis.

32. B R O M U S. 89.

86. *BROMUS secalinus* panicula patente: spiculis ovatis: aristis rectis. *P. Fl. Oec. 40.*

Dan. *Heyre.*

β, *Bromus* aristis prorsus carens & glaber. *Scheuchz. Agrost. p. 251.*

α, In agris & alibi vulgaris. β, rarior in horto.

87. *BROMUS mollis* panicula erectiuscula: spicis ovatis: aristis rectis: foliis mollissime villosis.

Morif. Hist. p. 213. t. 7. f. 18.

In siccis horti.

Num satis a Secalino distinctus?

88. *BROMUS sterilis* panicula patula: spiculis oblongis distichis: glumis subulato-aristatis.

Morif. Hist. p. 212. t. 7. f. 11.

In agris ad vicum *Kollekolle*.

89. * *BROMUS arvensis* panicula nutante: spiculis ovato-oblongis.

In agris.

90. *BROMUS tectorum* panicula nutante spiculis linearibus.

In sylvestribus pratis.

91. † *BROMUS giganteus* panicula nutante: spiculis quadrifloris: aristis brevioribus.

Vaill. Bot. Paris. t. 18. f. 3.

Ad *Gyldenlund*.

92. *BROMUS pinnatus* spiculis alternis subsessilibus teretibus: culmo indiviso. *Fl. Dan. t. 164.*

In sterilibus haud infrequens.

93. † *BROMUS triflorus* panicula patente: floribus subtrifloris.

In nemoribus.

33. A V E N A. 91.

* *Paniculata.*

94. *AVENA elatior* paniculata, calycibus bifloris: flosculo hermaphrodito mutico: masculo aristato.

Fl. Dan. t. 165.

In pratis haud infrequens, ac in paludibus.

95. *AVENA fatua* paniculata, calycibus trifloris: flosculo

in omnibus basi pilosis: aristis totis levibus.

P. Fl. Oec. 34. 96.

Dan. Wild Havre; Flare. Havre.

Inter segetes.

MONOGYNIA.

12. VALERIANA. 44.

29. VALERIANA *officinalis* floribus triandris: foliis omnibus pinnatis. *P. Fl. Oec.* 153.

Dan. *Oyen-rod*; BALDRIAN.

Pharmac. VALERIANÆ radix, herba.

In paludosis nemoris.

30. VALERIANA *Dioica* floribus triandris dioicis: foliis pinnatis.

In pratis uliginosis.

31. VALERIANA *locusta* caule dichotomo: foliis linearibus: floribus triandris. *P. Fl. Oec.* 152.

Dan. *Fettekaas*; *Feldkrop*.

In arvis.

13. IRIS. 59.

* *Barbata nectariis petalorum reflexorum.*

32. † IRIS *Germanica* corollis barbatis: caule-foliis longiore multifloro: floribus inferioribus pedunculatis.

In Insula *Saltholm*.

* * *Imberbes petalis deflexis levibus.*

33. IRIS *pseudacorus* corollis imberbibus: petalis interioribus: stigmatibus minoribus: foliis ensiformibus.

Dan. *Sverd-Lilie*.

Pharmac. ACORI PALUSTRIS radix.

In paludosis.

14. SCHOENUS. 65.

34. * SCHOENUS *Mariscus* culmo tereti, foliis marginate dorsoque aculeatis. *Morif. Hist.* p. 273. t. II. f. 24.

In lacu *Lyngby* copiose.

- 35 * *SCHOENUS albus* culmo subtriquetro folioso;
 floribus fasciculatis: foliis setaceis. *Morif. Hist.*
t. 9. f. 39.
 In paludosis.

15. CYPERUS. 66.

36. † *CYPERUS fuscus* culmo triquetro nudo: umbella
 trifida: pedunculis simplicibus inæqualibus: spi-
 cis confertis linearibus. *Fl. Dan. t. 179.*
 In pratis paludosis.
 Flora Danica huic assignat locum natalem *Fridrichsdal*; ibi-
 dem hucusque invenire nequeo.

16. SCIRPUS. 67.

* *Spica unica.*

37. *SCIRPUS palustris* culmo tereti nudo: spica sub-
 vata terminali. *Fl. Dan. t. 167.*
Dan. Sif. nomen generis.
Obs. Figura, *t. 167.* quam *Scirpum acicularem* vult
Cl. Oederus, potius offert palustrem; acicularis enim duplo
 vel triplo minor, tenuior &c. culmo plerumque curvato.
 In ripis ac inundatis frequens.
38. *SCIRPUS cespitosus* culmo striato nudo: spica bi-
 valvi terminali longitudine calycis: radicibus
 squamula interstinctis. *Scheuchz. Agrost. t. 7. f. 18.*
P. Fl. Oec. 132.
Dan. Myre-Sif.
 In paludibus turfosis.
39. *SCIRPUS acicularis* culmo tereti nudo setiformi;
 spica ovata terminali bivalvi: seminibus nudis.
Scheuchz. Agrostog. t. 7. f. 19. Morif. Hist. t. 10.
f. 37.
 Sub aqua piscinæ in Sylva *Hareskov* copiose.

40. † SCIRPUS *fluitans* culmis teretibus nudis alternis: caule folioso flaccido. *Scheuchz. Agrost. t. 7. f. 20.*
In inundatis.

* * *Culmo tereti polystachio.*

41. SCIRPUS *lacustris* culmo tereti nudo: spicis ovatis pluribus pedunculatis terminalibus.

In stagnis ubique.

42. * SCIRPUS *setaceus* culmo nudo setaceo: spicis lateralibus subsolitariis sessilibus. *Morif. Hist. t. 10. f. 23.*

Ad piscinas ac in ripa lacus *Lyngby.*

* * * *Culmo triquetro, panicula foliacea.*

43. SCIRPUS *maritimus* culmo triquetro: panicula conglobata foliacea, spicularum squamis trifidis: intermedia subulata.

In littore Maris Balthici.

44. SCIRPUS *sylvaticus* culmo triquetro folioso: umbella foliosa: pedunculis nudis supradecompositis: spicis confertis.

Dan. *Cyper-Gras.*

In paludosis ad *Valballe* vulgaris.

17. ERYOPHORUM. 68.

45. ERIOPHORUM *Polystachion* culmo folioso tereti: foliis planis: spicis pedunculatis. *Vaill. Bot. Paris. t. 16. f. 1. P. Fl. Oec. 67.*

Dan. *Enguld; Kier-uld.*

In paludibus uliginosis.

46. ERIOPHORUM *vaginatum* culmis teretibus vaginatis: spica paleacea. *Fl. Dan. t. 236. Scheuchz. Agrost. t. 7. f. 1.*

In paludibus.

18. NARDUS. 69.

47. NARDUS *stricta* spica recta setacea. *Morif. Hist.*
p. 217. t. 7. f. 8.
In sterilibus.

DIGYNIA.

19. PHALARIS. 74.

48. PHALARIS *arundinacea*, panicula oblonga ventricosa.
In ripa amnis, ac in prato humido nemoris copiosissima.
49. PHALARIS *phleoides* panicula cylindrica spiciformi.
In pratis siccis.

20. PANICUM. 76.

50. PANICUM *crus galli* spicis alternis conjugatisque:
spiculis subdivisis: glumis aristatis hispidis, rachi
quinquangulari. *Morif. Hist.* p. 189. t. 4. f. 15.
In cultis; passim.

21. PHLEUM. 77.

51. PHLEUM *pratense* spica cylindracea longissima
Morif. Hist. p. 188. t. 4. f. 1.
In agris frequens.
52. PHLEUM *nodosum* spica cylindrica basi sterili: culmo
adscendente, radice bulbifera.
In aridis.
53. * PHLEUM *alpinum* spica ovato-cylindracea. *Fl.*
Dan. t. 213.
Raro.

54. † *PHLEUM arenarium* spica ovata ciliata: caule
ramoso.
In arenosis.

22. ALOPECURUS. 78.

55. ALOPECURUS *pratensis* culmo spicato erecto.
P. Fl. Oec. 15.
Dan. *Kasle - Gras*; *Revrumpe - Gras*.
In pratis fertilibus.
56. ALOPECURUS *geniculatus* culmo spicato infracto.
Dan. *Suppe - Gras*.
In sterilioribus.

23. MILIUM. 79.

57. * MILIUM *effusum* floribus paniculatis dispersis
muticis. *Moris. Hist. t. 5. f. 10.*
In Nemore & Sylva frequens.

24. AGROSTIS. 80.

- * *Aristata.*

58. AGROSTIS *Spica venti* petalo exteriori aristam rectam, strictam, longissimam exsertente.
Dan. *Hvine - Gras*.
In agris Secalinis.
59. AGROSTIS *rubra* paniculae parte florente patentissima; petalo exteriori glabro, terminato arista tortili-recurva.
In pratis.
60. AGROSTIS *canina* calycibus coloratis: petalorum arista dorsali recurva: culmis prostratis subramosis.
In pascuis humidiusculis.

* * Mutica.

61. * AGROSTIS *stolonifera* panicula ramulis divaricatis - muticis: culmo repente: calycibus æqualibus.

In agris.

62. AGROSTIS *capillaris* panicula capillari patente: calycibus subulatis æqualibus hispidiusculis coloratis: flosculis muticis. *Fl. Dan. T. 1. t. 163.*

In pratis.

25. A I R A. 81.

* Mutica.

63. AIRA *cærulea* foliis planis: panicula coarctata: floribus pedunculatis muticis convoluto-subulatis.

Fl. Dan. t. 239. P. Fl. Oec. 10.

Dan. *Blaatoppet - Græs.*

In pascuis aquosis.

64. AIRA *aquatica* foliis planis: panicula patente: floribus muticis lævibus calyce longioribus.

Vaill. Bot. Paris. t. 17. f. 7.

In aquosis pascuis.

* * Aristata.

65. AIRA *cespitosa* foliis planis: panicula patente: petalis basi villosis aristatisque arista recta brevi.

Fl. Dan. t. 240. P. Fl. Oec. 11.

In pratis.

66. AIRA *flexuosa* foliis setaceis: culmis subnudis panicula divaricata: pedunculis flexuosis.

Fl. Dan. t. 157.

In Sylva.

67. AIRA *canescens* foliis setaceis, summo spathaceo paniculam inferne obvolvante.

Dan. *Hvas Sive - Græs.*

In arvis.

68. *AIRA precox* foliis fetaceis: vaginis angulatis: floribus paniculato-spicatis: flosculis basi aristatis.
In arenosis.

26. MELICA. 82.

69. *MELICA nutans* petalis imberbibus: panicula nutante simplici.
Dan. *Havre-Gras*.
In Nemore passim.
70. † *MELICA ciliata* flosculi inferioris petalo exteriori ciliato.
In sterilibus.

27. POA. 83.

71. *POA aquatica* panicula diffusa: spiculis sexfloris linearibus. *Morif. Hist. p. 201. t. 6. f. 25*.
Dan. *Vand-Gras*.
Obs. Spiculæ communiter quinquefloræ, raro sexfloræ, sæpe quadrifloræ, interdum trifloræ; culmus humanam superans altitudinem.
In ripa stagni *Lyngby* æe amnis.
72. *POA trivialis* panicula subdiffusa: spiculis trifloris, quadrifloris pubescentibus: culmo erecto tereti.
In pascuis ubique.
73. * *POA angustifolia* panicula diffusa: spiculis quadrifloris pubescentibus: culmo erecto tereti.
In pascuis & alibi, copiose in horto.
74. *POA pratensis* panicula diffusa: spiculis quinquefloris glabris: culmo erecto tereti.
In pratis.
75. *POA annua* panicula diffusa angulis rectis: spiculis obtusis: culmo obliquo compresso.
In pratis.

76. POA compressa panicula secunda coarctata: culmo obliquo compresso. *Vaill. Bot. Paris. t. 18. f. 5.*
In sterilibus.

77. * POA nemoralis panicula attenuata: spiculis sub-
bifloris, mucronatis scabris: culmo incurvo.
In nemore.

28. BRIZA. 84.

78. BRIZA media spiculis ovatis: calyce flosculis brevior. *Morif. Hist. p. 203. t. 6. f. 45.*
Dan. Befver-Gras, Kiarligheds-Gras.
In pratis frequens.

29. DACTYLIS. 86.

79. DACTYLIS glomerata, panicula secunda glomerata.
Morif. Hist. p. 202. t. 6. f. 38.
Dan. Heas-Gras.
Vulgaris.

30. CYNOSURUS. 87.

80. CYNOSURUS cristatus bracteis pinnatifidis.
Fl. Dan. t. 238.
Dan. Hane, Korns-Gras.
In pratis frequens.

† Cynosurus paniceus panicula subspicata: flosculis simplicibus biaristatis.

Gramen hoc in *Fl. Su.* an. 1755. descriptum, altera species in *Linnaei* editione an. 1762. exulat.

31. FESTUCA. 88.

* Panicula secunda.

81. FESTUCA ovina panicula secunda coarctata aristata: culmo tetragono nudiusculo: foliis setaceis.
P. Fl. Oec. 71. Morif. Hist. p. 182. t. 3. f. 13.
In collibus aridis frequens.

82. * *FESTUCA rubra* panicula secunda scabra: spiculis
septifloris aristatis: flosculo ultimo mutico: culmo
semitereti.
In sterilibus:

* * *Panicula equali.*

83. *FESTUCA decumbens* panicula erecta: spiculis sub-
ovatis muticis: calyce flosculis majore: caule de-
cumbente. *Fl. Dan. t. 162.*
Dan. *Heyre-Gras.*
In pascuis.
84. *FESTUCA elatior* panicula secunda erecta: spiculis
subaristatis: exterioribus teretibus.
Buxh. Cent. 5. t. 16.
In pratis humidiusculis.
85. *FESTUCA fluitans* panicula ramosa erecta: spiculis
subseffilibus teretibus muticis.
Fl. Dan. t. 237. P. Fl. Oec. 70.
Dan. *Manne-Gras; Suppe-Gras.*
In paludosis.

32. B R O M U S. 89.

86. *BROMUS secalinus* panicula patente: spiculis ova-
tis: aristis rectis. *P. Fl. Oec. 40.*
Dan. *Heyre.*
 β , *Bromus* aristis prorsus carens & glaber. *Scheuchz.*
Agrost. p. 251.
 α , In agris & alibi vulgaris. β , rarior in horto.
87. *BROMUS mollis* panicula erectiuscula: spicis ovatis:
aristis rectis: foliis mollissime villosis.
Morif. Hist. p. 213. t. 7. f. 18.
In siccis hortis.
Num satis a *Secalino* distinctus?

88. *BROMUS sterilis* panicula patula: spiculis oblongis distichis; glumis subulato-aristatis.

Morif. Hist. p. 212. t. 7. f. 11.

In agris ad vicum *Kollekolle*.

89. * *BROMUS arvensis* panicula nutante: spiculis ovato-oblongis.

In agris.

90. *BROMUS tectorum* panicula nutante spiculis linearibus.

In sylvestribus pratis.

91. † *BROMUS giganteus* panicula nutante: spiculis quadrifloris: aristis brevioribus.

Vaill. Bot. Paris. t. 18. f. 3.

Ad *Goldenlund*.

92. *BROMUS pinnatus* spiculis alternis subsessilibus teretibus: culmo indiviso. *Fl. Dan. t. 164.*

In sterilibus haud infrequens.

93. † *BROMUS triflorus* panicula patente: floribus subtrifloris.

In nemoribus.

33. A V E N A. 91.

* *Paniculata.*

94. *AVENA elatior* paniculata, calycibus bifloris: flosculo hermaphrodito mutico: masculo aristato.

Fl. Dan. t. 165.

In pratis haud infrequens, ac in paludibus.

95. *AVENA fatua* paniculata, calycibus trifloris: flosculo hermaphrodito mutico: aristis totis levibus.

P. Fl. Oec. 34. 96.

Dan. Wild Havre; Fl. Havre.

Inter segetes.

MONOGYNIA.

12. VALERIANA. 44.

29. VALERIANA *officinalis* floribus triandris: foliis omnibus pinnatis. *P. Fl. Oec.* 153.

Dan. *Oyen-rod*; BALDRIAN.

Pharmac. VALERIANÆ radix, herba.

In paludosis nemoris.

30. VALERIANA *Dioica* floribus triandris dioicis: foliis pinnatis.

In pratis uliginosis.

31. VALERIANA *locusta* caule dichotomo: foliis linearibus: floribus triandris. *P. Fl. Oec.* 152.

Dan. *Fettekaas*; *Feldkrop*.

In arvis.

13. IRIS. 59.

* *Barbata nectariis petalorum reflexorum.*

32. † IRIS *Germanica* corollis barbatis: caule-foliis longiore multifloro: floribus inferioribus pedunculatis.

In Insula *Saltholm*.

* * *Imberbes petalis deflexis levibus.*

33. IRIS *pseudacorus* corollis imberbibus: petalis interioribus stigmatibus minoribus: foliis ensiformibus.

Dan. *Sverd-Lilie*.

Pharmac. ACORI PALUSTRIS radix.

In paludosis.

14. SCHOENUS. 65.

34. * SCHOENUS *Mariscus* culmo tereti, foliis marginate dorsoque aculeatis. *Morif. Hist.* p. 273.

t. II. f. 24.

In lacu *Lyngby* copiose.

35. * *SCHOENUS albus* culmo subtriquetro folioso, floribus fasciculatis: foliis setaceis. *Morif. Hist. t. 9. f. 39.*
In paludosis.

15. CYPERUS. 66.

36. † *CYPERUS fuscus* culmo triquetro nudo: umbella trifida: pedunculis simplicibus inæqualibus: spicis confertis linearibus. *Fl. Dan. t. 179.*
In pratis paludosis.

Flora Danica huic assignat locum natalem *Fridrichsdal*; ibidem hucusque invenire nequeo.

16. SCIRPUS. 67.

* *Spica unica.*

37. *SCIRPUS palustris* culmo tereti nudo: spica subovata terminali. *Fl. Dan. t. 167.*

Dan. *Sif.* nomen generis.

Obs. Figura, t. 167. quam *Scirpum acicularem* vult *Cl. Oederus*, potius offert *palustrem*; acicularis enim duplo vel triplo minor, tenuior &c. culmo plerumque curvato.

In ripis ac inundatis frequens.

38. *SCIRPUS cespitosus* culmo striato nudo: spica bivalvi terminali longitudine calycis: radicibus squamula interstinctis. *Scheuchz. Agrost. t. 7. f. 18. P. Fl. Oec. 132.*

Dan. *Myre-Sif.*

In paludibus turfosis.

39. *SCIRPUS acicularis* culmo tereti nudo setiformi: spica ovata terminali bivalvi: seminibus nudis. *Scheuchz. Agrostog. t. 7. f. 19. Morif. Hist. t. 10. f. 37.*

Sub aqua piscinæ in Sylva *Hareskov* copiose.

40. † SCIRPUS *fluitans* culmis teretibus nudis alternis :
caule folioso flaccido. *Scheuchz. Agrost. t. 7. f. 20.*
In inundatis.

* * *Culmo tereti polystachio.*

41. SCIRPUS *lacustris* culmo tereti nudo : spicis ovatis
pluribus pedunculatis terminalibus.

In stagnis ubique.

42. * SCIRPUS *setaceus* culmo nudo setaceo : spicis la-
teralibus subfolitariis sessilibus. *Morif. Hist. t. 10.*
f. 23.

Ad piscinas ac in ripa lacus *Lyngby.*

* * * *Culmo triquetro, panicula foliacea.*

43. SCIRPUS *maritimus* culmo triquetro : panicula
conglobata foliacea, spicularum squamis trifidis :
intermedia subulata.

In littore Maris Balthici.

44. SCIRPUS *ylvaticus* culmo triquetro folioso : um-
bella foliosa : pedunculis nudis supradecompositis :
spicis confertis.

Dan. *Cyper-Gras.*

In paludosis ad *Valbulla* vulgaris.

17. ERYOPHORUM. 68.

45. ERIOPHORUM *Polystachion* culmo folioso tereti :
foliis planis : spicis pedunculatis. *Vaill. Bot. Paris.*
t. 16. f. 1. P. Fl. Oec. 67.

Dan. *Enguld; Kier-uld.*

In paludibus uliginosis.

46. ERIOPHORUM *vaginatum* culmis teretibus vagina-
tis : spica paleacea. *Fl. Dan. t. 236. Scheuchz.*
Agrost. t. 7. f. 1.

In paludibus.

18. NARDUS. 69.

47. NARDUS *stricta* spica recta setacea. *Morif. Hist.*
p. 217. t. 7. f. 8.
In sterilibus.

DIGYNIA.

19. PHALARIS. 74.

48. PHALARIS *arundinacea*, panicula oblonga ventricosa.
In ripa amnis, ac in prato humido nemoris copiosissima.
49. PHALARIS *phleoides* panicula cylindrica spiciformi.
In pratis siccis.

20. PANICUM. 76.

50. PANICUM *crus galli* spicis alternis conjugatisque:
spiculis subdivisis; glumis aristatis hispidis, rachis
quinquangulari. *Morif. Hist.* p. 189. t. 4. f. 15.
In cultis; passim.

21. PHEUM. 77.

51. PHEUM *pratense* spica cylindracea longissima.
Morif. Hist. p. 188. t. 4. f. 1.
In agris frequens.
52. PHEUM *nodosum* spica cylindrica basi sterili: culmo
adscendente, radice bulbifera.
In aridis.
53. * PHEUM *alpinum* spica ovato-cylindracea. *Fl.*
Dan. t. 213.
Raro.

54. † *PHLEUM arenarium* spica ovata ciliata: caule ramofo.
In arenosis.

22. ALOPECURUS. 78.

55. ALOPECURUS *pratensis* culmo spicato erecto.
P. Fl. Oec. 15.
Dan. *Kasle-Gras*; *Revrumpe-Gras*.
In pratis fertilibus.
56. ALOPECURUS *geniculatus* culmo spicato infracto.
Dan. *Suppe-Gras*.
In sterilioribus.

23. MILIUM. 79.

57. * *MILIUM effusum* floribus paniculatis dispersis muticis. *Moris. Hist. t. 5. f. 10.*
In Nemore & Sylva frequens.

24. AGROSTIS. 80.

- * *Aristata.*
58. AGROSTIS *Spica venti* petalo exteriori aristam rectam, strictam, longissimam existentem.
Dan. *Hvine-Gras*.
In agris Secalinis.
59. AGROSTIS *rubra* paniculae parte florente patentissima: petalo exteriori glabro, terminato arista tortili-recurva.
In pratis.
60. AGROSTIS *canina* calycibus coloratis: petalorum arista dorsali recurva: culmis prostratis subramosis.
In pascuis humidiusculis.

* * Mutica.

61. * AGROSTIS *stolonifera* paniculae ramulis divaricatis - muticis: culmo repente: calycibus æqualibus.

In agris.

62. AGROSTIS *capillaris* panicula capillari patente: calycibus subulatis æqualibus hispidiusculis coloratis: flosculis muticis. *Fl. Dan. T. 1. t. 162.*

In pratis.

25. A I R A. 81.

* Mutica.

63. AIRA *cærulea* foliis planis: panicula coarctata: floribus pedunculatis muticis convoluto-subulatis. *Fl. Dan. t. 239. P. Fl. Oec. 10.*

Dan. *Blaatoppet - Græs.*

In pascuis aquosis.

64. AIRA *aquatica* foliis planis: panicula patente: floribus muticis lævibus calyce longioribus.

Vaill. Bot. Parif. t. 17. f. 7.

In aquosis pascuis.

* * Aristata.

65. AIRA *cespitosa* foliis planis: panicula patente: petalis basi villosis aristatisque arista recta brevi. *Fl. Dan. t. 240. P. Fl. Oec. 11.*

In pratis.

66. AIRA *flexuosa* foliis setaceis: culmis subnudis panicula divaricata: pedunculis flexuosis.

Fl. Dan. t. 157.

In Sylva.

67. AIRA *canescens* foliis setaceis, summo spathaceo paniculam inferne obvolvante.

Dan. *Hvas Sive - Græs.*

In arvis.

68. *AIRA precox* foliis fetaceis: vaginis angulatis: floribus paniculato-spicatis: flosculis basi aristatis.
In arenosis.

26. MELICA. 82.

69. *MELICA nutans* petalis imberbibus: panicula nutante simplici.
Dan. *Havre-Gras*.
In Nemore passim.
70. † *MELICA ciliata* flosculi inferioris petalo exteriori ciliato.
In sterilibus.

27. POA. 83.

71. *POA aquatica* panicula diffusa: spiculis sexfloris linearibus. *Morif. Hist. p. 201. t. 6. f. 25*.
Dan. *Vand-Gras*.
Obs. Spiculæ communiter quinquefloræ, raro sexfloræ, sæpe quadrifloræ, interdum trifloræ; culmus humanam superans altitudinem.
In ripa stagni *Lyngby* æ æmnis.
72. *POA trivialis* panicula subdiffusa: spiculis trifloris, quadrifloris pubescentibus: culmo erecto tereti.
In pascuis ubique.
73. * *POA angustifolia* panicula diffusa: spiculis quadrifloris pubescentibus: culmo erecto tereti.
In pascuis & alibi, copiose in horto.
74. *POA pratensis* panicula diffusa: spiculis quinquefloris glabris: culmo erecto tereti.
In pratis.
75. *POA annua* panicula diffusa angulis rectis: spiculis obtusis: culmo oblique compresso.
In pratis.

76. POA compressa panicula secunda coarctata: culmo obliquo compresso. *Vaill. Bot. Paris. t. 18. f. 5.*
In sterilibus.

77. * POA nemoralis panicula attenuata: spiculis sub-
bifloris, mucronatis scabris: culmo incurvo.
In nemore.

28. BRIZA 84.

78. BRIZA media spiculis ovatis: calyce floribus bre-
vior. *Morif. Hist. p. 203. t. 6. f. 45.*
Dan. Befver-Gras, Kiarligheds-Gras.
In pratis frequens.

29. DACTYLIS 86.

79. DACTYLIS glomerata, panicula secunda glomerata.
Morif. Hist. p. 202. t. 6. f. 38.
Dan. Heas-Gras.
Vulgaris.

30. CYNOSURUS 87.

80. CYNOSURUS cristatus bracteis pinnatifidis.
Fl. Dan. t. 238.
Dan. Hane, Korns-Gras.
In pratis frequens.

† Cynosurus paniceus panicula subspicata: fasciculis
simplicibus biaristatis.

Gramen hoc in *Fl. Su.* an. 1755. descriptum, altera spe-
ciem *Linnei* editione an. 1762. exulat.

31. FESTUCA 88.

* Panicula secunda.

81. FESTUCA ovina panicula secunda coarctata aristata;
culmo tetragono nudiusculo: foliis setaceis.
P. Fl. Oec. 71. Morif. Hist. p. 182. t. 3. f. 13.
In collibus aridis frequens.

82. * *FESTUCA rubra* panicula secunda scabra: spiculis septifloris aristatis: flosculo ultimo mutico: culmo semitereti.

In sterilibus.

* * *Panicula aequali.*

83. *FESTUCA decumbens* panicula erecta: spiculis subovatis muticis: calyce flosculis majore: caule decumbente. *Fl. Dan. t. 162.*

Dan. *Heyre-Gras.*

In pascuis.

84. *FESTUCA elatior* panicula secunda erecta: spiculis subaristatis: exterioribus teretibus.

Buxb. Cent. 5. t. 16.

In pratis humidiusculis.

85. *FESTUCA fluitans* panicula ramosa erecta: spiculis subseffilibus teretibus muticis.

Fl. Dan. t. 237. P. Fl. Oec. 70.

Dan. *Manae-Gras; Suppe-Gras.*

In paludosis.

32. B R O M U S. 89.

86. *BROMUS secalinus* panicula patente: spiculis ovatis: aristis rectis. *P. Fl. Oec. 40.*

Dan. *Heyre.*

β, *Bromus* aristis prorsus carens & glaber. *Scheuchz.*

Agrost. p. 251.

α, In agris & alibi vulgaris. β, rarior in horto.

87. *BROMUS mollis* panicula erectiuscula: spicis ovatis: aristis rectis: foliis mollissime villosis.

Morif. Hist. p. 213. t. 7. f. 18.

In siccis horti.

Num satis a *Secalino* distinctus?

88. *BROMUS sterilis* panicula patula: spiculis oblongis distichis; glumis subulato-aristatis.

Morif. Hist. p. 212. t. 7. f. 11.

In agris ad vicum *Kollekolle*.

89. * *BROMUS arvensis* panicula nutante: spiculis ovato-oblongis.

In agris.

90. *BROMUS tectorum* panicula nutante spiculis linearibus.

In sylvestribus pratis.

91. † *BROMUS giganteus* panicula nutante: spiculis quadrifloris: aristis brevioribus.

Vaill. Bot. Paris. t. 18. f. 3.

Ad *Goldenlund*.

92. *BROMUS pinnatus* spiculis alternis subsessilibus teretibus: culmo indiviso. *Fl. Dan. t. 164.*

In sterilibus haud infrequens.

93. † *BROMUS triflorus* panicula patente: floribus subtrifloris.

In nemoribus.

33. A V E N A. 91.

* *Paniculata.*

94. *AVENA elatior* paniculata, calycibus bifloris: flosculo hermaphrodito mutico: masculo aristato.

Fl. Dan. t. 165.

In pratis haud infrequens, ac in paludibus.

95. *AVENA fatua* paniculata, calycibus trifloris: flosculis omnibus basi pilosis: aristis totis levibus.

P. Fl. Oec. 34. 96.

Dan. Vild Havre; Flare. Havre.

Inter segetes.

96. * *AVENA flavescens* panicula laxa, calycibus trifloris brevibus: flosculis omnibus aristatis.

Morif. Hist. p. 215. t. 7. f. 42.

In horto frequens.

Spicula in nostra triflora & biflora.

Avena paniculata calycibus bifloris: flosculis dorso biaristatis. Num fatuae varietas?

* * *Spicata.*

97. *AVENA pratensis* spicata: calycibus quinquefloris!
In pratis apricis.

34. A R U N D O. 93.

98. *ARUNDO phragmites* calycibus quinquefloris: panicula laxa. *P. Fl. Oec. 29.*

Dan. *Rör.*

In lacubus.

99. *ARUNDO epigeios* calycibus unifloris: panicula erecta: foliis subtus glabris.

In paludibus.

100. *ARUNDO calamagrostis* calycibus unifloris: culmo ramoso. *Schaechz. Agrest. t. 3. f. 3. B.*

In paludosis lacus *Lyngby.*

101. *ARUNDO arenaria* calycibus unifloris: foliis involutis mucronato-pungentibus. *P. Fl. Oec. 28.*

Dan. *Klittetag; Hiehne; Seene-Gras.*

In elevatis arenosis lacus *Fuhsöe.*

35. L O L I U M. 95.

102. *LOLIUM temulentum* spica aristata: spiculis compressis aristatis.

Fl. Dan. t. 160. P. Fl. Oec. 24.

Dan. *Svingel.*

In agris.

103. *LOLIUM perenne* spica mutica: spiculis compressis multifloris.

Ad vias.

104. * *LOLIUM tenue* spica mutica tereti: spiculis trifloris.

In arvis.

36. ELYMUS. 96.

105. † *ELYMUS arenarius* spica erecta arcta: calycibus tomentosis flosculo longioribus. *P. Fl. Oec.* 62.

Dan. *Spids Strand - Græs.*

In arenosis maritimis.

37. HORDEUM. 98.

106. *HORDEUM murinum* flosculis lateralibus masculis aristatis: involucris intermediis ciliatis.

Vaill. Bot. Paris. t. 17. f. 6.

Dan. *Byg - Græs; Fandens - Ax.*

In ruderatis.

38. TRITICUM. 99.

107. *TRITICUM repens* calycibus subulatis quadrifloris acuminatis. *P. Fl. Oec.* 147.

Dan. *Hunde - Græs; Knoppe - Græs.*

Pharmac. GRAMINIS radix.

In cultis.

Obs. Valde variat calycibus trifloris, quadrifloris, quinquefloris, sexfloris acuminatis; addetur alia varietas spiculis sexfloris subaristatis.

TRIGYNIA.

39. MONTIA. 101.

108. *MONTIA fontana.*

Fl. Dan. t. 131. Mich. gen. t. 13. f. 2.

In inundatis frequens.

Classis IV.

TETRANDRIA.
MONOGYNIA.

* Flores *monopetali, monospermi, superi, aggregati.*

40. DIPSACUS. *Calyx communis polyphyllus. Receptaculum conicum paleaceum. Sem. columnaria.*
 41. SCABIOSA. *Cal. communis polyphyllus, proprius, duplex. Recept. elevatum, subfoliaceum. Sem. coronata, involuta.*

* * Flores *monopetali, monocarpii, inferi.*

46. CENTUNCULUS. *Cal. 4-fidus. Cor. 4-fida, patens. Stam. brevia. Caps. 1-locularis, circumscissa.*
 45. PLANTAGO. *Cal. 4-fidus. Cor. 4-fida, limbo reflexo. Stam. longissima. Caps. 2-locularis, circumscissa.*

† *Verbena officinalis.*

* * * Flores *monopetali, dicocci, superi. Stellata.*

44. GALIUM. *Cor. 1-petala, plana. Fructus subglobosi.*
 43. ASPERULA. *Cor. 1-petala, tubulosa. Fruct. subglobosi.*
 42. SHERARDIA. *Cor. 1-petala, tubulosa. Fruct. coronatus, Semin. 3-dentatis.*

* * * * Flores *tetrapetali*, *superi*.

48. CORNUS. *Cal.* 4-dentatus, deciduus. *Drupa*
nucleo 2-loculari.

† Flores *tetrapetali inferi*.

Casdamine hirsuta.

Euonymus Europæus.

* * * * * Flores *incompleti*, *inferi*.

49. ALCHEMILLA. *Cal.* 8-fidus. *Cor.* o. *Sem.* r.
calyci inclusum.

† *Convallaria bifolia*.

D I G Y N I A.

51. CUSCUTA. *Cal.* 4-fidus. *Cor.* 1-petala, 4-fida,
ovata. *Caps.* 2-locularis, circumscissa.

50. APHANES. *Cal.* 8-fidus. *Cor.* o. *Sem.* 2. nuda.

T E T R A G Y N I A.

52. ILEX. *Cal.* 4-dentatus. *Cor.* 1-petala, rotata.
Styl. o. *Bacca* 4-sperma.

55. SAGINA. *Cal.* 4-phyllus. *Cor.* 4-petala. *Caps.*
4-locularis, 4-valvis, polysperma.

53. POTAMOGETON. *Cal.* 4-phyllus. *Cor.* o.
Styl. o. *Sem.* 4. sessilia.

54. RUPPIA. *Cal.* o. *Cor.* o. *Sem.* 4. pedicellata.
† *Linum Radiola*.



M O N O G Y N I A.

40. DIPSACUS. 114.

109. DIPSACUS *fullonum* foliis sessilibus ferratis.

Dan. *Kardé - Tidsel.*

In nemore versus hortum copiose.

110. DIPSACUS *pilosus* foliis petiolatis appendiculatis.

41. S C A B I O S A. 115.

* *Corollulis quadrifidis.*

111. SCABIOSA *arvensis* corollulis quadrifidis radiantibus: caule hispido: foliis pinnatifidis: lobis distantibus.

Dan. *Skab - Urt.*

Pharmac. SCABIOSÆ RADIX, flores.

In agris.

112. SCABIOSA *succisa* corollulis quadrifidis: caule simplici: ramis approximatis: foliis lanceolato-ovatis integerrimis.

Dan. *Dievels - Bid.*

Pharmac. MORSUS DIABOLI radix, herba.

In subhumidis.

** *Corollulis quinquefidis.*

113. SCABIOSA *columbaria* corollulis quinquefidis radiantibus: foliis radicalibus ovatis crenatis: caulibus pinnatis setaceis.

In apricis aridis.

42. S H E R A R D I A. 120.

114. SHERARDIA *arvensis* foliis omnibus verticillatis: floribus terminalibus.

In arvis.

43. ASPERULA. 121.

115. ASPERULA *adorata* foliis octonis lanceolatis: floribus fasciculatis pedunculatis. *P. Fl. Oec.* 31.

Dan. *Skov-Merke; Myssike; Bokar.*

Pharmac. MATRISYLVÆ herba.

In Sylvis passim.

44. GALIUM. 125.

* Fructu *glabro.*

116. GALIUM *palustre* foliis quaternis obovatis inaequalibus: caulibus diffusis.

Dan. *Klammer-Urt.*

In udibus.

117. GALIUM *uliginosum* foliis senis lanceolatis retrosum ferrato-aculeatis, mucronatis rigidis: corollis fructu majoribus.

In pascuis humidiusculis.

118. † GALIUM *saxatile* foliis senis obovatis obtusis: caule ramosissimo procumbente.

119. GALIUM *verum* foliis octonis linearibus sulcatis: ramis floriferis brevibus. *P. Fl. Oec.* 75.

Dan. *Klammer-Urt, Marie Sengbalm.*

In pratis frequens.

120. † GALIUM *mollugo* foliis octonis ovato-linearibus subserratis patentissimis mucronatis: caule flaccido: ramis patentibus.

* * Fructu *hispido.*

121. GALIUM *boreale* foliis quaternis lanceolatis ternerviis: caule erecto feminibus hispidis.

In sterilibus.

122. *GALIMUM Aparine* foliis octonis lanceolatis carinatis scabris retrorsum aculeatis: geniculis villosis fructu hispido. *P. Fl. Oec.* 74.

Dan. *Beyler*; *Snerre*.

Ad sepes.

45. PLANTAGO. 142.

123. *PLANTAGO major* foliis ovatis glabris: scapo tereti: spica flosculis imbricatis.

Dan. *Vey-Bred*; *Faare-Tunge*.

Pharmac. PLANTAGINIS RADIX, Folia, Semina.
Ubique in viis.

124. *PLANTAGO media* foliis ovato-lanceolatis pubescentibus: spica cylindrica: scapo tereti.

In pratis.

125. *PLANTAGO lanceolata* foliis lanceolatis: spica subovata nuda: scapo angulato.

Ad vias.

126. *PLANTAGO maritima* foliis semi-cylindræis integerrimis basi lanatis: scapo tereti.

Dan. *Strand-Krage-Tæer*; *Ravne-Fod*.

In littore maris.

Maritimam hanc plantam ad viam publicam inter *Haderstebiam* & *Apenrade* copiose crescere haud parum miratus, duomiliaria circiter a mari remotam. Uti quoque ad viam inter *Slesvicum* & *Rensburgum* passim; folia vero basi minime lanata.

127. *PLANTAGO coronopus* foliis linearibus dentatis: scapo tereti.

Dan. *Strand veybred*.

In littore maris.

128. † *PLANTAGO uniflora* scapo unifloro.

Fl. Dan. t. 170.

In inundatis.

† *Plantago dubia* foliis lanceolatis longitudine scapi: spica oblonga.

In *Fl. Su.* 1755, descripta, in altera *Specier.* ed. 1762. omissa.

46. GENTUNCULUS. 145.

129. CENTUNCULUS *minimus*.

Mich. gen. t. 18. Vaill. Bot. Paris. t. 4. f. 2.

Fl. Dan. t. 177.

In agris arenosis raro.

47. CORNUS. 149.

130. CORNUS *sanguinea* arborea: cymis nudis.

P. Fl. Oec. 56.

Dan. Vild Cornel- Tra.

Ad lacum Lyngbye- Sæ.

131. † CORNUS *Sæcica* herbacea: ramis binis.

Fl. Dan. t. 5.

Dan. Hönse- Bar.

In Mona.

48. ALCHEMILLA. 165.

132. ALCHEMILLA *vulgaris* foliis lobatis.

Pauli Fl. Oec. 13.

Dan. Lövesod.

Pharmac. ALCHEMILLÆ RADIX, herba.

In pratis sylvaticis.

DIGYNIA.

49. APHANES. 166.

133. APHANES *arvensis*.

Dan. Jomfrue- Kam.

In arvo elatiori versus pagum Virum.

50. CUSCUTA. 170.

134. † CUSCUTA *Europæa* floribus sessilibus.

Fl. Dan. t. 199.

Dan. Hör- Silke.

Pharmac. CUSCUTÆ herba; EPITHYMI herba.

In Insula Thorstinge copiose crescit.

TETRAGYNIA.

51. ILEX. 172.

135. † ILEX *aquifolium* foliis ovatis acutis spinosis.
 Dan. *Christ-Torn; Mare-Torn; Skov-Tidsel.*
 In sylvis Ducatum *Slesvici ac Holsatiae* frequens.

52. POTAMOGETON. 174.

136. POTAMOGETON *natans* foliis oblongo-ovatis petiolatis natantibus.
 In lacu *Fuursee*.
137. POTAMOGETON *perfoliatum* foliis cordatis amplexicaulibus. *Fl. Dan. t. 196.*
 Ibidem.
138. POTAMOGETON *lucens* foliis lanceolatis planis in petiolos desinentibus. *Fl. Dan. t. 195.*
139. POTAMOGETON *compressum* foliis linearibus obtusis: caule compresso. *Fl. Dan. t. 203.*
 In fossa ad Molendinum.
140. POTAMOGETON *pectinatum* foliis setaceis parallelis approximatis distichis.
 In paludibus ad lacum *Bagvard*.
141. POTAMOGETON *pusillum* foliis linearibus oppositis alternisque distantibus, basi patentibus: caule tereti. *Poill. Bot. Paris. t. 32. f. 4.*
 In fossa ad Molendinum.
142. † POTAMOGETON *crispum* foliis lanceolatis alternis undulatis ferratis.
143. † POTAMOGETON *gramineum*, foliis lineari-lanceolatis alternis sessilibus stipula latioribus.
Fl. Dan. t. 222.
 In *Bornholmia*.

144. † POTAMOGETON *marinum* foliis linearibus alternis distinctis inferne vaginantibus.

Fl. Dan. t. 186.

54. RUPPIA. 175.

145. † RUPPIA *maritima*.

Mich. gen. t. 35.

In littoribus marinis.

55. SAGINA. 176.

146. SAGINA *procumbens*, ramis procumbentibus petalis brevissimis.

In agris.



Classis V.

PENTANDRIA.

MONOGYNIA.

Flores monopetali, inferi, tetraspermi. Asperifolia.

64. ECHIUM. Cor. fauce nuda, irregularis, campanulata.
60. PULMONARIA. Cor. fauce nuda, pervia, infundibulif. Cal. prismaticeo-5-gonum.
57. LITHOSPERMUM. Cor. fauce nuda, perforata, infundibulif. Cal. 5-partitus.
61. SYMPHYTUM. Cor. fauce clausa radiis subulatis: limbo tubulato-ventricoso.
63. LYCOPSIS. Cor. fauce fornicata infundibulif.: tubo curvato.
62. ASPERUGO. Cor. fauce fornicata infundibulif.: Cal. fructus compresso-palmatus, lamellis plano-parallelis, sinuatis.
59. CYNOGLOSSUM. Cor. fauce clausa fornicibus, infundibulif. Semina depressa interiore tantum latere stylo affixa.
58. ANCHUSA. Cor. fauce clausa fornicibus, infundibulif.: tubo basi prismaticeo.
56. MYOSOTIS. Cor. fauce clausa fornicibus, hypocraterif. 5-fida: lobis emarginatis.


Flores monopetali, inferi, angiospermi.

59. ANAGALLIS. Caps. 1 - locularis, circumscissa. Cor. rotata. Stigm. capitatum.
68. LYSIMACHIA. Caps. 1 - locularis, 10-valvis, globosa, mucronata. Cor. rotata. Stig. obtusum,

65. PRIMULA. *Capsf.* 1 - locularis. *Cor.* infundibulif. fauce pervia. *Stigm.* globosum. *Involucrum* umbellulæ.
67. HOTTONIA. *Capsf.* 1 - locularis. *Cor.* hypocraterif. *Stamina* tubo corollæ imposita. *Stigm.* globosum.
66. MENYANTHES. *Capsf.* 1 - locularis. *Corolla* villosa. *Stig.* 2 - fidum.
70. CONVOLVULUS. *Capsf.* 2 - locularis : loculis dispermis. *Cor.* campanulata, plicata. *Stig.* bifidum.
76. DATURA. *Capsf.* 2 - locularis, 4 - valvis. *Cor.* infundibulif. plicata. *Cal.* tubulosus, angulatus, deciduus.
77. HYOSCYAMUS. *Capsf.* 2 - locularis, operculata. *Cor.* infundibulif. obtusa. *Stamina* inclinata. *Stig.* capitatum.
75. VERBASCUM. *Capsf.* 2 - locularis, 2 - valvis. *Cor.* rotata, subinæqualis. *Stigm.* obtusum.
78. SOLANUM. *Bacca* 2 - locularis. *Cor.* rotata. *Anthera* subcoalitæ poro gemino dehiscentes.
79. PHYSALIS. *Bacca* 2 - locularis intra calycem inflatum. *Cor.* rotata. *Anthera* conniventes.

Flores *monopetali*, *superi.*

72. SAMOLUS. *Capsf.* 1 - locularis, apice 5 - valvis, infera. *Cor.* hypocraterif. rotata. *Stamina* munita squamulis corollæ. *Stig.* capitatum.
73. PHYTEUMA. *Capsf.* 2 - 3 - locularis, infera, perforata. *Cor.* rotata, laciniis linearibus 5 - partita. *Stigma* 2 - f. 3 - fidum.
71. CAMPANULA. *Capsf.* 3 - f. 5 - locularis, perforata, infera. *Cor.* campanulata fundo clauso valvis staminiferis. *Stigma* 3 - fidum.



74. LONICERA. *Bacca* 2 - locularis : subrotunda, infera, polysperma. *Cor.* monopetala inæqualis. *Stigm.* capitatum.

Flores *pentapetali*, *inferi*.

80. RHAMNUS. *Bacca* 3 - locularis, rotunda. *Cal.* tubulosus, corollifer. *Squamae* oris 5. convergentes, *stamina* munientes. *Cor.* nulla.

81. EVONYMUS. *Bacca* capsularis, lobata. *Cal.* patens. *Cor.* 5 - petala. *Caps.* 5 - gona, 5 - locularis, 5 - valvis, colorata. *Sem.* calyptrata.

antehort + Viole.

Flores *pentapetali*, *superi*.

82. RIBES. *Bacca* polysperma, infera. *Cal.* corollifer. *Petala* 5. & *stamina* calyci inserta. *Stylus* 2 - fidus.

82. HEDERA. *Bacca* 5 - sperma. *Cal.* cingens fructum. *Stigm.* simplex.

Flores *incompleti*, *inferi*.

84. GLAUX. *Caps.* 1 - locularis, 5 - valvis, 5 - sperma. *Cal.* 1 - phyllus, rudior, campanulatus. *Cor.* nulla.

DIGYNIA.

Flores *monopetali*, *inferi*.

85. ASCLEPIAS. *Folliculi* 2. *Cor.* reflexa. *Nettaris* 5. auriformibus, unguiculatis.

90. GENTIANA. *Caps.* 1 - locularis, 2 - valvis, receptaculis 2. longitudinalibus. *Cor.* tubulosa, indeterminata.

Flores

Flores *incompleti*.

88. SALSOLA. *Sem.* 1. cochleatum, tectum. *Cal.* 5 - phyllus. *Cor.* 0. *Caps.* 1 - sperma.
87. CHENOPODIUM. *Sem.* 1. lenticulare, superum. *Cal.* 5 - phyllus, 5 - gonus, foliolis concavis. *Cor.* 0.
86. HERNIARIA. *Sem.* 1, ovatum, tectum. *Cal.* 5 - partitus. *Stamina* 5 sterilia. *Cor.* 0.
89. ULMUS. *Bacca* exsucca, compresso - membranaea. *Cal.* 5 - fidus emarcescens. *Cor.* 0.

Flores pentapetali, superi, dispermi. *Umbellatae*.

*) *Involucro universali partialique.*

91. ERYNGIUM. *Fl.* capitati. *Receptaculo* paleaceo.
92. HYDROCOTYLE. *Fl.* subumbellati, fertiles. *Umbella* simplici: *Involucro* 4 - phyllo. *Petala* integra. *Sem.* semiorbiculata, compressa.
93. SANICULA. *Fl.* subumbellati, abortivi, disco steriles: *Umbellis* confertis, subcapitatis. *Sem.* muricata.
101. HERACLEUM. *Flores* radiati, fertiles. *Involucr.* deciduum. *Cor.* difformis. *Sem.* membranacea.
106. OENANTHE. *Fl.* radiati, abortivi, disco steriles: fessiles, difformes. *Involucr.* simplex. *Sem.* coronata.
97. DAUCUS. *Fl.* radiati, fertiles. *Invol.* pinnatum. *Sem.* hispida.
95. TORDYLIUM. *Fl.* radiati, fertiles. *Invol.* longa, indivisa. *Sem.* suborbiculata, margine crenata.
96. CAUCALIS. *Flores* radiati, abortivi. *Invol.* simplex. *Sem.* muricata.

98. CONIUM. *Fl.* flosculosi, fertiles. *Petala* cordata. *Sem.* gibba costato - fulcata.
94. BUPLEURUM. *Fl.* flosculosi, fertiles. *Pet.* cordata. *Involucra* umbellulæ majora, 5 - phylla. *Fructus* subrotundus, compressus, striatus.
100. ATHAMANTA. *Fl.* flosculosi, fertil. *Pet.* cordata. *Sem.* ovato - oblonga, convexa, striata.
104. SIUM. *Fl.* flosc. fert. *Pet.* cordata. *Semina* subovata, striata. *Invol.* polyphyllum.
99. SELINUM. *Fl.* flosc. fert. *Pet.* cordata æqualia. *Sem.* ovali - oblonga, compressa, plana, in medio striata. *Invol.* reflexum.
102. LIGUSTICUM. *Flores* floscul. fertiles. *Pet.* planiuscula. *Fruct.* oblongus, 5 - sulcatus utrinque. *Involucr.* membranacea.
103. ANGELICA. *Fl.* flosc. fert. *Pet.* planiuscula, incurva. *Umbellula* globosæ. *Corollæ* æquales. *Fructus* subrotundus, angulatus, solidus, sulcis utrinque 3.
105. SISON. *Fl.* flosc. fert. *Pet.* planiuscula. *Umbellula* depauperata. *Fructus* ovatus, striatus. *Invol.* subtetraphyllum.

β) *Involucris partialibus; universali nullo,*

110. SCANDIX. *Fl.* subrad. abortivi. *Fruct.* oblongi, subulati. *Cor.* difformis. *Petala* integra.
111. CHEROPHYLLUM. *Fl.* flosc. abortivi. *Invol.* 5 - phyllum, reflexum, concavum. *Petala* cordata. *Fructus* oblongus, lævis.
109. ÆTHUSA. *Fl.* flosc. fertiles. *Invol.* dimidiatum, 3 - phyllum, pendulum. *Fructus* striatus.
107. PHELLANDRIUM. *Fl.* flosc. fertiles. *Fruct.* ovatus, lævis, coronatus perianthio & pistillo.
112. IMPERATORIA. *Fl.* flosc. fertiles. *Umbella* expanso - plana. *Fructus* subrotundus, compressus, medio gibbus, margine cinctus.

108. **CICUTA.** *Fl.* flosc. fertiles. *Pet.* planiuscula, ovata, inflexa. *Styli* corolla longiores. *Fructus* subovatus, sulcatus, bipartibilis.

2) *Involucro* nullo; nec universali, nec partiali.

113. **PASTINACA.** *Fl.* flosc. fertil. *Sem.* elliptica, depresso-plana.

117. **ÆGOPodium.** *Fl.* flosc. fert. *Pet.* cordata, concava. *Sem.* gibba, striata.

116. **APIUM.** *Fl.* flosc. fert. *Sem.* ovata, minuta, striata. *Invol.* univers. 1-phyllum. *Petala* inflexa.

115. **PIMPINELLA.** *Fl.* flosc. fert. subradiati. *Umbellae* ante florescentiam nutantes: *Sem.* ovato-oblonga.

114. **CARUM.** *Fl.* flosc. fert. *Petala* cordata, emarginata, carinata. *Sem.* gibba, striata. *Invol.* 1-phyllum.

TRIGYNIA.

* Flores *superi.*

118. **VIBURNUM.** *Cor.* 5-fida, campanulata. *Bacca* 1-sperma. *Cal.* 5-partitus.

119. **SAMBUCUS.** *Cor.* 5-fida, rotato-concava. *Bacca* 3-sperma. *Cal.* 5-partitus.

* * Flores *inferi.*

120. **ALSINE.** *Cor.* 5-petala. *Caps.* 1-locularis, 3-valvis. *Cal.* 5-phyllus. *Petala* 2-fida.

121. **CORRIGIOLA.** *Cor.* 5-petala. *Cal.* 5-phyllus. *Sem.* 1, triquetrum.

† *Polygonum amphibium.*

TETRAGYNIA.

122. PARNASSIA. *Cal.* 5 - partitus. *Pet.* 5. *Nectaria* 5, cordata, ciliata apicibus globosis. *Caps.* 4 - valvis.

PENTAGYNIA.

124. LINUM. *Cal.* 5 - phyllus. *Cor.* 5 - petala. *Caps.* 5 - valvis, 10 - locularis. *Sem.* solitaria.
 125. DROSERA. *Cal.* 5 - fidus. *Cor.* 5 - petala. *Caps.* 1 - locularis, apice 5 - valvi. *Sem.* plurima.
 123. STATICE. *Cal.* 1 - phyllus, integer, plicatus. *Cor.* 5 - petala, *Sem.* 1. calyce infundibulif. vestitum.

† *Cerastium pentandrum.*
Spergula pentandra.

POLYGYNIA.

126. MYOSURUS. *Cal.* 5 - phyllus, basi adnatus. *Nect.* 5, subulata, petaliformia. *Sem.* numerosa.
 † *Ranunculus hederaceus.*



MONOGYNIA.

56. MYOSOTIS. 180.

147. MYOSOTIS *scorpioides* feminibus nudis: foliorum apicibus callofis. P. Fl. Oec. 101.

a) *Arvensis*, foliis hirsutis.

β) *Palustris*, foliis glabris.

Dan. *Forgat-Migej*; *Skorpions-Urt.*

In pratis, in nemore ubique. β, ad aquas.

148. MYOSOTIS *Lappula* feminibus aculeis triglochidibus: foliis lanceolatis pilosis.

Dan. *Hunde-Tunge.*

In argillosis & ruderatis.

57. LITHOSPERMUM. 181.

149. LITHOSPERMUM *officinale* feminibus lævibus: corollis vix calycem superantibus: foliis lanceolatis.

Dan. *Steenbrak.*

Pharmac. LITHOSPERMI semina.

In nemore ad hertum.

150. † LITHOSPERMUM *arvense* feminibus rugosis, corollis vix calycem superantibus.

P. Fl. Oec. 93.

Dan. *Steen-Klinte.*

In agris.

58. ANCHUSA. 182.

151. ANCHUSA *officinalis* foliis lanceolatis: spicis imbricatis secundis.

Dan. *Oortunge.*

Pharmac. EUGLOSSI radix, herba.

In agris.

152. † *ANCHUSA angustifolia* racemis subnudis conjugatis.

59. CYNOGLOSSUM. 183.

153. *CYNOGLOSSUM officinale* staminibus corolla brevioribus: foliis lato-lanceolatis tomentosis sessilibus.

Dan. *Hunde-Tunge*; *Uldhorrer*.

Pharmac. CYNOGLOSSI radix, herba.

Ad pagos in macerie, rudetatis.

60. PULMONARIA. 184.

154. *PULMONARIA officinalis* foliis radicalibus ovato-cordatis scabris.

Dan. *Lunge-Urt*.

Pharmac. PULMONARIÆ herba.

In sylvaticis.

155. *PULMONARIA angustifolia* foliis radicalibus lanceolatis.

Ad *Dronninggaard* copiose.

61. SYMPHYTUM. 185.

156. *SYMPHYTUM officinale* foliis ovato-lanceolatis.

P. Fl. Oec. 141.

Dan. *Kulsukker-Rod*; *Gongsalve-Rod*.

Pharmac. CONSOLIDÆ MAJORIS radix, herba, flores.

In subhumidis.

62. ASPERUGO. 189.

157. *ASPERUGO procumbens* calycibus fructus compressis. *Morif. Hist. III. p. 439. S. XI. t. 26. f. 13.*

In rudetatis, rarior.

63. LYCOPSIS. 190.

158. LYCOPSIS *arvensis* foliis lanceolatis hispidis: calycibus florescentibus erectis.

Dan. *Liden Oxetunge*.

Inter segetes.

64. ECHIUM. 191.

159. ECHIUM *vulgare* caule tuberculato hispido: foliis caulinis lanceolatis hispidis; floribus spicatis lateralibus. P. Fl. Oec. 61.

Dan. *Vild Oxetunge*; *Slange-Hoved*.

In maceriebus, ac ad sepes frequens.

65. PRIMULA. 197.

160. PRIMULA *veris* foliis denticulatis rugosis.

P. Fl. Oec. 114.

a) *Officinalis* - - limbo corollarum concavo.

β) *Elatior* - - limbo corollarum plano.

† γ) *Acaulis* - - scapo nullo. Fl. Dan. t. 194.

Dan. *Koe-Driver*.

Pharmac. PRIMULÆ VERIS radix, herba, flores.

a) In pratis versus lacum *Fuursøe* copiose.

β) passim.

161. † PRIMULA *farinosa* foliis crenatis glabris: limbo florum plano.

Fl. Dan. t. 125.

Dan. *Enge-Stierne*; *Blaa-May*.

In campis sterilibus.

66. MENYANTHES. 202.

162. MENYANTHES *trifoliata* foliis ternatis.

P. Fl. Oec. 99.

Dan. *Bukke-Blade*; *Beske-Klever*.

Pharmac. TRIFOLII AQUATICI seu FIBRINI herba.

Ad margines paludum frequens.

67. HOTTONIA. 203.

163. HOTTONIA *palustris* pedunculo verticillato, multifloro.

Dan. Vand-röllike.

In paludibus copiosissime.

68. LYSIMACHIA. 205.

* *Pedunculis multifloris.*

164. LYSIMACHIA *vulgaris* paniculata : racemis terminalibus.

Dan. Fred-Lös.

In paludibus nemorosis.

165. LYSIMACHIA *thyrsiflora* racemis lateralibus pedunculatis.

Dan. Due-Urt.

Ibidem.

* * *Pedunculis unifloris.*

166. † LYSIMACHIA *Nummularia* foliis subcordatis : floribus solitariis : caule repente.

Dan. Gul Petje-Blad.

Pharmac. NUMMULARIÆ herba.

In depressis locis ditionis *Kiloniensis* frequens.

167. † LYSIMACHIA *nemorum* foliis ovatis acutis : floribus solitariis : caule procumbente.

fl. Dan. t. 174.

Dan. Gul-Arfve.

In nemorosis.

69. ANAGALLIS. 206.

168. ANAGALLIS *arvensis* foliis indivisis : caule procumbente. fl. Dan. t. 88.

Dan. Röd-arfve ; Grine ved Middag.

In arvis haud infrequens.

70. CONVULVULUS. 215.

169. CONVULVULUS *arvensis* foliis sagittatis utrinque acutis: pedunculis unifloris.

Dan. *Snerrer*.

Inter Segetes.

170. CONVULVULUS *sepium* foliis sagittatis, postice truncatis: pedunculis tetragonis unifloris.

Dan. *Stor Snerle*; *vor Frues Særk*.

In alneto Valhalla.

71. CAMPANULA. 218.

† *Foliis laevioribus, angustioribus.*

171. CAMPANULA *rotundifolia* foliis radicalibus reniformibus: caulinis linearibus.

Dan. *Smaa Klokke*.

In pratis.

172. CAMPANULA *patula* foliis strictis lanceolato-ovatis: panicula patula.

In colle *Brankebakken* raro.

173. † CAMPANULA *Rapunculus* foliis undulatis: radicalibus lanceolato-ovalibus: panicula coarctata.

P. Fl. Qec. 43.

Dan. *Rapuntzel*.

174. CAMPANULA *decurrens* foliis radicalibus lobatis: caulinis lanceolato-linearibus subserratis sessilibus, remotis.

In dumetis.

In Flora utraque Suecia & in priori Spec. editione nomen hujus tribus suis *Perfoliata*, in secunda Spec. editione *decurrens*.



Pentandria: Monogynia.

* * *Foliis scabris, latioribus.*

175. CAMPANULA *Trachelium* caule angulato: foliis petiolatis: calycibus ciliatis: pedunculis trifloris.

Dan. *vilde Klokke*.

In nemore.

176. † CAMPANULA *glomerata* caule angulato simplici: floribus sessilibus: capitulo terminali.

177. † CAMPANULA *cervicaria* hispida: floribus sessilibus: capitulo terminali: foliis lanceolato-linearibus undulatis.

178. † CAMPANULA *thyrsoides* hispida racemo ovato-oblongo terminali: caule simplicissimo: foliis lanceolato-linearibus.

72. SAMOLUS. 222.

179. † SAMOLUS *Valerandi*. Fl. Dan. t. 198.

In subhumidis maritimis.

73. PHYTEUMA. 220.

180. † PHYTEUMA *spicata* spica oblonga: capsulis bilocularibus: foliis radicalibus cordatis.

In montosis sylvaticis.

74. LONICERA. 233.

181. LONICERA *Periclymenum* capitulis ovatis imbricatis terminalibus: foliis omnibus distinctis.

Dan. *Löbe-Rose*.

Pharmac. CAPRIFOLII herba.

In dumetis.

182. * LONICERA *Xylesteum* pedunculis bifloris: bacis distinctis: foliis integerrimis pubescentibus.

In sepibus raro.

75. VERBASCUM. 245.

183. VERBASCUM *Thapsus* foliis decurrentibus utrinque tomentosis.

Dan. *Konge - Lys.*

Pharmac. VERBASCI folia, radix.

In glareosis ac sepibus copiose.

184. * VERBASCUM *Lychnitis* foliis cuneiformi-oblongis.

In ruderatis villæ veteris.

† * VERBASCUM *phlomoïdes* foliis ovatis utrinque tomentosis: inferioribus petiolatis?

In montosis promontorii insulæ *Mona* legi.

185. VERBASCUM *nigrum* foliis oblongo-cordatis petiolatis.

Ad sepes & vias frequens.

186. † VERBASCUM *Blattaria* foliis amplexicaulibus oblongis glabris: pedunculis solitariis.

Circa *Hanoviam* in Germania vulgarem offendi.

In argillaceis.

76. DATURA. 246.

187. DATURA *Stramonium* pericarpis spinosis erectis ovatis: foliis ovatis glabris.

Pharmac. Semenibus: *cautendum.*

In ruderatis ad littus maris frequens.

77. HYOSCYAMUS. 247.

188. HYOSCYAMUS *niger* foliis amplexicaulibus sinuatis: floribus sessilibus. *P. Fl. Oec. 84.*

Dan. *Bulme - Urt; Horse - Tender; Soe - Bønn.*

Pharmac. HYOSCYAMI radix, herba, semina.

Ad pagos & maceries.

78. SOLANUM. 251.

189. SOLANUM *nigrum* caule inermi herbaceo: foliis ovatis angulatis: umbellis nutantibus.

Dan. *Swine-Urt*; *Sov-Bær*.

In ruderalis.

190. SOLANUM *Dulcamara* caule inermi frutescente flexuoso: foliis superioribus hastatis: racemis cy-mosis. P. Fl. Oec. 137

Dan. *Bitter-Søde*; *Hunde-Bær*.

Pharmac. SOLANI folia.

In alnetis ad amnem.

79. PHYSALIS. 250.

191. * PHYSALIS *Alkekengi* foliis geminis integris actis: caule herbaceo inferne subramoso.

Pharmac. ALKEKENGII herba, baccæ.

In nemore ad hortum.

80. RHAMNUS. 265.

192. † RHAMNUS *catharticus* spinis terminalibus: floribus quadrifidis dioicis: foliis ovatis.

P. Fl. Oec. 123.

Dan. *Korsbær-Torn*; *Vrie-Torn*.

Pharmac. SPINÆ QERVINÆ baccæ.

193. RHAMNUS *Frangula* inermis floribus monogynis hermaphroditis: foliis integerrimis.

P. Fl. Oec. 124.

Dan. *Tørste-Træ*.

Pharmac. FRANGULÆ cortex.

In nemorosis ad amnem.

81. EVONYMUS. 271.

194. EVONYMUS *Europæus* floribus plerisque quadrididis. P. Fl. Oec. 68.

Dan. *Beenved.*

In nemorosis raro.

82. RIBES. 281.

* *Ribesia inermia.*

195. RIBES *rubrum* inerme, racemis glabris pendulis: floribus planiusculis. P. Fl. Oec. 125.

Dan. *Ribs.*

Pharmac. RIBIUM bacca.

Passim ad amnem.

196. RIBES *nigrum* inerme, racemis pilosis: floribus oblongis.

Dan. *Soel-Bar.*

Pharmac. RIBIS NIGRI bacca.

In præcipitiis lacus *Fuursøe.*

* * *Grossularia aculeata.*

197. * RIBES *uva crispa* ramis aculeatis: baccis glabris: pedicellis bractea monophylla.

Ad sepes.

83. HEDERA. 283.

198. HEDERA *helix* foliis ovatis lobatisque.

Dan. *Ved-Bende.*

Pharmac. HEDERÆ folia, bacca, gummi.

In Nemoris trunco Fagi, raro.

84. GLAUX. 291.

199. GLAUX *maritima*.Dan. *Strand-Isop*.In littore maris ad *Webek*.

DIGYNIA.

85. ASCLEPIAS. 306.

200. ASCLEPIAS *Vincetoxicum* foliis ovatis basi barbatis: caule erecto: umbellis proliferis.Dan. *Svaleroed*.*Pharmac.* HIRUNDINARIÆ radix, herba.

Circa molendina.

86. HERNIARIA. 308.

201. HERNIARIA *glabra* herbacea glabra.Dan. *Brid-Urt*; *Pisse-Urt*.*Pharmac.* HERNIARIÆ herba.

In collibus e regione horti.

87. CHENOPODIUM. 309.

* *Foliis angulosis.*202. CHENOPODIUM, *bonus Henricus*, foliis triangulari-fagittatis integerrimis: spicis compositis aphyllis. *P. Fl. Oec.* 50.Dan. *Stolt Henrik*.*Pharmac.* BONI HENRICI herba.

In ruderatis.

203. CHENOPODIUM *urbicum* foliis triangularibus subdentatis: racemis confertis strictissimis, cauli approximatis longissimis.

Ad pagos.

204. *CHENOPODIUM rubrum* foliis cordato-triangularibus obtusiusculis dentatis: racemis erectis compositis foliosis, caule brevioribus.

Pharmac. ATRIPLICIS sylvestris herba.

In ruderatis.

205. *CHENOPODIUM album* foliis rhomboïdeo-triangularibus erosis, postice integris: summis oblongis, racemis erectis.

Vaill. Bot. Par. t. 7. f. 1.

In cultis.

206. *CHENOPODIUM viride* foliis rhomboïdeis dentato-sinuatis: racemis ramosis subfoliatis.

In oleraceis.

207. *CHENOPODIUM hybridum* foliis cordatis angulato-acuminatis: racemis ramosis nudis.

Vaill. Bot. Paris. t. 7. f. 2.

Inter olera.

208. *CHENOPODIUM glaucum* foliis ovato-oblongis repandis: racemis nudis simplicibus glomeratis.

In ruderatis.

* * *Foliis integris.*

209. † *CHENOPODIUM vulvaria* foliis integerrimis rhombeo-ovatis: floribus conglomeratis axillaribus.

210. † *CHENOPODIUM polyspermum* foliis integerrimis ovatis: caule decumbente: cymis dichotomia aphyllis axillaribus.

211. *CHENOPODIUM maritimum* foliis subulatis semicylindricis.

In littore maris Balthici; ad *Lundsgaard & Hindsgavl Fionia* frequens.

88. SALSOLA. 311.

212. SALSOLA *Kali* herbacea decumbens: foliis subulatis spinosis: calycibus marginatis axillaribus.
Meris. Hist. III. p. 611. S. V. t. 33. f. 11.
 In littore sabuloso maris, ad *Webeck, Lundsgaard, Snoghoy.*
213. † SALSOLA *Tragus* herbacea erecta: foliis subulatis spinosis lævibus: calycibus ovatis.
214. SALSOLA *hirsuta* diffusa herbacea: foliis teretibus obtusis tomentosis. *Fl. Dan. t. 187.*
 In littore maris.

89. ULMUS. 316.

215. † ULMUS *campestris* foliis duplicato-ferratis, basi inæqualibus. *Pauli Fl. Oec. 157.*
Dan. Alm.
Pharmac. ULMI cortex medius.

90. GENTIANA. 322.

* *Corollis quinquefidis, subcampaniformibus.*

216. † GENTIANA *pneumonanthe* corollis quinquefidis campanulatis oppositis: pedunculatis: foliis linearibus.
 In subhumidis ericetis ad *Rensburgum* legi.

* *Corollis quinquefidis infundibuliformibus.*

217. GENTIANA *Centaurium* corollis quinquefidis infundibuliformibus: caule dichotomo.
Dan. Tusindgylden.
Pharmac. CENTAURII minoris herba.
 In sterilibus.

218. † GENTIANA *Amarella* corollis quinquefidis hypocrateriformibus, fauce barbatis.

219. GENTIANA *campestris* corollis quadrifidis fauce barbatis.

Dan. *Enzian*.

Pharmac. GENTIANELLÆ herba.

Circa hortum in pratis aridis; floret ante & post æquinoctium.

91. ERYNGIUM. 324.

220. † ERYNGIUM *maritimum* foliis radicalibus subrotundis, plicatis, spinosis: capitulis pedunculatis.

Dan. *Strandmands-Troe; Hundrede-Hoved*.

Pharmac. ERYNGII radix, herba.

In littore arenoso ad oppidum *Præste* legi.

221. † ERYNGIUM *campestre* foliis amplexicaulibus pinnato-laciniatis.

Dan. *Biergmands-Troe*.

Pharmac. ERYNGII radix.

92. HYDROCOTYLE. 325.

222. HYDROCOTYLE *vulgaris* foliis peltatis: umbellis quinquefloris.

Fl. Dan. t. 90.

Dan. *Vand-Krage-Taer*.

In paludoso ad hortum.

93. SANICULA. 326.

223. SANICULA *Europæa* foliis radicalibus simplicibus: flosculis omnibus sessilibus.

Dan. *Sanikel*.

Pharmac. SANICULÆ herba.

In Sylvaticis versus lacum *Fuursjøe*.

D

94. BUPLEURUM. 328.

224. † BUPLEURUM *rotundifolium* involucris universalibus nullis: foliis perfoliatis.

Dan. *Gienem vaxt.*

95. TORDYLIUM. 330.

225. TORDYLIUM *Anthriscus* umbellis confertis: foliolis ovato-lanceolatis pinnatifidis.

Morif. Hist. p. 30. t. 14. f. 8.

In dumetis.

96. CAUCALIS. 331.

226. CAUCALIS *leptophylla* feminum aculeis triglochidibus uncinatis: pilis verticillatis hispidis.

Dan. *Bettler - Luus.*

97. DAUCUS. 333.

227. DAUCUS *Carota* feminibus hispidis: petiolis subtus nervosis.

Dan. *Vilde Gule - Rödder; wild Pastinakel.*

Pharmac. DAUCI SYLVESTRIS femina.

In campis.

98. CONIUM. 336.

228. CONIUM *maculatum* feminibus striatis.

P. Fl. Oec. 54.

Dan. *Skarntyde.*

Pharmac. CICUTÆ herba, radix.

Inter fegetes.

99. SELINUM. 337.

229. SELINUM *palustre* sublaetescens, radice unica.

In paludibus Nemoris.

100. ATHAMANTA. 338.

230. ATHAMANTA *Oreoselinum* foliis divaricatis.

P. Fl. Oec. 32.

Dan. *Vilde Gule* - Rödder.

In collibus passim.

101. HERACLEUM. 345.

231. HERACLEUM *Sphondylium* foliolis pinnatifidis.

P. Fl. Oec. 79.

Dan. *Biörne* - Kloe; *Biörne* - Lap.

Ad margines agrorum passim.

102. LIGUSTICUM. 346.

232. † LIGUSTICUM *levisticum* foliis multiplicibus:
foliolis superne ineis.

Crescit quoque in ruderatis horti veteris, vix tamen
pro indigena habenda planta.

233. † LIGUSTICUM *Scoticum* foliis biternatis.

Fl. Dan. t. 207.

Ad littora maris.

103. ANGELICA. 347.

234. ANGELICA *Archangelica* foliorum impari lobato.

Fl. Dan. t. 206. P. Fl. Oec. 19.

Dan. *Angelik*.

Pharmac. ANGELICÆ SATIVÆ radix, herba, se-
mina.

Ad hortum in culta.

235. ANGELICA *sylvestris* foliolis æqualibus ovato-
lanceolatis ferratis.

Pharmac. ANGELICÆ SYLVESTRIS radix.

In paludibus nemorosis.

104. S I U M. 348.

236. SIUM *latifolium* foliis pinnatis : umbellis terminalibus.

Dan. *Vand Pastinak.*

Ad amnem.

237. † SIUM *nodiflorum* foliis pinnatis : umbellis axillaribus sessilibus.

105. S I S O N. 349.

238. SISON *inundatum* repens : umbellis bifidis.
Fl. Dan. t. 89.

In inundatis prati, vulgo *Ladegaard.*

106. O E N A N T H E. 352.

239. OENANTHE *fistulosa* stolonifera, foliis caulinis pinnatis filiformibus fistulosis.

Dan. *Vand - Stenbrek.*

In inundato paludoso Vivarii.

240. † OENANTHE *crocata* foliis omnibus multifidis obtusis subæqualibus.

107. P H E L L A N D R I U M. 353.

241. PHELLANDRIUM *aquaticum* foliorum ramificationibus divaricatis. *P. Fl. Oec. 106.*

Dan. *Skarn - Tyde.*

Ad amnem.

108. C I C U T A. 354.

242. CICUTA *virosa* umbellis oppositifoliis : petiolis marginatis obtusis. *Fl. Dan. t. 208. P. Fl. Oec. 52.*

Dan. *Vand Pastinak.*

Pharmac. CICUTÆ herba.

In ripa paludosa amnis.

109. ÆTHUSA. 355.

243. ÆTHUSA *Cynapium*. *Pauli Fl. Oec.* 7.
 Dan. *Vild Petersilie*.
 Inter olera.

110. SCANDIX. 357.

244. † SCANDIX *pecten Veneris* feminibus lævibus,
 rostro longissimo.
 245. SCANDIX *Cerrefolium* feminibus nitidis, ovato-
 subulatis: umbellis sessilibus lateralibus.
 Dan. *Körvel*.
Pharmac. CEREFOLII herba, semina.
 In agris.
 246. SCANDIX *Anthriscus* feminibus ovatis hispidis:
 corollis uniformibus: caule lævi.
 Dan. *Vild Körvel*.
 In ruderatis.

111. CHÆROPHYLLUM. 358.

247. CHÆROPHYLLUM *sylvestre* caule striato: genu-
 lis tumidiusculis. *P. Fl. Oec.* 49.
 Dan. *Vild Körvel; Skarutyle*.
Pharmac. CICUTARIÆ herba.
 Ad septimenta, in pratis, arvis, sylvis vulgarissi-
 mum.
 248. CHÆROPHYLLUM *temulentum* caule scabro: ge-
 niculis tumidis: umbellis cernuis.
Meris. Hist. III. p. 302. t. 101. f. 7.
 Intra maceriem arboreti, vulgo *Deyligheden*.
 Involucrum, s. phyllum, reflexum, concavum.

112. IMPERATORIA. 359.

249. † IMPERATORIA *Ostruthium*.

113. PASTINACA. 362.

250. PASTINACA *fativa* foliis simpliciter pinnatis.
P. Fl. Oec. 105.
 Dan. *Vild Pastinak*.
 In sepimento *Deyligbeden*.

114. CARUM. 365.

251. CARUM *Carvi*. *P. Fl. Oec.* 46.
 Dan. *Kummen*.
Pharmac. CARVI femina.
 In pratis raro.

115. PIMPINELLA. 366.

252. PIMPINELLA *saxifraga* foliis pinnatis; foliolis radicalibus subrotundis; summis linearibus.
 Dan. *Quase-Urt*; *Bierg-Rad*; *vild Pimpinelle*.
Pharmac. PIMPINELLA *alba* radix, herba, femina.
 In glareosis.

116. ARIUM. 367.

253. ARIUM *graveolens* foliolis caulinis cuneiformibus; umbellis sessilibus.
 Dan. *Peterfilie*; *Vand-Merke*.
Pharmac. ARII radix, herba, femina.

117. EGOPodium. 368.

254. EGOPodium *podagraria* foliis caulinis summis ternatis.
Pauli Fl. Oec. 6.
 Dan. *Busmekker*; *Squalder-Kaal*.
 In dumetis collis *Branckebakke*.

TRIGYNIA.

118. VIBURNUM. 370.

255. VIBURNUM *Opulus* foliis lobatis : petiolis glandulosis. P. Fl. Oec. 155.

Dan. Fugle - Bär ; Vand - Hyld.

In pratis subhumidis.

119. SAMBUCUS. 372.

256. SAMBUCUS *nigra* cymis quinquepartitis : caule arboreo. P. Fl. Oec. 131.

Dan. Hyld.

Pharmac. SAMBUCI cortex, folia, flores, baccæ, femina.

Ad sepes ubique.

257. SAMBUCUS *Ebulus* cymis trifidis : stipulis foliaceis : caule herbaceo. P. Fl. Oec. 130.

Dan. Sommer - Hyld ; Attik.

Pharmac. EBULI radix, folia, flores, baccæ, femina.

In umbrosis nemoris.

120. ALSINE. 380.

258. ALSINE *media* petalis bipartitis : foliis ovato-cordatis. Pauli Fl. Oec. 16.

Dan. Fugle - Græs.

In cultis ubique.

121. CORRIGIOLA. 378.

259. † CORRIGIOLA *littoralis*.

In campis arenosis Slesvici circa Rensburgum copiosissima.

104. S I U M. 348.

236. *SIUM latifolium* foliis pinnatis : umbellis terminalibus.

Dan. *Vand Pastinak.*

Ad amnem.

237. † *SIUM nodiflorum* foliis pinnatis : umbellis axillaribus sessilibus.

105. S I S O N. 349.

238. *SISON inundatum* repens : umbellis bifidis.

Fl. Dan. t. 89.

In inundatis prati, vulgo *Ladegaard.*

106. O E N A N T H E. 352.

239. *OENANTHE fistulosa* stolonifera, foliis caulinis pinnatis filiformibus fistulosis.

Dan. *Vand - Stenbrek.*

In inundato paludoso Vivarii.

240. † *OENANTHE crocata* foliis omnibus multifidis obtusis subæqualibus.

107. P H E L L A N D R I U M. 353.

241. *PHELLANDRIUM aquaticum* foliorum magnificationibus divaricatis. *P. Fl. Occ. 106.*

Dan. *Skarn - Tyde.*

Ad amnem.

108. C I C U T A. 354.

242. *CICUTA virosa* umbellis oppositifoliis : petiolis marginatis obtusis. *Fl. Dan. t. 208. P. Fl. Occ. 52.*

Dan. *Vand Pastinak.*

Pharmac. CICUTÆ herba.

In ripa paludosa amnis.

109. ÆTHUSA. 355.

243. ÆTHUSA *Cynapium*. *Pauli Fl. Oec.* 7.
 Dan. *Vild Petersilie*.
 Inter olera.

110. SCANDIX. 357.

244. † SCANDIX *pecten Veneris* feminibus lævibus,
 rostro longissimo.
 245. SCANDIX *Cerefolium* feminibus nitidis, ovato-
 subulatis: umbellis sessilibus lateralibus.
 Dan. *Körvel*.
Pharmac. CEREFOLII herba, semina.
 In agris.
 246. SCANDIX *Anthriscus* feminibus ovatis hispidis:
 corollis uniformibus: caule lævi.
 Dan. *Vild Körvel*.
 In ruderatis.

111. CHÆROPHYLLUM. 358.

247. CHÆROPHYLLUM *sylvestre* caule striato: genu-
 lis tumidiusculis. *P. Fl. Oec.* 49.
 Dan. *Vild Körvel; Skarstjle*.
Pharmac. CICUTARIÆ herba.
 Ad sepimenta, in pratis, arvis, sylvis vulgarissi-
 mum.
 248. CHÆROPHYLLUM *temulentum* caule scabro: ge-
 niculis tumidis: umbellis cernuis.
Morif. Hist. III. p. 302. t. 101. f. 7.
 Intra maceriem arboreti, vulgo *Deyligheden*.
 Involucrum, 8. phyllum, reflexum, concavum.

112. IMPERATORIA. 359.

249. † IMPERATORIA *Ostrachium*.

113. PASTINACA. 362.

250. PASTINACA *sativa* foliis simpliciter pinnatis.
P. Fl. Oec. 105.
 Dan. *Vild Pastinak.*
 In sepimento *Deyligbeden.*

114. CARUM. 365.

251. CARUM *Carvi.* *P. Fl. Oec.* 46.
 Dan. *Kummen.*
Pharmac. CARVI femina.
 In pratis raro.

115. PIMPINELLA. 366.

252. PIMPINELLA *saxifraga* foliis pinnatis; foliolis radicalibus subrotundis: summis linearibus.
 Dan. *Quase-Urt; Bierg-Rad; vild Pimpinelle.*
Pharmac. PIMPINELLE *alba* radix, herba, femina.
 In glareosis.

116. APIUM. 367.

253. APIUM *graveolens* foliolis caulinis cuneiformibus: umbellis sessilibus.
 Dan. *Peterfilie; Vand-Merke.*
Pharmac. APII radix, herba, femina.

117. EGOPodium. 368.

254. EGOPodium *podagraria* foliis caulinis summis ternatis.
Pauli Fl. Oec. 6.
 Dan. *Busmekker; Squaller-Kaal.*
 In dumetis collis *Brankebakke.*

TRIGYNIA.

118. VIBURNUM. 370.

255. VIBURNUM *Opulus* foliis lobatis : petiolis glandulosis. P. Fl. Oec. 155.

Dan. *Fugle - Bär* ; *Vand - Hyld*.

In pratis subhumidis.

119. SAMBUCUS. 372.

256. SAMBUCUS *nigra* cymis quinquepartitis : caule arboreo. P. Fl. Oec. 131.

Dan. *Hyld*.

Pharmac. SAMBUCI cortex, folia, flores, baccæ, femina.

Ad sepes ubique.

257. SAMBUCUS *Ebulus* cymis trifidis : stipulis foliaceis : caule herbaceo. P. Fl. Oec. 130.

Dan. *Sommer - Hyld* ; *Attik*.

Pharmac. EBULI radix, folia, flores, baccæ, femina.

In umbrosis nemoris.

120. ALSINE. 380.

258. ALSINE *media* petalis bipartitis : foliis ovato-cordatis. Pauli Fl. Oec. 16.

Dan. *Fugle - Græs*.

In cultis ubique.

121. CORRIGIOLA. 378.

259. † CORRIGIOLA *littoralis*.

In campis arenosis *Slesvici* circa *Rensburgum* copiosissima.

TETRAGYNIA.

122. PARNASSIA. 384.

260. PARNASSIA *palustris*.

Dan. *Lever-Blomster*.

Pharmac. HEPATICÆ ALBÆ herba, flores.

In pratis humidis circa *Nyebroc*.

PENTAGYNIA.

123. STATICE. 388.

261. STATICE *armeria* scapo simplici capitato: foliis linearibus.

Dan. *Bierg Nellike*.

In pascuis ad semitas.

262. † STATICE *linonium* scapo paniculato tereti: foliis lævibus.

124. LINUM. 389.

263. LINUM *catharticum* foliis oppositis ovato-lanceolatis: caule dichotomo: corollis acutis.

P. Fl. Oec. 92.

In pratis subhumidis.

264. LINUM *Radiola* foliis oppositis: caule dichotomo: floribus tetrandris tetragynis. *Fl. Dan. t.* 178.

In arenosis.

125. DROSERA. 391.

265. DROSERA *rotundifolia* scapis radicatis: foliis orbiculatis. *P. Fl. Oec.* 60.

Dan. *Soel-Dug; Himmel-Dug*.

Pharmac. RORIS SOLIS herba.

In paludibus turfosis.

266. *Diosera longifolia* scapis radicatis: foliis oblongis.
Cum priore in paludosis.

POLYGYNIA.

126. MYOSURUS 394.

267. *MYOSURUS minimus*.
Dan. *Muse Rump.*
In maceris frequens.



Classis VI.

HEXANDRIA.
MONOGYNIA.* Flores *spathacei*.

127. NARCISSUS. *Cor.* supera 6 - petala. *Nectarium* infundibulif. 1 - phyllum. *Stam.* intra nectarium.
 128. ALLIUM. *Cor.* infera 6 - partita, patens. *Spatha* multiflora. *Umbella* congesta. *Capsf.* supera.

* * Flores *nudi*.

134. CONVALLARIA. *Cor.* infera 6 - fida. *Bacca* maculosa, 3 - locularis.
 135. HYACINTHUS. *Cor.* infera 6 - fida, campanulata. *Germinis* ad apicem pori 3 melliferi.
 132. ANTHERICUM. *Cor.* infera 6 - petala, patens. *Capsf.* ovata.
 131. ORNITHOGALUM. *Cor.* infera 6 - petala, erecta, persistens, supra medium patens. *Filamenta* alterna basi dilatata.
 133. ASPARAGUS. *Cor.* infera 6 - partita, erecta: petalis 3bus interioribus apice reflexis. *Bacca* 3 - locularis, 6 - sperma.
 129. LILIUM. *Corolla* infera hexapetala. *Petalis* basi incumbenibus.
 130. TULIPA. *Cor.* infera 6 - petala, campanulata. *Stylus* o.

Flores *apetali*.

136. ACORUS. *Spadix* cylindricus tectus flosculis. *Cor.* 6 - petalæ, nudæ. *Styl.* o. *Folliculi* 1 - spermi.
 137. JUNCUS. *Cal.* 6 - phyllus. *Cor.* o. *Capsf.* 1 - locularis.

138. PEPLIS. *Cal.* 12-fidus. *Petala* 6, calyci inserta, *Capsf.* 2-locularis.

D I G Y N I A.

† *Rumex digynus.*
Polygonum Persicaria.
Polygonum hydropiper.

T R I G Y N I A.

141. TRIGLOCHIN. *Cal* 3-phyllus. *Cor.* 3-petala, calyciformis. *Styl.* o. *Capsf.* basi dehiscens.
139. RUMEX. *Cal.* 3-phyllus. *Cor.* 3-petala, conniventia. *Sem.* 1. triquetrum.
140. SCHEUCHZERIA. *Cal.* 6-phyllus. *Cor.* o. *Styl.* o. *Capsf.* 3, inflatae, 1-spermæ.

P O L Y G Y N I A.

142. ALISMA. *Cal.* 3-phyllus. *Cor.* 3-petala. *Sem.* plura.



MONOGYNIA.

127. N A R C I S S U S. 403.

268. † NARCISSUS *pseudo-Narcissus* spatha uniflora: nectario campanulato erecto, crispo, æquante: petala ovata.

Dan. *Paaske-Lilie*.

In Insula *Amagria*.

128. A L L I U M. 409.

* Foliis caulinis planis.

269. † ALLIUM *sativum* caule planifolio bulbifero: radice composita: staminibus tricuspидatis.

Pauli Fl. Oec. 14.

270. ALLIUM *Scorodoprasum* caule planifolio bulbifero: foliis crenulatis: vaginis ancipitibus: staminibus tricuspидatis.

271. † ALLIUM *carinatum* caule planifolio bulbifero: staminibus subulatis.

* * Foliis caulinis teretibus.

272. † ALLIUM *Spherocephalum* caule teretifolio umbellifero: foliis semiteretibus: staminibus tricuspидatis corolla longioribus.

273. † ALLIUM *vineale* caule teretifolio bulbifero, staminibus tricuspидatis.

274. ALLIUM *oleraceum* caule tereti bulbifero: foliis scabris semiteretibus subtus fulcatis: staminibus simplicibus.

Dan. *Vild Løg*.

In Nemore raro; ad villam *Fionia Hindsgavl* frequens.

275. ALLIUM *Schænoprasum* scapo nudo adæquante
folia teretia filiformia.

Dan. *Græsløg*.

Raro; in littore villæ *Hindsgavl* passim.

* * * Foliis radicalibus, scapo nudo.

276. † ALLIUM *urfinum* scapo nudo semicylindræo;
foliis lanceolatis, petiolatis: umbella fastigiata.

Dan. *Rams*.

In Sylva prædii *Hindsgavl Fionia* nimis copiosum,
haud extirpandum.

Lac & carnem bovinam tanto ALLII sapore inficit, ut pro-
sus edi nequeant; post Festum vero Johannis impune in
locis pascuntur boves.

129. LILIUM. 410.

277. LILIUM *Martagon* foliis verticillatis: floribus
reflexis: corollis revolutis.

Ad hortum veterem quotannis sponte pervenit.

130. TULIPA. 415.

278. TULIPA *sylvestris* flore subnutante: foliis lanceo-
latis.

In Vivario semel inventa.

131. ORNITHOGALUM. 418.

279. ORNITHOGALUM *luteum* scapo anguloso diphy-
llo: pedunculis umbellatis simplicibus.

Dan. *Fugle-Melk*.

In vivario; in prato ad *Nyebroc* & in sylvestri ad
Dronninggaard.

280. ORNITHOGALUM *minimum* scapo angulato di-
phylo: pedunculis umbellatis ramosis.

Rarius,

281. * ORNITHOGALUM *nutans* floribus secundis pendulis: nectario stamineo campaniformi.
Totum hortum occupat.
Planta Neapolitana nullo modo extirpanda, nec extra eundem.

132. ANTHERICUM. 422.

282. † ANTHERICUM *ossifragum* foliis ensiformibus: filamentis lanatis. *Fl. Dan. t. 42. P. Fl. Oec. 21. Dan. Beenbrek.*
• In diœcesi Wiburgensi.

283. ANTHERICUM *ramosum* foliis planis: scapo ramoso: corollis planis: pistillo recto.
Dan. Ederkop - Urt.
In collibus ad molendina copiose floret planta hæc Auctore *Linnaeo* in orbe rarissima; in quibusdam individuis scapus simplex, minime ramosus.

133. ASPARAGUS. 424.

284. ASPARAGUS *officinalis* caule herbaceo erecto: foliis fetaceis: stipulis duabus interioribus, una exteriore. *P. Fl. Oec. 30.*
Dan. Aspargis.
Pharmac. ASPARAGI radix, femina.
In nemore ex horto forte illuc delata.

134. CONVALLARIA. 425.

285. CONVALLARIA *majalis* scapo nudo.
Dan. Lilie Konval.
Pharmac. LILII CONVALLIUM flores.
• In vivario raro.
286. † CONVALLARIA *verticillata* foliis verticillatis.
Fl. Dan. t. 86.

287. † CONVALLARIA *polygonatum* foliis alternis amplexicaulibus, caule ancipiti: pedunculis axillaribus subunifloris. *P. Fl. Oec. 55.*

Dan. *Værk-urt; Hvidroed; Bukkebar.*

Pharmac. SIGILLI SALOMONIS radix.

288. CONVALLARIA *multiflora* foliis alternis amplexicaulibus: caule tereti: pedunculis axillaribus multifloris. *Fl. Dan. t. 152.*

In alneto ad viam vulgo *Nyebroc.*

♂) - - - caule tereti: pedunculis axillaribus unifloris.

Dan. *Bukke - Bar; Salomons Segl.*

Hanc varietatem copiose legi in alneto pagi *Holte.*

289. CONVALLARIA *bifolia* foliis cordatis.

In nemorosis sterilibus vulgaris.

135. HYACINTHUS. 427.

290. † HYACINTHUS *botryoïdes* corollis globosis uniformibus: foliis canaliculato - cylindricis strictis.

136. ACORUS. 434.

291. ACORUS *calamus.* *Pauli Fl. Oec. p. 17.*

Dan. *Kalmus.*

Pharmac. CALAMI VULGARIS radix.

In paludosis ad amnem copiose.

137. JUNCUS. 437.

* Culmis *nudis.*

292. JUNCUS *conglomeratus* culmo nudo stricto: capitulo laterali.

Dan. *Glat Lyse - Sif.*

In uliginosis.

293. *JUNCUS effusus* culmo nudo stricto : panicula laterali.

Dan. *Skarp Lyse-Sif*.

In sylvestribus paludosis.

294. *JUNCUS filiformis* culmo nudo filiformi nutante : panicula laterali.

In paludosis turfosis.

295. * *JUNCUS squarrosus* culmo nudo : foliis setaceis : capitulis glomeratis aphyllis.

Morif. Hist. III. p. 228. t. 9. f. 13.

In turfosis frequens.

* * Culmis foliosis.

296. *JUNCUS articulatus* foliis nodoso-articulatis : petalis obtusis.

Dan. *Sif-Gras*, nomen plerisque commune.

In ripa.

297. *JUNCUS bulbosus* foliis linearibus canaliculatis : capsulis obtusis.

Morif. Hist. III. p. 227. t. 9. f. 11.

In ericetis.

298. *JUNCUS bufonius* culmo dichotomo : foliis angulatis : floribus solitariis sessilibus.

In inundatis.

299. *JUNCUS pilosus* foliis planis pilosis : corymbo ramofo.

In Sylvis.

300. *JUNCUS campestris* foliis planis subpilosis : spicis sessilibus pedunculatisque.

In ficcis.

138. PEPLIS. 446.

301. *PEPLIS Portula* floribus apetalis. *Fl. Dan. t. 64.*

Dan. *Vand-Portulac.*

In inundatis; copiosissime ad dextram, qua itur in villam,

TRIGYNIA.

TRIGYNIA.

139. RUMEX. 451.

* Floribus *Hermaphroditis*.

302. RUMEX *crispus* floribus hermaphroditis: valvulis integris graniferis: foliis lanceolatis, undatis, acutis.

Dan. *Krused - Skreppe*.

Ad aggeres.

303. RUMEX *maritimus* floribus hermaphroditis: valvulis dentatis graniferis: foliis linearibus.

Dan. *Strand - Skreppe*.

In littore maris.

304. RUMEX *acutus* floribus hermaphroditis: valvulis dentatis graniferis: foliis cordato-oblongis acuminatis. *P. Fl. Oec. 127.*

Dan. *Röd Skreppe; Rhabur*.

Pharmac. LAPATHI ACUTI radix, herba, semina.

In humidis.

305. RUMEX *aquaticus* floribus hermaphroditis: valvulis integerrimis nudis: foliis cordatis acutis.

In ripa.

* * Floribus *diclinis*.

306. RUMEX *acetosa* floribus dioicis: foliis oblongis sagittatis. *P. Fl. Oec. 126.*

Dan. *Syrer, Suur - Kaal*.

Pharmac. ACETOSÆ NOSTRATIS herba, radix.

In nemore ac in pratis.

307. RUMEX *acetosella* floribus dioicis: foliis lanceolato-hastatis.

In pratis ubique.

140. SCHEUCHZERIA. 452.

308. * SCHEUCHZERIA *palustris*. Fl. Dan. t. 76.
In infulis lacuum Bagsværd & Lyngbye frequens.

141. TRIGLOCHIN. 453.

309. TRIGLOCHIN *palustre* capsulis trilocularibus sublinearibus.
In pratis humidis.
310. TRIGLOCHIN *maritimum* capsulis sexlocularibus ovatis.
Dan. Strand - Siv - Gras.
In maritimis rarius.

POLYGYNIA.

142. ALISMA. 460.

311. ALISMA *plantago aquatica* foliis ovatis acutis : fructibus obtuse trigonis.
Dan. Gotberaad.
In paludosis vulgaris.
312. † ALISMA *natans* foliis ovatis obtusis : pedunculis solitariis. Act. Paris. 1719. t. 4. f. 8.
313. ALISMA *Ranunculoides* foliis lineari-lanceolatis ; fructibus globofo-squarrosis.
Fl. Dan. t. 122.
In foliis rarius.



Classis VII.

HEPTANDRIA.

MONOGYNIA.

43. TRIENTALIS. *Cal.* 7 - phyllus. *Cor.* 7 - partita
æqualis plana. *Bacca* 1 - locularis, sicca.

MONOGYNIA.

143. TRIENTALIS. 461.

14. TRIENTALIS *Europæa* foliis lanceolatis integerrimis. *Fl. Dan.* t. 84.

Dan. Vinter - Grøn.

In Sylva ac. in nemore vulgatissima.



Classis VIII.

OCTANDRIA
MONOGYNIA.

* Flores completi.

145. EPILOBIUM. Cor. 4 - petala. Cal. 4 - phyllus, superus. Caps. oblonga, 4 - locularis. Sem. papposa.
144. OENOTHERA. Cal. 4 - fidus. Petala 4. Caps. cylindrica, infera. Sem. nuda.
146. VACCINIUM. Cor. 1 - petala; Filamenta receptaculo inserta. Cal. 4 - dentatus, superus. Bacca 4 - locularis, polysperma.
147. ERICA. Cor. 1 - petala, 4 - fida. Cal. 4 - phyllus, inferus. Filament. receptaculo inserta. Anthera 2 - fidæ. Caps. 4 - locularis.

† *Monotropa Hypopithys.*

** Flores incompleti.

148. DAPHNE. Cal. o. Cor. 4 - fida, marcescens; Stamina includens. Bacca pulposa, 1 - sperma.

TRIGYNIA.

149. POLYGONUM. Cal. o. Cor. 5 - partita calycina. Sem. 1 - angulatum, nudum.

TETRAGYNIA.

151. ADOXA. Cor. 4 - seu 5 - fida, supera. Cal. 2 - fidus, inferus. Bacca 4 - seu 5 - locularis, calyci coalita.
152. ELATINE. Cor. 4 - petala. Cal. 4 - phyllus. Caps. 4 - locularis, 4 - valvis, depressa.
150. PARIS. Cor. 4 - petala, subulata. Cal. 4 - phyllus. Bacca 4 - locularis.

MONOGYNIA.

144. OENOTHERA. 469.

15. * OENOTHERA *biennis* foliis ovato - lanceolatis planis: caule muricato subvillosa.

In cespitosis horti Virginica planta sponte provenit; nunquam in vicinitate culta.

145. EPILOBIUM. 471.

* *Staminibus declinatis.*

16. EPILOBIUM *angustifolium* foliis sparsis linearilanceolatis: floribus inæqualibus. *P. Fl.* 63.

Dan. *Due - Urt.*

In margine agri prope *Othiniam Fionia* tria individua, in petris Molis *Hassia*, vulgo *Winterkastan*, numerosa offendi. Planta hortis digna.

** *Staminibus erectis regularibus: Petalis bifidis.*

17. EPILOBIUM *hirsutum* foliis oppositis lanceolatis ferratis decurrenti - amplexicaulibus.

In paludoso ad lacum *Bagsværd*.

18. EPILOBIUM *montanum* foliis oppositis ovatis dentatis.

In sylvestribus frequens.

19. EPILOBIUM *tetragonum* foliis lanceolatis denticulatis, imis oppositis: caule tetragono.

In sylvestribus rarius.

20. EPILOBIUM *palustre* foliis oppositis lanceolatis integerrimis: petalis emarginatis: caule erecto.

In humidiusculis.

146. VACCINIUM. 483.

* Foliis *annotinis deciduis*.

321. VACCINIUM *Myrtillus* pedunculis unifloris: foliis ovatis ferratis deciduis: caule angulato.

P. Fl. Oec. 149.

Dan. *Blaa-Bar*.

Pharmac. MYRTILLI baccæ.

In sylva ad lacum *Fuursøe*, & in alnetis.

322. VACCINIUM *uliginosum* pedunculis unifloris: foliis integerrimis ovalibus basi ciliatis.

Dan. *Bölle-Bar*.

In uliginosis.

* * Foliis *sempervirentibus*.

323. † VACCINIUM *Vitis idææ* racemis terminalibus nutantibus: foliis obovatis revolutis integerrimis subtus punctatis. Fl. Dan. t. 41. P. Fl. Oec. 151.

Dan. *Tytte-Bar*, *Kröfing-Bar*.

In sterilibus, sabulosis.

324. VACCINIUM *Oxycoccus* foliis integerrimis revolutis ovatis: caulibus repentibus filiformibus nudis. Fl. Dan. t. 80. P. Fl. Oec. 150.

Dan. *Trane-Bar*.In paludibus Sphagno obsitis in sylva vulgo *Hareskov*, ac lacus *Lyngbye*.

147. ERICA. 484.

325. ERICA *vulgaris* antheris bicornibus inclusis: corollis inæqualibus campanulatis mediocribus: foliis oppositis sagittatis. P. Fl. Oec. 66.

Dan. *Lyng*.

In campis sterilibus.

326. † ERICA *Tetralix* antheris bicornibus inclusis: corollis subglobosis confertis folio longioribus: foliis quaternis ciliatis patentibus. *Fl. Dan. t. 81.*

Copiose in ericetis inter *Apenrade & Flensburgum.*

327. † ERICA *cinerea* antheris bicornibus inclusis: corollis ovatis racemosis: foliis ternis glabris linearibus. *Fl. Dan. t. 39.*

148. D A P H N E. 485.

328. DAPHNE *Mezereum* floribus sessilibus ternis caulinis: foliis lanceolatis deciduis.

Pharmac. LAUREOLÆ cortex.

Rarissime, semel modo legi.

T R I G Y N I A.

149. P O L Y G O N U M. 495.

* Spica *unica.*

329. POLYGONUM *bistorta* caule simplicissimo, monostachyo: foliis ovatis in petiolum decurrentibus.

P. Fl. Oec. 109.

Dan. Slange - Urt.

* * Pistillo *bifido.*

330. POLYGONUM *amphibium* floribus pentandris semidigynis: staminibus corolla longioribus.

Dan. Pile - Urt.

In aquosis glabrum; in siccis scabrum.

331. POLYGONUM *hydropiper* floribus hexandris semidigynis: foliis lanceolatis: stipulis submuticis.

P. Fl. Oec. 110.

Dan. Vand - Peber; Bitter Pile - Urt.

Pharmac. PERSICARIÆ herba.

In subhumidis,

332. POLYGONUM *persicaria* floribus hexandris digynis: spicis ovato-oblongis: foliis lanceolatis: stipulis ciliatis.

In ruderatis.

* * * Foliis *indivisis*; Floribus *oſtandris*.

333. POLYGONUM *aviculare* floribus oſtandris trigynis axillaribus: foliis lanceolatis: caule procumbente herbaceo. *P. Fl. Oec.* 108.

Dan. *Vey-Græs*; *Hönſe-Græs*; *Hundrede-Knuder*, *Pharmac.* CENTUMNODIÆ herba.

In agris & alibi.

* * * * Foliis *subcordatis*.

334. POLYGONUM *Convolutulus* foliis cordatis: caule volubili: floribus planiusculis.

Dan. *Snerle*.

In agris aq in horto frequens.

TETRAGYNIA.

150. PARIS. 500.

335. PARIS *quadrifolia*. *Fl. Dan.* t. 132.

Dan. *Ulf-Bær*; *Fire-Blad*.

Pharmac. Baccæ herb. PARIDIS.

In prato alneo ad lacum *Bagsværd*; in insula *Thorsinge* passim in nemorosis.

151. ADOXA. 501.

336. ADOXA *Moschatellina*. *Fl. Dan.* t. 94.

Dan. *Desmer-Urt*.

In nemorosis passim. In horto regio *Fridrichsberg* copiose.

152. ELATINE. 502.

337. ELATINE *Hydropiper* foliis oppositis.

Fl. Dan. t. 156. *Vqill. Bot. Paris.* t. 2. f. 2.

In rivulo.

Classis IX.

ENNEANDRIA.

HEXAGYNIA.


153. BUTOMUS. *Cal. 0. Cor. 6-petala. Caps. 6.*
polyspermæ.

HEXAGYNIA.

153. BUTOMUS. 507.

338. BUTOMUS *umbellatus.*

In amne, qua hortum permanat.



Classis X.

DECANDRIA

MONOGYNIA.

Flores *polypetali aequales*.

154. MONOTROPA. *Cal.* 0. *Pet.* 10, quorum 5 exteriora basi excavato-mellifera. *Capsf.* 5-valvis, polysperma.

Quinta pars numeri excluditur omnibus lateralibus floribus.

158. PYROLA. *Cal.* 5-partitus. *Petala* 5. *Antherae* sursum bicornes. *Capsf.* 5-locularis, angulis dehiscens, polysperma.
155. LEDUM. *Cal.* 5-fidus. *Cor.* plana, 5-partita. *Capsf.* 5-locularis, basi dehiscens, polysperma.

Flores *monopetali aequales*.

156. ANDROMEDA. *Cal.* 5-partitus. *Cor.* ovata: ore 5-fido: *Capsf.* 5-locularis.
157. ARBUTUS. *Cal.* 5-partitus. *Cor.* ovata, ore basi pellucida. *Bacca* 5-locularis.

DIGYNIA.

161. SCLERANTHUS. *Cor.* 0. *Cal.* 5-fidus, superus. *Sem.* 2. calyci inclusa.
159. CHRYSOSPLENIUM. *Cor.* 0. *Cal.* 4 seu 5-fidus superus, coloratus. *Capsf.* 2-rostris, 1-locularis, polysperma.
160. SAXIFRAGA. *Cor.* 5-petala. *Cal.* 5-partitus. *Capsf.* 2-rostris, 1-locularis.

162. *SAPONARIA*. *Cor.* 5 - petala. *Pet.* unguiculata.
Cal. 1 - phyllus, nudus. *Capsf.* oblonga, 1 - locularis.

163. *DIANTHUS*. *Cor.* 5 - petala. *Pet.* unguiculata.
Cal. cylindricus, 1 - phyllus: basi squamis 4.
Capsf. cylindrica, 1 - locularis.

TRIGYNIA.

167. *ARENARIA*. *Capsf.* 1 - locularis, polysperma.
Petal. 5. integra patentia. *Cal.* 5 - phyllus, patens.

166. *STELLARIA*. *Capsf.* 1 - locularis, polysperma.
Petala 5 - bipartita, patentia. *Cal.* 5 - phyllus, patens.

164. *CUCUBALUS*. *Capsf.* 3 - locularis. *Petal.* 5, unguiculata: fauce nuda. *Cal.* inflatus.

165. *SILENE*. *Capsf.* 3 - locularis. *Pet.* 5. unguiculata: fauce coronata. *Cal.* ventricosus.

† *Tamarix Germanica*.

PENTAGYNIA.

168. *SEDUM*. *Capsf.* 5. ad nectaria. *Cor.* 5 - petala.
Squamæ nectariferæ 5, ad basin germinis. *Cal.* 5 - fidus.

173. *SPERGULA*. *Capsf.* 1 - locularis, 5 - valvis, ovata. *Pet.* 5, integra. *Cal.* 5 - phyllus.


172. *CERASTIUM*. *Capsf.* 1 - locularis, apice dehiscent. *Petala* bifida. *Cal.* 5 - phyllus.

170. AGROSTEMMA. *Capsf.* 1-locularis, oblonga. *Petala* 5, unguiculata: limbo obtuso, indiviso. *Cal.* 1-phyllus, tubulosus, coriaceus.

171. LYSHNIS. *Capsf.* 3-locularis, oblonga. *Pet.* 5, unguiculata: limbo bifido. *Cal.* 1-phyllus, oblongus, lævis.

169. OXALIS. *Capsf.* 5-locularis, 5-gona, angulis dehiscens. *Petala* unguibus connexa. *Cal.* 5-phyllus.

† Adoxa,



MONOGYNIA.

154. MONOTROPA. 536.

339. MONOTROPA *Hipopithys* floribus lateralibus octandris: terminali decandro. *Fl. Dan. t. 232.*

In vivario frequens; in Sylvis vulgo *Hareskov*, *Trueskov*, & in luco *Kalven* ad *Dronninggaard* sæpius legi.

Parasitica radiculæ fagi, uti conspicitur in icone optime *Fl. Dan.*, nullam ipsa radicem habens.

155. LEDUM. 546.

340. † LEDUM *palustre*.

Pharmac. RORISMARINI SYLVESTRIS herba.

156. ANDROMEDA. 549.

341. † ANDROMEDA *polifolia* pedunculis aggregatis: corollis ovatis: foliis alternis lanceolatis revolutis,

Fl. Dan. t. 54.

Dan. Vild Rosmarin.

Ad *Kollekolle* crescere quidem dicitur, mihi in paludibus ejus regionis assidue quovis tempore quærenti nunquam se obtulit.

157. ARBUTUS. 522.

342. † ARBUTUS *uva ursi* caulibus procumbentibus: foliis integerrimis. *Fl. Dan. t. 33. P. Fl. Oec. 23.*

Dan. Meelbar - Rths.

158. PYROLA. 554.

343. PYROLA *rotundifolia* staminibus adscendentibus: pistillo declinato. *Fl. Dan. t. 110.*

Dan. Vinter - Grön.

Pharmac. PYROLÆ herba.

In Insulis lacus *Lyngbye* paludosis.

18 *Decandria. Mono. - & Digynia.*

344. *PYROLA minor* floribus racemosis dispersis: staminibus pistillisque rectis. *Fl. Dan. t. 55.*

Locis umbrosis passim; in vivario, in Sylva *Hareskov* & in *Næffet*, ac in Insula *Kvæberholm* frequens.

345. *PYROLA secunda* racemo unilaterali.
In præcipitiis Sylvæ *Hareskov* frequens.

DIGYNIA.

159. CHRYSOSPLENIUM. 558.

346. *CHRYSOSPLENIUM alternifolium* foliis alternis.
Frequens in uliginosis Vivarii ac Sylvæ *Næffet*.

160. SAXIFRAGA. 559.

347. * *SAXIFRAGA Hirculus* foliis caulinis lanceolatis alternis nudis inermibus: caule erecto.
Fl. Dan. t. 200.

In prato paludoso *Jonstrup* frequens.

348. *SAXIFRAGA granulata* foliis reniformibus caulinis lobatis: caule ramoso: radice granulosa.
Dan. Steenbrek.

Pharmac. *SAXIFRAGÆ ALBÆ* radix sub nomine
seminum, herba, flores.

In arvis sterilibus frequens.

349. † *SAXIFRAGA tridactylites* foliis cuneiformibus trifidis alternis: caule erecto ramoso.

In *Amagria* frequens.

161. SCLERANTHUS. 562.

350. *SCLERANTHUS annuus* calycibus fructus patulis.
P. Fl. Oec. 133.

Dan. Tand-Urt; Knavel.

In agris.

51. *SCLERANTHUS perennis* calycibus fructus clausis.
In arvis siccis ac in cespitosis.

162. S A P O N A R I A. 564.

52. *SAPONARIA officinalis* calycibus cylindricis: foliis ovato-lanceolatis.
Extra hortum copiose.

163. D I A N T H U S. 565.

* Flores aggregati.

53. *DIANTHUS armeria* floribus aggregatis fasciculatis: squamis calycinis lanceolatis villosis, tubum æquantibus. *Fl. Dan. t. 230.*

Dan. *Vilde Nelliker.*

In arvis raro.

54. † *DIANTHUS prolifer* floribus aggregatis capitatis: squamis calycinis ovatis obtusis muticis tubum superantibus. *Fl. Dan. t. 201.*

In *Bornholmia*.

* * Flores solitarii.

55. *DIANTHUS deltoides* floribus solitariis: squamis calycinis lanceolatis binis: corollis crenatis.

In aridis, loco elatiori vulgo *Brankebakke*.

56. † *DIANTHUS superbus* floribus paniculatis: squamis calycinis brevibus acuminatis: corollis multifido-capillaribus: caule erecto.

57. † *DIANTHUS plumarius* floribus solitariis: squamis calycinis subovatis brevissimis: corollis multifidis fauce pubescentibus.

* * * Caule unifloro herbaceo.

58. *DIANTHUS arenarius* caulibus subunifloris: squa-

mis calycinis ovatis obtusis: corollis multifidis:
foliis linearibus.
In arenosis.

TRIGYNIA.

164. CUCUBALUS. 566.

359. † CUCUBALUS *bacciferus* calycibus campanulatis:
petalis distantibus: pericarpis coloratis: ramis di-
varicatis.

360. CUCUBALUS *Behen* calycibus subglobosis glabris
reticulato venosis: capsulis trilocularibus: corol-
lis subnudis.

In siccis vulgo *Deyligheden*.

165. SILENE. 567.

361. SILENE *nutans* petalis bifidis: floribus lateralibus
secundis cernuis: panicula nutante.

Dan. *Tiare-Urt*.

Passim in montosis, ad Molendinum ærarium fre-
quens.

362. † SILENE *armeria* floribus fasciculatis fastigiatis:
foliis superioribus cordatis glabris.

166. STELLARIA. 568.

363. STELLARIA *nemorum* foliis cordatis petiolatis:
panicula pedunculis ramosis.

In vivario, ac in luco ad lacum *Væreløse*.

364. STELLARIA *holostea* foliis lanceolatis serrulatis:
petalis bifidis.

Dan. *Oeyentröst*.

In Nemore.

365. STEL-

355. STELLARIA *graminea* foliis linearibus integerrimis; floribus paniculatis.
In sylvestribus vulgaris.

167. A R E N A R I A. 569.

366. ARENARIA *peploides* foliis ovatis acutis carnosis.
In littore maris.
367. * ARENARIA *trinervia* foliis ovatis acutis petiolatis, nervosis.
In Nemore.
368. ARENARIA *serpyllifolia* foliis subovatis acutis sessilibus: corollis calyce brevioribus.
In glareosis.
369. ARENARIA *rubra* foliis filiformibus: stipulis membranaceis vaginantibus.
Dan. *Kna - Gras*.
In arvis copiose.

P E N T A G Y N I A.

168. SEDUM. 579.

* * *Planifolia.*

370. † SEDUM *Telephium* foliis planiusculis serratis: corymbo folioso: caule erecto.
Dan. *St. Hans - Urt*; *Kraft - Urt*.
In sabulosis præcipitiis maritimis ad villam *Fionia Lundsgaard*.

* * *Teretifolia.*

371. † SEDUM *album* foliis oblongis obtusis teretiusculis sessilibus patentibus: cyma ramosa.
Fl. Dan. t. 66.

372. *SEDUM acre* foliis subovatis adnato - sessilibus
gibbis erectiusculis alternis: cyma trifida.

Dan. *Helleknop*.

Pharmac. SEDI MINORIS herba.

In sterilissimis fræquens.

373. *SEDUM sexangulare* foliis subovatis adnato - sessilibus, gibbis erectiusculis sexfariam imbricatis.

In campis lîceis.

169. O X A L I S. 582.

* *Scapo radicali.*

374. *OXALIS acetosella* scapo unifloro: foliis ternatis:
radice squamoso - articulata. *P. Fl. Oec.* 104.

• Dan. *Suur - Klever*; *Giöge - Mad*.

Pharmac. ACETOSELLÆ herba.

In Sylvestribus.

* * *Foliis caulinis alternis.*

375. † *OXALIS corniculata* caule ramoso diffuso: pedunculis umbelliferis.

170. AGROSTEMMA. 583.

376. *AGROSTEMMA githago* hirsuta, calycibus corollam æquantibus: petalis integris nudis.

Pauli Fl. Oec. 9.

Dan. *Klinte*.

• Inter Secale.

171. L Y C H N I S. 584.

377. *LYCHNIS flos cuculi* petalis quadrifidis: fructu subrotundo.

Dan. *Hofmod*; *Græs - Nellike*.

In pratis humidis frequens.

378. LYCHNIS *viscaria* petalis integris.

Dan. *Tiare - Urt*; *Beeg - Urt*.

In pratis elevatis, *Brankebakke*.

In quibusdam flosculis quatuor modo inveniuntur pistilla.

379. LYCHNIS *dioica* floribus dioicis.

Passim; in prato ad lacum *Bagfverd* vulgaris, flore purpureo.

172. CERASTIUM. 585.

* Capfulis *oblongis*.

380. CERASTIUM *vulgatum* foliis ovatis: petalis calyci æqualibus: caulibus diffusis.

Vaill. Bot. Paris. t. 30. f. 1.

In ruderatis.

381. * CERASTIUM *viscosum* erectum villoso-viscosum. *Vaill. Bot. Paris. t. 30. f. 3.*

In maceritiis.

382. CERASTIUM *semidetandrum* floribus pentandris: petalis emarginatis. *Vaill. Bot. Paris. t. 20. f. 2.*

In pascuis.

383. CERASTIUM *arvense* foliis lineari-lanceolatis acutis glabris: corollis calyce majoribus.

Vaill. Bot. Paris. t. 30. f. 4.

In arenosis.

* * Capfulis *subrotundis*.

384. CERASTIUM *aquaticum* foliis cordatis sessilibus: floribus solitariis: fructibus pendulis.

In umbrosis subhumidis.

173. SPERGULA. 586.

385. SPERGULA *arvensis* foliis verticillatis: floribus decandris.

P. Fl. Oec. 140.

Dan. Kna - Gras.

In arvis frequens.

* SPERGULA *pentandra* foliis verticillatis: floribus pentandris.

In arvis rarior. Num sola præcedentis varietas?

386. SPERGULA *nodosa* foliis oppositis subulatis levibus: caulibus simplicibus.

Fl. Dan. t. 96.

In campis depressis.



Classis XI.

DODECANDRIA.

MONOGYNIA.

174. ASARUM. *Cor.* o. *Cal.* 3 - seu 4 - fidus, superus. *Caps.* 6 - seu 8 - locularis, coriacea, coronata.
 175. LYTHRUM. *Cor.* 6 - petala, calyci inserta. *Cal.* 12 - fidus, inferus. *Caps.* 2 - locularis, polysperma.

DIGYNIA.

176. AGRIMONIA. *Cor.* 5 - petala, *Cal.* 5 - dentatus, altero obvallatus, *Sem.* 1. & 2. in fundo calycis.

TRIGYNIA.

177. RESEDA. *Corolla* petala laciniata. *Cal.* 1 - phyllus, partitus. *Caps.* 1 - locularis, ore dehiscens.
 178. EUPHORBIA. *Cor.* 4 - seu 5 - petala. *Petalis* peltatis. *Cal.* 1 - phyllus, ventricosus. *Caps.* 3 - cocca.

DODECAGYNIA.

179. SEMPERVIVUM. *Cor.* 12 - petala. *Cal.* 12 - partitus. *Caps.* 12, polysperma.



MONOGYNIA.

174. ASARUM. 589.

387. † ASARUM *Europæum* foliis reniformibus obtusis
binis.

Pharmac. ASARI radix, folia, femina.

175. LYTHRUM. 604.

388. LYTHRUM *Salicaria* foliis oppositis cordato-lanceolatis: floribus spicatis dodecandris,

Dan. *Katte - Urt.*

Ad ripam amnis in Nemore frequens.

DIGYNIA.

176. AGRIMONIA. 607.

389. AGRIMONIA *Eupatoria* foliis caulinis pinnatis:
impari sessili: fructibus hispidis.

Dan. *Agermaane.*

Pharmac. AGRIMONIE herba.

In pratis apicis.

TRIGYNIA.

177. RESEDA. 608.

390. RESEDA *Luteola* foliis lanceolatis integris: calycibus quadrifidis.

P. Fl. Oec. 122.

Dan. *Vau.*

In arvis & ad sepes.

178. EUPHORBIA. 609.

391. EUPHORBIA *Helioscopia umbella quinquefida* : trifida : dichotoma, involucellis obovatis : foliis cuneiformibus ferratis.

Dan. *Ulve-Melk*; *Fandens-Melk*.

Pharmac. ESULÆ herba.

In cultis frequens.

392. EUPHORBIA *Peplus umbella trifida* : dichotoma, involucellis ovatis : foliis integerrimis obovatis petiolatis.

Dan. *Fugle-Melk*; *Vorte-Urt*.

In oleraceis.

DODECAGYNIA.

179. SEMPERVIVUM. 612.

393. SEMPERVIVUM *tectorum foliis ciliatis* : propagibus patentibus.

Dan. *Huus-Lög*.

Pharmac. SEMPERVIVI MAJORIS herba.

In tectis tuguriorum.



Classis XII.

ICOSANDRIA.

MONOGYNIA.

180. PRUNUS. *Cal.* inferus, 5-fidus. *Cor.* 5-petala. *Drupe* nux futuris prominatis.

DIGYNIA.

181. CRATEGUS. *Cal.* superus, 5-fidus. *Cor.* 5-petala. *Bacca* infera, 2-sperma.

TRIGYNIA.

182. SORBUS. *Cal.* superus, 5-fidus. *Cor.* 5-petala. *Bacca* infera 3-sperma.

PENTAGYNIA.

183. MESPILUS. *Cal.* superus, 5-fidus. *Cor.* 5-petala. *Bacca* infera, 5-sperma.
 184. PYRUS. *Cal.* superus 5-fidus. *Cor.* 5-petala. *Pomum* inferum, 5-loculare, polyspermum.
 185. SPIRÆA. *Cal.* inferus, 5-fidus. *Cor.* 5-petala. *Capsule* congestæ, polyspermæ.

POLYGYNIA.

186. ROSA. *Cal.* 5-fidus, urceolatus, carnosus, collo coarctatus. *Cor.* 5-petala. *Sem.* plurima, hispida, calycis interiori lateri affixa.

87. RUBUS. *Cal.* 5 - fidus. *Cor.* 5 - petala. *Bacca* composita acinis monospermis.
88. TORMENTILLA. *Cal.* 8 - fidus. *Cor.* 4 - petala. *Sem.* 8, mutica, subrotunda, nuda, receptaculo parvo exsucco affixa.
88. FRAGARIA. *Cal.* 10 - fidus. *Cor.* 5 - petala. *Sem.* plurima supra receptaculum baccatum, decidua.
89. POTENTILLA. *Cal.* 10 - fidus. *Cor.* 5 - petala. *Sem.* plurima mutica, subrotunda, nuda, receptaculo parvo exsucco affixa.
91. GEUM. *Cal.* 10 - fidus. *Cor.* 5 - petala. *Sem.* plurima, caudata, pilosa.
92. COMARUM. *Cal.* 10 - fidus. *Cor.* 5 - petala, calyce minora. *Recept.* seminum ovatum, spongiosum, persistens.

† Spiræa filipendula.
Spiræa Ulmaria.



MONOGYNIA.

180. PRUNUS. 620.

394. PRUNUS *Padus* floribus racemosis: foliis deciduis, basi subtus biglandulosis.

Fl. Dan. t. 205. P. Fl. Oec. 115.

Dan. Hage-Bær; vilde Kirse-Bær.

In dumeto extra septum, vulgo Deyligheden.

395. PRUNUS *spinosa* pedunculis solitariis: foliis lanceolatis glabris: ramis spinosis. *P. Fl. Oec. 116.*

Dan. Slaaen-Torn.

Pharmac. ACACIÆ NOSTRÆ cortex, flores, fructus.
Ubique vulgaris.

DIGYNIA.

181. CRATÆGUS. 622.

396. † CRATÆGUS *Aria* foliis ovatis inæqualiter serratis: subtus tomentosis.

Dan. Axel-Bær; Bornholms Rosiner.

397. CRATÆGUS *Oxyacantha* foliis obtusis subtrifidis serratis. *P. Fl. Oec. 58.*

Dan. Hage-Torn; Havn-Torn; Hvid-Torn.
Rassim.

TRIGYNIA.

182. SORBUS. 623.

398. SORBUS *aucuparia* foliis pinnatis utrinque glabris.
P. Fl. Oec. 139.

Dan. Røn.

Pharmac. SORBI AUCUPARIÆ baccæ.

In Sylvis frequens.

PENTAGYNIA.

183. MESPILUS. 625.

329. † MESPILUS *Cotoneaster inermis*: foliis ovatis integerrimis. *Fl. Dan. t. 112.*

184. PYRUS. 626.

400. PYRUS *malus* foliis serratis: umbellis fessilibus. *F. Fl. Oec. 118.*

Dan. Skov-Æble.

In Nemore ad paludem magnam.

185. SPIRÆA. 630.

401. SPIRÆA *filipendula* foliis pinnatis: foliolis uniformibus serratis: caule herbaceo: floribus cymosis.

Dan. Bierg-Mandel; Bønder-Muskat.

Pharmac. SAXIFRAGIÆ RUBRÆ radix, herba.

In septo Deyligheden ac in colle Brankbakke vulgaris.

402. SPIRÆA *Ulmaria* foliis pinnatis: impari majore lobato: floribus cymosis.

Dan. Mød-Urt; Giedde-Skiæg.

Pharmac. ULMARIÆ radix, herba.

Cum præcedente vulgaris.

POLYGYNIA.

186. ROSA. 631.

403. ROSA *canina* germinibus ovatis pedunculisque glabris: caule petiolisque aculeatis.

Pharmac. ROSÆ SYLVESTRIS flores, fungi CYNOSBATI f. Bedeguar.

Vulgaris.

404. ROSA *spinosissima* germinibus globosis glabris: pedunculis, caule, petiolisque aculeatissimis.

Ad sepes minus frequens.

405. ROSA *Eglanteria* germinibus globosis pedunculisque glabris: caule aculeis sparsis rectis: petiolis scabris: foliolis acutis.

Dan. *Engel-Torn*; *Hyven-Torn*,
Vulgaris.

In dumetis nec rara; rarissima in Suecia planta.

187. RUBUS. 632.

* *Frutescentes.*

406. RUBUS *idæus* foliis quinato-pinnatis ternatisque: caule aculeato: petiolis canaliculatis.

Pharmac. RUBI IDÆI fructus.

Passim.

407. RUBUS *castus* foliis ternatis subnudis: lateralibus bilobis: caule tereti aculeato.

Dan. *Kor-Bar*.

In dumetis passim.

408. RUBUS *fruticosus* foliis quinato-digitatis ternatisque: caule petiolisque aculeatis.

Dan. *Bramber*.

In sepibus.

* * *Herbacei.*

409. RUBUS *saxatilis* foliis ternatis: flagella nudis repantibus herbaceis. *Fl. Dan. t. 134.*

Dan. *Frue-Bar*.

Inter lapides rarus.

410. † RUBUS *Chamamorus* foliis simplicibus lobatis: caule inermi unifloro. *Fl. Dan. t. 1.*

Dan. *Multe-Bar*.

Pharmac. CHAMÆMORI fructus conditi.

188. FRAGARIA 633.

411. FRAGARIA *vesca* flagellis reptans.

Dan. *Jord-Bar.*

Pharmac. FRAGARIÆ herba, radix, fructus.

In Sylvaticis passim.

189. POTENTILLA 634.

* Foliis *pinnatis*.

412. POTENTILLA *anserina* foliis pinnatis serratis;
caule repente. *P. Fl. Oec.* 113.

Dan. *Sölv-Urt*; *Gaafe-Urt.*

Pharmac. ANSERINÆ herba.

Ad vias.

413. † POTENTILLA *rupestris* foliis pinnatis alternis;
foliolis quinis ovatis, crenatis: caule erecto.

* * Foliis *digitatis*.

414. POTENTILLA *argentea* foliis quinatis cuneiformibus incisus subtus tomentosis: caule erecto.

Ad margines agrorum.

415. POTENTILLA *verna* foliis radicalibus quinatis acute serratis retusis: caulinis ternatis: caule declinato.

In collibus asperis aprisis; in *Brankebakke* vulgaris.

416. POTENTILLA *reptans* foliis digitatis: caule repente: pedunculis unifloris.

Pharmac. PENTAPHYLLI radix, herba.

In campo ac horto.

417. † POTENTILLA *Norvegica* foliis ternatis: caule dichotomo: pedunculis axillaribus.

Fl. Dan. t. 171.

190. TORMENTILLA. 635.

418. TORMENTILLA *erecta*, caule erectiusculo: foliis sessilibus. P. Fl. Oec. 144.

Dan. *Tormentil*; *Blod-Roed*.

Pharmac. TORMENTILLÆ radix, herba.

In pascuis sterilibus.

191. GEUM. 636.

419. GEUM *urbanum* floribus erectis: fructu globoso villoso: aristis uncinatis nudis: foliis lyratis.

P. Fl. Oec 77.

Dan. *Nellike-Roed*; *Benedikt-Urt*.

Pharmac. CARYOPHYLLATÆ radix, herba.

In dumetis.

420. GEUM *rivale* floribus nutantibus: fructu oblongo: aristis plumosis.

In pratis humidis.

192. COMARUM. 638.

421. COMARUM *palustre*.

In uliginosis frequens.



Classis XIII.

POLYANDRIA: MONOGYNIA.

Flores tetrapetali.

195. PAPAVER. *Cal.* 2 - phyllus. *Cor.* 4 - petala.
Capsf. 1 - locularis, coronata; sub stigmate persi-
stente poris dehiscens.
194. CHELIDONIUM. *Cal.* 2 - phyllus. *Cor.* 4 - pe-
tala. *Siliqua* 1 - locularis, linearis.
193. ACTÆA. *Cal.* 4 - phyllus. *Cor.* 4 - petala. *Bacca*
1 - locularis. *Sem.* semiorbiculata, gemino ordine.

Flores pentapetali.

198. CISTUS. *Cal.* 5 - phyllus: *Foliolis* duobus ma-
noribus. *Cor.* 5 - petala. *Capsf.* subrotunda.
197. THIA. *Cal.* 5 - partitus, deciduus. *Cor.* 5 - pe-
tala. *Bacca* sicca, globosa, 5 - locularis, 5 - val-
vis, basi dehiscens, 1 - sperma.
† Delphinium Consolida.

Flores polypetali.

196. NYMPHÆA. *Cal.* 4 - seu 5 - phyllus, magnus.
Cor. polypetala. *Bacca* multilocularis, corticosa,
truncata.

TRIGYNIA.

199. DELPHINIUM. *Cal.* 0. *Cor.* 5 - petala: supre-
mo cornuto. *Nectarium* bifidum, sessile. *Siliqua*
3. f. 1.
200. ACONITUM. *Cal.* 0. *Cor.* 5 - petala: suprema
galeato, *Nect.* 2 - pedunculata, recurva.

PENTAGYNIA.

201. AQUILEGIA. *Cal.* 0. *Cor.* 5 - petala. *Nectaris* 5, corniculata inter petala. *Capsf.* 5, distinctæ.

HEXAGYNIA.

202. STRATIOTES. *Spatha* diphylla. *Perianth.* 3-6-
dum, *Petala* 3. *Bacca* 6-locularis, infera.

POLYGYNIA.

204. THALICTRUM. *Cal.* 0. *Cor.* 4 - petala. *Sem.* plurima, submutica, nuda.
207. CALTHA. *Cal.* 0. *Cor.* 5 - petala. *Capsf.* plures, polyspermæ.
206. TROLLIUS. *Cal.* 0. *Cor.* circiter 14 - petala, *Capsf.* plurimæ, ovatæ, monospermæ.
203. ANEMONE. *Cal.* 0. *Cor.* 6 - 9 - petala, *Sem.* plurima.

† A. Hepatica. *Cal.* 3 - phyllus.

205. RANUNCULUS. *Cal.* 5 - phyllus. *Cor.* 5 - petala, intra ungues poro mellifero. *Sem.* plurima.



MONO.

MONOGYNIA.

193. A C T Æ A. 644.

422. ACTÆA *spicata* racemo ovato: fructibus baccatis;
Pauli Fl. Oec. 5.
 Dan. *Drue - Munke; Christophers - Urt.*
 In nemore ac sylvis passim.

194. CHELIDONIUM. 647.

423. CHELIDONIUM *majus* pedunculis umbellatis.
 Dan. *Svale - Urt.*
Pharmac. CHELIDONII MAJORIS herba, radix.
 In maceriis frequens.
424. † CHELIDONIUM *glaucium* pedunculis unifloris;
 foliis amplexicaulibus sinuatis: caule glabro.

195. P A P A V E R. 648.

* Capsulis *hispidis*.

425. PAPAVER *Argemone* capsulis clavatis hispidis;
 caule folioso multifloro.
 Dan. *Valdmue.*
 In agris.

* * Capsulis *glabris*.

426. PAPAVER *dubium* capsulis oblongis glabris: caule
 multifloro setis adpressis: foliis pinnatifidis incisis.
 Inter segetes.
427. PAPAVER *Rhæas* capsulis glabris globosis: caule
 piloso multifloro: foliis pinnatifidis incisis.
 Dan. *Korn - Rose.*
Pharmac. PAPAVERIS RHÆADOS herba, flores.
 Inter segetes.

G

428. * *PAPAVER somniferum* calycibus capsulisque glabris: foliis amplexicaulibus incis. *ac*

Pharm. PAPAVERIS ALBI & NIGRI semina, capitula. In septo *Deyligheden*, in macerie horti veteris, *ac* in horto frequens.

Tota glabra, parte caulis inter fructificationem & supremum folium fetis raris prostantibus.

196. N Y M P H Æ A. 653.

429. *NYPHÆA lutea* foliis cordatis integerrimis: calyce petalis majore, pentaphyllo.

Dan. *Aakande*; *Kierlinge - Kruus*.

In amne ac lacubus frequens.

430. *NYPHÆA alba* foliis cordatis integerrimis: calyce quadrifido.

Pharmac. NYPHÆÆ ALBÆ radix, flores.

Cum præcedente vulgaris.

197. T I L I A. 660.

431. *TILIA Europæa* floribus nectario destitutis.

P. Fl. Oec. 143.

Dan. *Lind*.

Pharmac. TILIÆ flores, cortex interior, carbones. In nemoroso *Cathrinenholm*.

198. C I S T U S. 673.

432. *CISTUS Helianthemum* suffruticosus procumbens: stipulis lanceolatis: foliis oblongis revolutis subpilosis. *Fl. Dan. t.* 101.

In aridis passim.

TRIGYNIA.

199. DELPHINIUM. 681.

33. DELPHINIUM *Consolida nectariis monophyllis*;
caule subdiviso. *P. Fl. Oec.* 59.

Dan. *Ridder - Spore.*

Pharmac. CONSOLIDÆ REGALIS flores.

In agris raro.

200. ACONITUM. 682.

34. † ACONITUM *Napellus* foliorum laciniis linearibus
superne latioribus linea exaratis.

Fl. Dan. t. 123. *Pauli Fl. Oec.* 3.

Dan. *Storm - Hat*; *Munke - Hætte.*

PENTAGYNIA.

201. AQUILEGIA. 684.

35. AQUILEGIA *vulgaris* nectariis incurvis.

Dan. *Akkeley.*

Pharmac. AQUILEGIÆ herba, flores, femina.

In nemorosis raro; in hortis frequens.

HEXAGYNIA.

202. STRATIOTES. 687.

36. STRATIOTES *Aloides* foliis ensiformi-triangularibus
ciliato-aculeatis.

Dan. *Vand - Aloë.*

In amne ac paludosis.

POLYGYNIA,

203. ANEMONE. 694.

* Flore calyculato.

437. ANEMONE *Hepatica* foliis trilobis integerrimis.
Dan. *Adel-Klefver*.

Pharmac. HEPATICÆ NOBILIS herba, flores.

In umbrosis nemoris ac Sylvæ *Hareskov* frequens.

* * Pedunculo involucrato.

438. ANEMONE *Pulsatilla* pedunculo involucrato: petalis rectis: foliis bipinnatis.

Fl. Dan. t. 153. Paul. Fl. Oec. 18.

Dan. *Koe-Bielde; Oxe-Oere.*

Pharmac. PULSATILLÆ herba.

In colle aprico.

439. * ANEMONE *pratensis* pedunculo involucrato: petalis apice reflexis: foliis bipinnatis.

In aridis ad lacum *Fuursjøe Farum* versus vulgarissima.

440. † ANEMONE *vernalis* pedunculo involucrato: foliis pinnatis: flore erecto. *Fl. Dan. t. 29.*

* * * Flore nudo.

441. ANEMONE *nemorosa* feminibus acutis: foliolis incisis: caule unifloro. *Pauli Fl. Oec. 17.*

Dan. *Hvid-Sippe; Hvide-Ved.*

Pharmac. RANUNCULI ALBI flores.

In nemorosis vulgaris.

442. † ANEMONE *Ranunculoïdes* feminibus acutis: foliolis incisis: petalis subrotundis: caule subbifloro. *Fl. Dan. t. 140.*

In prato nemoroso ad lacum *Vareløse* frequens; in luco *Ermelund* vulgarissima.

204. THALICTRUM. 697.

13. † THALICTRUM *minus* foliis sexpartitis : floribus cernuis.

In fabuloso ad diverforium *Kiöge - Kroe.*

Folia quædam sexpartita, quædam tripartita. *Petala* 4. purpurefcentia. *Stamina* 14. 16. vel 18. *Pistilla* 3. 4. vel 5.

14. THALICTRUM *flavum* caule folioso sulcato : panicula multiplici erecta.

Dan. *Knepi - Panden.*

In subhumidis pratis.

15. † THALICTRUM *aquilegifolium* fructibus pendulis triangularibus rectis : caule tereti.

205. RANUNCULUS. 699.

* Foliis *simplicibus*.

46. RANUNCULUS *flammula* foliis ovato-lanceolatis petiolatis : caule declinato.

Dan. *Hane - Fod.*

In pascuis udis frequens.

47. RANUNCULUS *reptans* foliis linearibus : caule repente. *Fl. Dan. t. 108.*

In ripa lacus *Lyngbye - Söe.*

48. RANUNCULUS *Lingua* foliis lanceolatis : caule erecto.

In lacu *Lyngbye - Söe.*

49. † RANUNCULUS *amplexicaulis* foliis ovatis acuminatis amplexicaulibus : caule multifloro : radice fasciculata.

50. RANUNCULUS *Ficaria* foliis cordatis angulatis petiolatis.

Pharmac. CHELIDONII MINORIS radix, herba.

Vulgaris ubique primo vere.

* * Foliis dissectis & divisis.

451. RANUNCULUS *auricomus* foliis radicalibus reniformibus, crenato-incisis: caulinis digitatis linearibus: caule multifloro. B.
In pascuis subhumidis.

452. RANUNCULUS *sceleratus* foliis inferioribus palmatis, summis digitatis: fructibus oblongis.
In paludibus & ad ripam amnis.

453. RANUNCULUS *bulbosus* calycibus retroflexis: pedunculis fulcatis: caule erecto: foliis compositis.
In pascuis vulgaris.

454. RANUNCULUS *repens* calycibus patulis: pedunculis fulcatis: stolonibus repentibus: foliis compositis.
In oleraceis.

455. † RANUNCULUS *polyanthemos* calycibus patulis: pedunculis fulcatis: caule erecto: foliis multipartitis.

456. RANUNCULUS *acris* calycibus patulis: pedunculis teretibus: foliis tripartito-multifidis: summis linearibus. P. Fl. Oec. 120.

Dan. Smör - Urt; Krage - Taer; Hane - Fod.

457. RANUNCULUS *arvensis* feminibus aculeatis: foliis superioribus decompositis linearibus.
In agris.

458. † RANUNCULUS *hederaceus* foliis subrotundis trilobis integerrimis: caule repente.

459. RANUNCULUS *aquatilis* foliis submersis capillaribus: emerfis subpeltatis.

In paludosis Seminarii Deyligheden.

In horto offendi loco elatiori ab aquosis remotissimo: foliis omnibus capillaribus.

206. TROLLIUS. 700.

460. TROLLIUS *Europæus* corollis conniventibus: nectariis longitudine staminum.

Fl. Dan. t. 133.

Dan. Eng - Blomme ; Kabbeleje.

In pratis subhumidis ad lacum *Bagvaard*, & ad turguria *Ravnebusene*.

207. CALTHA. 703.

461. CALTHA *palustris*. *P. Fl. Oec.* 42.

Dan. Smør - Urt ; Koe - Blomme.

In uliginosis vulgaris.



Classis XIV.

DIDYNAMIA.

GYMNOSPERMIA.

* Calyces *subquinquefidi*.

219. LEONURUS. *Anthera* glandulis nitidis ossis ad-
spersæ.
213. GLECOMA. *Anthearum* singulum par in for-
mam crucis connivens. *Calyx* 5-fidus.
212. MENTHA. *Filamenta* distantia, recta. *Corolla*
subæqualis; 4-fida: lacinia latiore emarginata.
208. TEUCRIUM. *Corollæ* labium superius (nullum)
ultra basin bipartitum, divaricatum pro egressu
staminum.
210. AJUGA. *Corollæ* labium superius minimum. *Sta-*
mina labio superiore longiora.
211. BETONICA. *Corollæ* labium superius adscen-
dens planiusculum, tubus cylindricus. *Calyx* ari-
status.
214. LAMIUM. *Corollæ* Labium superius integrum,
fornicatum; labium inferius bilobum; *faux* utrin-
que margine dentata.
215. GALEOPSIS. *Corollæ* labium superius crenatum,
fornicatum; labium inferius supra bi-dentatum.
216. STACHYS. *Corollæ* labium sup. fornicatum;
lab. inferius lateribus reflexum: lacinia interme-
dia majore emarginata. *Stamina* deflorata versus
latera reflexa.
209. NÉPETA. *Corollæ* lab. inferius lacinula interme-
dia crenata. *Faux* margine reflexo. *Stamina* ap-
proximata.

217. **BALLOTA.** *Cal.* hypocrateriformis, 5-dentatus, 10-striatus. *Corollæ* lab. sup. crenatum, concavum.
218. **MARRUBIUM.** *Cal.* hypocrateriformis rigidus, 10-striatus. *Cor.* lab. sup. 2-fidum; lineare, rectum.

* *Calyces bilabiati.*

223. **SCUTELLARIA.** *Calyx* ore integro post florescentiam clauso, operculato.
222. **THYMUS.** *Calycis* bilabiati faux villis clausa.
224. **PRUNELLA.** *Filamenta* bifurca, altero apice antherifera.
221. **ORIGANUM.** *Strobilus* tetragonus, spicatus, calyces colligens.
220. **CLINOPODIUM.** *Involucrum* multifetum verticillo subjectum, calyces colligens.

ANGIOSPERMIA.

Calyces quadrifidi.

228. **LATHRÆA.** *Cal.* 4-fidus. *Cor.* personata: glandula depressa ad basin futuræ germinis. *Caps.* 1-locularis.
226. **EUPHRASIA.** *Cal.* 4-fidus, cylindricus. *Cor.* personata. *Antheræ* inferiores altero tubo basi spinosæ.
225. **RHINANTHUS.** *Cal.* 4-fidus, ventricosus. *Cor.* personata. *Caps.* 2-locularis, obtusa, compressa.
227. **MELAMPYRUM.** *Cal.* 4-fidus. *Coroll.* personata: lab. superius compressum, margine replicato. *Caps.* 2-locularis, obliqua, dehiscens. *Semina* 2-gibba.

Calyces quinquefidi.

230. LIMOSELLA. *Cal.* 5 - fidus. *Cor.* 5 - fida, æqualis. *Stam.* per paria approximata. *Caps.* 1 - locularis, 2 - valvis, polysperma.
231. SCROPHULARIA. *Cal.* 5 - fidus. *Cor.* subglobosa, resupinata. Labii segmento intermedio interno. *Caps.* 2 - locularis.
232. ANTIRRHINUM. *Cal.* 5 - phyllus. *Corolla* personata basi deorsum prominens, nectarifera. *Caps.* 2 - locularis.
229. PEDICULARIS. *Cal.* 5 - fidus. *Cor.* personata. *Caps.* 2 - locularis, mucronata, obliqua. *Semina* tunicata.



GYMNOSPERMIA.

208. AJUGA. 705.

462. AJUGA *pyramidalis* tetragono - pyramidalis.
Fl. Dan. t. 185. Pauli Fl. Oec. 12.
Pharmac. CONSOLIDÆ MEDIÆ herba.
 463. † AJUGA *reptans* stolonibus reptantibus.

209. TEUCRIUM. 706.

464. † TEUCRIUM *Scordium* foliis oblongis sessilibus
 dentato - ferratis: floribus geminis lateralibus pe-
 dunculatis: caule diffuso.
Dan. Skordium, Lögs - Urt.
Pharmac. SCORDII herba.
 In pratis humidis.

210. NEPETA. 710.

465. NEPETA *Cataria* floribus spicatis: verticillis sub-
 pedicellatis: foliis petiolatis cordatis dentato - fer-
 ratis.
Dan. Kette - Urt.
 In ruderatis.

211. BETONICA. 718.

466. † BETONICA *officinalis* spica interrupta: corolla-
 rum lacinia labii intermedia emarginata.
Dan. Betonie.
Pharmac. BETONICÆ herba, flores.

212. MENTHA. 713.

* *Spicata.*

467. † MENTHA *viridis* spicis oblongis: foliis lanceo-
 latis nudis ferratis sessilibus: staminibus corolla
 longioribus.
Dan. Mynthe.

468. † MENTHA *sylvestris* spicis oblongis: foliis oblongis tomentosis ferratis sessilibus: staminibus corolla longioribus. *Spec. Plant.* p. 804.

* * Capitata.

469. MENTHA *aquatica* spicis capitatis: foliis ovatis ferratis petiolatis: staminibus corolla longioribus.

Dan. *Heste-Myntbe.*

Ad ripam amnis ac lacuum frequens.

470. MENTHA *arvensis* floribus verticillatis: foliis ovatis acute ferratis: staminibus corollam æquantibus. *Morif Hist.* III. p. 369. t. 7. f. 5.

Pharmac. MENTHÆ SYLVESTRIS herba.

In pratis humidis agrisque vulgaris.

471. † MENTHA *pulegium* floribus verticillatis: foliis ovatis obtusis subcrenatis: caulibus subteretibus repentibus: staminibus corolla longioribus.

Dan. *Vand-Poley.*

213. GLECOMA. 714.

472. GLECOMA *hederacea* foliis reniformibus crenatis. *P. Fl. Oec.* 78.

Dan. *Jord-Vedbende.*

Pharmac. HEDERÆ TERRESTRIS herba.

Ad sepes & alibi vulgaris.

214. LAMIUM. 716.

473. LAMIUM *album* foliis cordatis acuminatis petiolatis: verticillis vigintifloris.

Dan. *Döv-Nelde.*

Pharmac. LAMII ALBI herba.

In cultis.

474. *LAMIUM purpureum* foliis cordatis obtusis petiolatis.

In cultis.

475. *LAMIUM amplexicaule* foliis floralibus sessilibus amplexicaulibus obtusis.

Dan. *Katte-øyne*.

In arvis ac hortis vulgaris.

215. GALEOPSIS. 717.

476. *GALEOPSIS Ladanum* internodiis caulinis æqualibus: verticillis omnibus remotis.

Dan. *Hampe-Nelde*.

In arvis fabulosis.

477. *GALEOPSIS Tetrabit* internodiis caulinis superne incrassatis: verticillis summis subcontiguis.

β, *GALEOPSIS* corolla flava: labio inferiore maculato.

Inter olera & segetes frequens; β, in agris *Bagvard* vulgatissima.

478. *GALEOPSIS Galeobdolon* verticillis sexfloris: involucro tetraphyllo.

In nemorosis foliis.

216. STACHYS. 719.

479. *STACHYS sylvatica* verticillis sexfloris: foliis cordatis petiolatis.

In nemore ad hortum ac in foveis ad maceriem frequens.

480. *STACHYS palustris* verticillis sexfloris: foliis lineari-lanceolatis amplexicaulibus.

In agris subhumidis.

481. *STACHYS arvensis* verticillis sexfloris: caule debili: foliis obtusis nudiusculis: corollis calyce vix longioribus.

217. BALLOTA. 720.

482. *BALLOTA nigra* foliis cordatis indivisis ferratis :
calycibus acuminatis.

Ad pagum *Wirum*.

218. MARRUBIUM. 721.

483. *MARRUBIUM vulgare* dentibus calycinis setaceis
uncinatis.

Dan. *Marrube*.

Pharmac. MARRUBII ALBI herba.

In ruderatis.

219. LEONURUS. 722.

484. *LEONURUS Cardiaca* foliis caulinis lanceolato-tri-
lobis.

Dan. *Hierte - Span; Sey-Urt*.

Pharmac. CARDIACÆ herba.

In ruderatis.

220. CLINOPODIUM. 725.

485. *CLINOPODIUM vulgare* capitulis subrotundis hi-
spidis : bracteis setaceis.

In collibus aridis.

221. ORIGANUM. 726.

486. *ORIGANUM vulgare* spicis subrotundis panicula-
tis conglomeratis : bracteis calyce longioribus
ovatis. *P. Fl. Oec.* 103.

Dan. *Vild-Meyran; Toft; Konning*.

Pharmac. ORIGANI VULGARIS herba.

In montosis sylvaticis.

222. T H Y M U S. 727.

187. THYMUS *Serpyllum* floribus capitatis: caulibus repentibus: foliis planis obtusis basi ciliatis.

P. Fl. Oec. 142.

Dan. Vild-Thymian; Vor - Frues Sengebalm.

Pharmac. SERPYLLI herba.

In ficcis apricis frequens.

188. THYMUS *Acinos* verticillis sexfloris: caulibus erectis subramosis: foliis acutis ferratis.

Dan. Vild Basilikum.

In aridis rarior.

223. S C U T E L L A R I A. 734.

189. SCUTELLARIA *galericulata* foliis cordato-lanceolatis crenatis, floribus axillaribus.

Dan. Feber - Urt.

Ad ripas amnis ac paludum frequens.

224. P R U N E L L A. 735.

190. PRUNELLA *vulgaris* foliis omnibus ovato-oblongis petiolatis.

In pascuis frequens.

ANGIOSPERMIA.

225. RHINANTHUS. 740.

191. RHINANTHUS *crista galli*, corollarum labio superiore compresso brevior.

Dan. Hane - Kamm.

In pratis ficcis.

226. EUPHRASIA. 741.

492. EUPHRASIA *officinalis* foliis ovatis lineatis argute dentatis.

Dan. Oeyen - Tröst.

Pharmac. EUPHRASIE herba.

In sterilibus ubique.

493. EUPHRASIA *Odontites* foliis linearibus: omnibus ferratis.

In agris rarior.

227. MELAMPYRUM. 742.

494. MELAMPYRUM *cristatum* spicis quadrangularibus: bracteis cordatis compactis denticulatis imbricatis.

Dan. Koe - Huede.

Passim, in colle ad Molendinum ærarium frequentius.

495. † MELAMPYRUM *arvense* spicis conicis laxis: bracteis dentato - setaceis. P. Fl. Oec. 98.

Dan. Koe - Huede; Ræve - Rumpe.

Inter segetes.

In Præfectura *Hafniensi* exulat; in *Fridericoburgensi* numero Secale fere superat.

496. MELAMPYRUM *nemorosum* floribus secundis lateralibus, bracteis dentatis, cordato - lanceolatis: summis coloratis sterilibus: calycibus lanatis.

In prato nemoroso *Bagssvard*, & ad Molendina vulgaris.

497. MELAMPYRUM *pratense* floribus secundis lateralibus: conjugationibus remotis: corollis clausis.

In pratis.

498. MELAMPYRUM *sylvaticum* floribus secundis lateralibus: conjugationibus remotis: corollis hiantibus. Fl. Dan. t. 145.

In nemore.

228. LAZ

228. LATHRÆA. 743.

229. † LATHRÆA *Squamaria* caule simplicissimo: corollis pendulis: labio inferiore trifido.

Fl. Dan. t. 134.

Dan. Tand - Urt.

Kylling quidem olim *Fridrichsdal* locum huic natalem indicit; jam vero mihi assiduissime quærenti legere non contigit.

229. PEDICULARIS. 746.

230. PEDICULARIS *palustris* caule ramoso: calycibus calloso-punctatis: corollis labio obliquis.

Dan. Luus - Urt; Eng - Troid.

In pratis paludosis frequens ad lacus *Lyngbye & Bagsværd.*

231. PEDICULARIS *sylvatica* caule ramoso: calycibus oblongis angulatis lævibus: corollis labio cordato.

Fl. Dan. t. 225.

Cum præcedente passim.

230. ANTIRRHINUM. 750.

232. ANTIRRHINUM *Linaria* foliis lanceolato-linearibus confertis: caule erecto: spicis terminalibus sessilibus: floribus imbricatis.

P. Fl. Oec. 22.

Dan. Vild - Hör: Torske - Mund.

Pharmac. LINARIÆ herba.

In ruderalis ac in ipsa area Villæ.

233. ANTIRRHINUM *minus* foliis plerisque alternis lanceolatis obtusis: caule ramosissimo diffuso.

In campis rarior.

504. † ANTIRRHINUM *Orontium* corollis ecaudatis
floribus subspicatis: calycibus digitatis corolla lo-
gioribus.

Pharmac. ANTIRRHINI herba.

231. SCROPHULARIA. 756.

505. SCROPHULARIA *nodosa* foliis cordatis trinervi-
tis: caule acutangulo.

Dan. Orme - Urt; Bruun - Roed.

Pharmac. SCROPHULARIÆ radix.

Ad maceries ac in foveis haud infrequens.

506. † SCROPHULARIA *aquatica* foliis cordatis petio-
latis decurrentibus obtusis: caule membranis an-
gulato: racemis terminalibus.

232. LIMOSELLA. 776.

507. LIMOSELLA *aquatica*.

Fl. Dan. t. 69.

In ripa amnis ad molendinum, ac in piscina sylva
frequens.



Classis XV.

ETRADYNAMIA.

SILICULOSÆ.

Silicula integra nec apice emarginata.

- DRABA. *Silicula* integra, ovali-oblonga: *Valvulis* planiusculis, dissepimento parallelis. *Stylus* nullus.
- LUNARIA. *Silicula* integra, elliptica, compresso-plana: *Valvis* dissepimento æqualibus, parallelis, planis. *Stylus* exsertus.
- SUBULARIA. *Silicula* integra, ovata, *Valvis* ovatis, concavis, dissepimento contrariis. *Stylus* silicula brevior.
- MYAGRUM. *Silicula* stylo conico terminata persistente: *Valvulis* concavis.

Silicula apice emarginata.

- IBERIS. *Silicula* polysperma, emarginata. *Cor.* irregularis: petalis 2 exterioribus majoribus.
- COCHLEARIA. *Silicula* emarginata, turgida, scabra: *Valvulis* gibbis, obtusis.
- LEPIDIUM. *Silicula* emarginata cordata, polysperma: *Valvulis* carinatis, contrariis.
- THLASPI. *Silicula* emarginata, obcordata, polysperma: *Valvulis* navicularibus, marginatis, carinatis.

SILIUOSÆ.

Calyx clausus: Foliolis longitudinaliter conniventibus.

249. RAPHANUS. *Cal.* clausus. *Siliqua* torosa & articulata, teres. *Glandula* melliferæ 2 interamina breviora & pistillum, totidem inter staminibus longiora & calycem.
244. ERYSIMUM. *Cal.* clausus. *Siliqua* columnaris exacte tetraëdra.
245. ARABIS. *Glandula* nectariferæ 4, singulæ intercalycis foliola squamæ instar reflexæ. *Stigma* simplex.
247. BRASSICA. *Cal.* erectus connivens. *Sem.* globosa. *Glandula* inter stamina breviora & pistillum interque longiora & calycem.
246. TURRITIS. *Cal.* erectus, connivens. *Cor.* ætæta. *Siliqua* longissima, angulata.
241. DENTARIA. *Cal.* longitudinaliter connivens. *Stigma* emarginatum. *Siliqua* elastice diffiliens. *Valvulis* revolutis.

Calyx hians: Foliolis superne distantibus.

252. CRAMBE. *Siliqua* decidua, globosa, sicco-baccata. *Filamenta* 4, apice bifurca: altero antherifero.
251. ISATIS. *Siliqua* lanceolata, 1-locularis, 1-sperma, decidua, bivalvis. *Valvulis* navicularibus.
250. BUNIAS. *Siliqua* decidua, tetraëdra, angulis inæqualibus acuminatis muricata.
242. CARDAMINE. *Siliqua* elastice diffiliens, *Valvulis* revolutis. *Stigma* integrum. *Cal.* subhians.
248. SINAPIS. *Siliqua* dehiscens. *Cal.* pateris. *Corolla* unguis rectæ. *Glandula* inter stamina breviora & pistillum interque longiora & calycem.
243. SISYMBRIUM. *Siliqua* dehiscens valvulis reclinatis. *Calyx* patens. *Corolla* patula.

SILICULOSA.

233. MYAGRUM. 796.

08. MYAGRUM *sativum* filiculis obovatis pedunculatis, polyspermis.

Dan. *Gier*; *Vild - Hör*.

In agris.

09. MYAGRUM *paniculatum* filiculis lentiformibus orbiculatis punctato-rugosis. *Fl. Dan. t. 204.*

In agris haud infrequens.

234. SUBULARIA. 799.

10. † SUBULARIA *aquatica*. *Fl. Dan. t. 35.*

In lacustribus fabulosis.

235. DRABA. 800.

* *Caule nudo.*

11. DRABA *verna* scapis nudis: foliis lanceolatis subincisis.

Dan. *Gæslinge - Blomster*.

In apricis aridis ubique.

236. LEPIDIUM. 801.

12. LEPIDIUM *sativum* floribus tetradynamiis: foliis oblongis multifidis.

Pharmac. NASTURTII semen.

Ad hortum.

13. LEPIDIUM *ruderales* floribus diandris apetalis: foliis radicalibus dentato-pinnatis: ramiferis linearibus integerrimis. *Fl. Dan. t. 184.*

In ruderales.

237. THLASPI. 802.

- § 14. THLASPI *arvense* filiculis orbiculatis: foliis oblongis, dentatis, glabris. *P. Fl. Oec.* 142.

Dan. Penge - Urt.

Pharmac. THLASPEOS femina.

In agris.

- § 15. THLASPI *campestre* filiculis subrotundis: foliis sagittatis dentatis incanis.

In arvis apricis.

- § 16. THLASPI *Bursa pastoris* filiculis obcordatis: foliis radicalibus pinnatifidis.

Dan. Hyrde - Taske.

Pharmac. BURSÆ PASTORIS herba.

In cultis ubique.

238. COCHLEARIA. 803.

- § 17. COCHLEARIA *Danica* foliis hastatis angulatis.

Fl. Dan. t. 100.

Ad littora maris.

- § 18. † COCHLEARIA *officinalis* foliis radicalibus subrotundis; caulinis oblongis sublinuatis.

Fl. Dan. t. 135. *P. Fl. Oec.* 53.

Dan. Cochlearie; Skiörbugs - Urt.

Pharmac. COCHLEARIÆ herba.

- § 19. † COCHLEARIA *Coronopus* foliis pinnatifidis.

Fl. Dan. t. 202.

- § 20. † COCHLEARIA *Armoracia* foliis radicalibus lanceolatis crenatis: caulinis incis.

Dan. Vild Peber - Roed.

Pharmac. ARMORACIÆ radix.

239. IBERIS. 804.

- § 21. IBERIS *nudicaulis* herbacea foliis sinuatis: caule nudo simplici.

Dan. Steenkarse.

In arenosis.

240. LUNARIA. 809.

22. † LUNARIA *annua* foliis oppositis.

SILIKUOSA.

241. DENTARIA. 811.

523. DENTARIA *bulbifera* foliis inferioribus pinnatis :
summis simplicibus.In nemore; raro. * In præcipitiis promontorii Insulæ
Mona vulgaris.

242. CARDAMINE. 812.

524. CARDAMINE *impatiens* foliis pinnatis incisfis sti-
pulatis: floribus apetalis.*Morif. Hist. III. p. 221. t. 4. f. 1.*In umbrosis ad ripam, in Sylva *Nasset*.525. CARDAMINE *hirsuta* foliis pinnatis: floribus te-
trandris. *Fl. Dan. t. 148.*

In nemore ad ripam hortum versus, & alibi passim.

526. CARDAMINE *pratensis* foliis pinnatis: foliolis ra-
dicalibus subrotundis: caulinis lanceolatis.*Dan. Enge - Karse.*

In pascuis aquosis ubique.

527. CARDAMINE *amara* foliis pinnatis: axillis stolo-
niferis. *P. Fl. Occ. 44.**Dan. Vand - Karse.*

In aquosis umbrosis passim.

243. SISYMBRIUM. 813.

* *Siliquis declinatis brevibus.*528. SISYMBRIUM *Nasturtium aquat.* siliquis decli-
natis: foliis pinnatis: foliolis subcordatis.*Dan. Vand - Karse.**Pharmac. NASTURTHI AQUATICI herba.*

Minus frequens.

529. SISYMBRIUM *sylvestre* filiquis declinatis: foliis pinnatis: foliolis lanceolatis ferratis.

In humidis agrorum.

530. SISYMBRIUM *amphibium* filiquis declinatis oblongo - ovatis: foliis pinnatifidis ferratis.

Dan. *Vand - Peberroed*.

In palustribus.

* * Caule nudo.

531. † SISYMBRIUM *arenosum* caule subfoliofo ramofo: foliis lyratis rectangulo - dentatis hispidis: pilis ramosis.

* * * Foliis pinnatis.

532. SISYMBRIUM *Sophia* petalis calyce minoribus: foliis decomposito - pinnatis.

Pharmac. SOPHIÆ femina.

In maceriis passim.

244. ERYSIMUM. 814.

533. † ERYSIMUM *officinale* filiquis spicæ appressis: foliis runcinatis.

Dan. *Vild Senip*.

Pharmac. ERYSIMI herba.

534. ERYSIMUM *Barbarea* foliis lyratis: extimo subrotundo. Dan. *Vinter - Karse*.

Ad ripas raro.

535. ERYSIMUM *Alliaria* foliis cordatis.

Dan. *Hvidlögs - Urt*.

Pharmac. ALLIARIÆ herba.

In umbrosis frequens.

536. † ERYSIMUM *Cheiranthoides* foliis lanceolatis integerrimis.

245. A R A B I S. 818.

537. † ARABIS *Alpina* foliis amplexicaulibus dentatis.
Fl. Dan. t. 62.

538. ARABIS *Thaliana* foliis petiolatis lanceolatis integerrimis.
In aridis.

246. T U R R I T I S. 819.

539. TURRITIS *glabra* foliis radicalibus dentatis hispidis: caulinis integerrimis amplexicaulibus glabris.

Dan. *Vild Kaal.*

In siccis glareosis.

540. TURRITIS *hirsuta* foliis omnibus hispidis, caulinis amplexicaulibus.

In pascuis sylvaticis.

247. B R A S S I C A. 820.

541. BRASSICA *campestris* radice cauleque tenui: foliis caulinis uniformibus cordatis sessilibus.

Dan. *Vild Kaal.*

Inter Segetes.

248. S I N A P I S. 821.

542. SINAPIS *arvensis* filiquis multangulis torosoturgidis rostro longioribus. P. Fl. Oec. 136.

Dan. *Ager - Kaal.*

In agris.

543. SINAPIS *nigra* filiquis glabris apice tetragonis.

Dan. *Senip.*

Pharmac. SINAPIS femina.

In agris.

H 5

249. R A P H A N U S. 822.

544. RAPHANUS *Raphanistrum* filiquis teretibus articulatis lævibus unilocularibus. *P. Fl. Oec.* 121.

Dan. *Ager - Kaal*.

Inter hordeum.

250. B U N I A S. 823.

545. † BUNIAS *Cakile* filiculis ovatis lævibus ancipitibus.

Dan. *Strand - Karse*.

In sabuloso maritimo ad villas *Fionie Lundsgaard* & *Hindsgavl*.

251. I S A T I S. 824.

546. † ISATIS *tinctoria* foliis radicalibus crenatis: caulinis sagittatis: filiculis oblongis.


P. Fl. Oec. 86.

252. C R A M B E. 825.

547. CRAMBE *maritima* foliis cauleque glabris.

Dan. *Strand - Kaal*.

In littore maris.



Classis XVI.


MONADELPHIA.

DECANDRIA.

253. GERANIUM. *Monogyna. Stigmata 5. Caps.*
rostrata, 5 - cocca.

POLYANDRIA.

255. MALVA. *Calyx duplex: exterior 3 - phyllus.*
Caps. loculis 1 - spermis, verticillatis.
254. ALTHÆA. *Calyx duplex: exterior 9 - fidus.*
Caps. loculis 1 - spermis, verticillatis.



504. † ANTIRRHINUM *Orontium* corollis ecaudatis
floribus subspicatis: calycibus digitatis corolla lon-
gioribus.

Pharmac. ANTIRRHINI herba.

231. SCROPHULARIA. 756.

505. SCROPHULARIA *nodosa* foliis cordatis trinerve-
tis: caule acutangulo.

Dan. Orme - Urt; *Bruun - Roed.*

Pharmac. SCROPHULARIÆ radix.

Ad maceries ac in foveis haud infrequens.

506. † SCROPHULARIA *aquatica* foliis cordatis petio-
latis decurrentibus obtusis: caule membranis ad-
gulato: racemis terminalibus.

232. LIMOSELLA. 776.

507. LIMOSELLA *aquatica*.

Fl. Dan. t. 69.

In ripa amnis ad molendinum, ac in piscina sylva
frequens.



Classis XV.

ETRADYNAMIA.

SILICULOSÆ.

Silicula integra nec apice emarginata.

5. DRABA. *Silicula* integra, ovali-oblonga: *Valvulis* planiusculis, dissepimento parallelis. *Stylus* nullus.
6. LUNARIA. *Silicula* integra, elliptica, compresso-plana: *Valvis* dissepimento æqualibus, parallelis, planis. *Stylus* exsertus.
4. SUBULARIA. *Silicula* integra, ovata, *Valvis* ovatis, concavis, dissepimento contrariis. *Stylus* silicula brevior.
3. MYAGRUM. *Silicula* stylo conico terminata persistente: *Valvulis* concavis.

Silicula apice emarginata.

39. IBERIS. *Silicula* polysperma, emarginata. *Cor.* irregularis: petalis 2 exterioribus majoribus.
38. COCHLEARIA. *Silicula* emarginata, turgida, scabra: *Valvulis* gibbis, obtusis.
36. LEPIDIUM. *Silicula* emarginata cordata, polysperma: *Valvulis* carinatis, contrariis.
37. THLASPI. *Silicula* emarginata, obcordata, polysperma: *Valvulis* navicularibus, marginatis, carinatis.

SILIUOSÆ.

Calyx clausus: Foliolis longitudinaliter conniventibus.

249. RAPHANUS. *Cal.* clausus. *Siliqua* torosa f. articulata, teres. *Glandula* melliferæ 2 inter stamina breviora & pistillum, totidem inter stamina longiora & calycem.
244. ERYSIMUM. *Cal.* clausus. *Siliqua* columnari exacte tetraëdra.
245. ARABIS. *Glandula* nectariferæ 4, singulæ inter calycis foliola squamæ instar reflexæ. *Stigma* simplex.
247. BRASSICA. *Cal.* erectus connivens. *Sem.* globosa. *Glandula* inter stamina breviora & pistillum interque longiora & calycem.
246. TURRITIS. *Cal.* erectus, connivens. *Cor.* æcta. *Siliqua* longissima, angulata.
241. DENTARIA. *Cal.* longitudinaliter connivens. *Stigma* emarginatum. *Siliqua* elastice diffiliens. *Valvulis* revolutis.

Calyx hians: Foliolis superne distantibus.

252. CRAMBE. *Siliqua* decidua, globosa, sicco-baccata. *Filamenta* 4, apice bifurca: altero antherifero.
251. ISATIS. *Siliqua* lanceolata, 1-locularis, 1-sperma, decidua, bivalvis. *Valvulis* navicularibus.
250. BUNIAS. *Siliqua* decidua, tetraëdra, angulis inæqualibus acuminatis muricata.
242. CARDAMINE. *Siliqua* elastice diffiliens, *Valvulis* revolutis. *Stigma* integrum. *Cal.* subhians.
248. SINAPIS. *Siliqua* dehiscens. *Cal.* patens. *Cor.* unguis rectæ. *Glandula* inter stamina breviora & pistillum interque longiora & calycem.
243. SISYMBRIUM. *Siliqua* dehiscens valvulis ætiusculis. *Calyx* patens. *Corolla* patula.

SILICULOSA.

233. MYAGRUM. 796.

08. MYAGRUM *sativum* filiculis obovatis pedunculatis, polyspermis.

Dan. *Gier*; *Vild - Hör*.

In agris.

09. MYAGRUM *paniculatum* filiculis lentiformibus orbiculatis punctato-rugosis. *Fl. Dan. t. 204.*

In agris haud infrequens.

234. SUBULARIA. 799.

10. † SUBULARIA *aquatica*. *Fl. Dan. t. 35.*

In lacustribus sabulosis.

235. DRABA. 800.

* *Caule nudo.*

11. DRABA *verna* scapis nudis: foliis lanceolatis subincisis.

Dan. *Gæslinge - Blomster*.

In apricis aridis ubique.

236. LEPIDIUM. 801.

12. LEPIDIUM *sativum* floribus tetradynamiis: foliis oblongis multifidis.

Pharmac. NASTURTII semen.

Ad hortum.

13. LEPIDIUM *ruderales* floribus diandris apetalis: foliis radicalibus dentato-pinnatis: ramiferis linearibus integerrimis. *Fl. Dan. t. 184.*

In ruderatis.

H 3.

237. THLASPI. 802.

- §14. THLASPI *arvense* filiculis orbiculatis: foliis oblongis, dentatis, glabris. *P. Fl. Oec.* 142.

Dan. *Penge-Urt.*

Pharmac. THLASPEOS femina.

In agris.

- §15. THLASPI *campestre* filiculis subrotundis: foliis sagittatis dentatis incanis.

In arvis apricis.

- §16. THLASPI *Bursa pastoris* filiculis obcordatis: foliis radicalibus pinnatifidis.

Dan. *Hyrde-Taske.*

Pharmac. BURSÆ PASTORIS herba.

In cultis ubique.

238. COCHLEARIA. 803.

- §17. COCHLEARIA *Danica* foliis hastatis angulatis.

Fl. Dan. t. 100.

Ad littora maris.

- §18. † COCHLEARIA *officinalis* foliis radicalibus subrotundis; caulinis oblongis sublinuatis.

Fl. Dan. t. 135. *P. Fl. Oec.* 53.

Dan. *Cochlearie*; *Skiörbugs-Urt.*

Pharmac. COCHLEARIÆ herba.

- §19. † COCHLEARIA *Coronopus* foliis pinnatifidis.

Fl. Dan. t. 202.

- §20. † COCHLEARIA *Armoracia* foliis radicalibus lanceolatis crenatis: caulinis incis.

Dan. *Vild Peber-Roed.*

Pharmac. ARMORACIÆ radix.

239. IBERIS. 804.

- §21. IBERIS *nudicaulis* herbacea foliis sinuatis: caulibus nudo simplic.

Dan. *Steenkarse.*

In arenosis.

240. L U N A R I A. 809.

22. † LUNARIA *annua* foliis oppositis.

SILIKUOSA.

241. D E N T A R I A. 811.

23. DENTARIA *bulbifera* foliis inferioribus pinnatis :
summis simplicibus.

In nemore; raro. * In præcipitiis promontorii Insulæ
Mona vulgaris.

242. C A R D A M I N E. 812.

24. CARDAMINE *impatiens* foliis pinnatis incisfis sti-
pulatis: floribus apetalis.

Morif. Hist. III. p. 221. t. 4. f. 1.

In umbrosis ad ripam, in Sylva *Nasset*.

25. CARDAMINE *hirsuta* foliis pinnatis: floribus te-
trandris. *Fl. Dan. t. 148.*

In nemore ad ripam hortum versus, & alibi passim.

26. CARDAMINE *pratensis* foliis pinnatis: foliolis ra-
dicalibus subrotundis: caulinis lanceolatis.

Dan. Enge - Karse.

In pascuis aquosis ubique.

27. CARDAMINE *amara* foliis pinnatis: axillis stolo-
niferis. *P. Fl. Occ. 44.*

Dan. Vand - Karse.

In aquosis umbrosis passim.

243. S I S Y M B R I U M. 813.

* *Siliquis declinatis brevibus.*

28. SISYMBRIUM *Nasturtium aquat.* siliquis decli-
natis: foliis pinnatis: foliolis subcordatis.

Dan. Vand - Karse.

Pharmac. NASTURTH AQUATICI herba.

Minus frequens.

H 4

529. *SISYMBRIUM sylvestre* filiquis declinatis: foliis pinnatis: foliolis lanceolatis ferratis.

In humidis agrorum.

530. *SISYMBRIUM amphibium* filiquis declinatis oblongo-ovatis: foliis pinnatifidis ferratis.

Dan. *Vand - Peberroed*.

In palustribus.

* * Caule nudo.

531. † *SISYMBRIUM arenosum* caule subfolioso ramo: foliis lyratis rectangulo-dentatis hispidis: pilis ramosis.

* * * Foliis pinnatis.

532. *SISYMBRIUM Sophia* petalis calyce minoribus: foliis decomposito-pinnatis.

Pharmac. SOPHIÆ femina.

In maceriis passim.

244. ERYSIMUM. 814.

533. † *ERYSIMUM officinale* filiquis spicæ appressis: foliis runcinatis.

Dan. *Vild Senip*.

Pharmac. ERYSIMI herba.

534. *ERYSIMUM Barbaraa* foliis lyratis: extimo subrotundo. Dan. *Vinter - Karse*.

Ad ripas raro.

535. *ERYSIMUM Alliaria* foliis cordatis.

Dan. *Hvidlögs - Urt*.

Pharmac. ALLIARIÆ herba.

In umbrosis frequens.

536. † *ERYSIMUM Cheiranthoides* foliis lanceolatis integerrimis.

245. A R A B I S. 818.

537. † ARABIS *Alpina* foliis amplexicaulibus dentatis.
Fl. Dan. t. 62.

538. ARABIS *Thaliana* foliis petiolatis lanceolatis integerrimis.
In aridis.

246. T U R R I T I S. 819.

539. TURRITIS *glabra* foliis radicalibus dentatis hispidis: caulinis integerrimis amplexicaulibus glabris.

Dan. *Vild Kaal.*

In siccis glareosis.

540. TURRITIS *hirsuta* foliis omnibus hispidis, caulinis amplexicaulibus.

In pascuis sylvaticis.

247. B R A S S I C A. 820.

541. BRASSICA *campestris* radice cauleque tenui: foliis caulinis uniformibus cordatis sessilibus.

Dan. *Vild Kaal.*

Inter Segetes.

248. S I N A P I S. 821.

542. SINAPIS *arvensis* filiquis multangulis torosoturgidis rostro longioribus. P. Fl. Oec. 136.

Dan. *Ager - Kaal.*

In agris.

543. SINAPIS *nigra* filiquis glabris apice tetragonis.

Dan. *Senip.*

Pharmac. SINAPIS femina.

In agris.

H 5

226. EUPHRASIA. 741.

492. EUPHRASIA *officinalis* foliis ovatis lineatis argute dentatis.

Dan. Oeyen - Tröst.

Pharmac. EUPHRASIE herba.

In sterilibus ubique.

493. EUPHRASIA *Odontites* foliis linearibus: omnibus ferratis.

In agris rarior.

227. MELAMPYRUM. 742.

494. MELAMPYRUM *cristatum* spicis quadrangularibus: bracteis cordatis compactis denticulatis imbricatis.

Dan. Koe - Huede.

Passim, in colle ad Molendinum ærarium frequentius.

495. † MELAMPYRUM *arvense* spicis conicis laxis: bracteis dentato - fetaceis. P. Fl. Oec. 98.

Dan. Koe - Huede; Rave - Rumpé.

Inter segetes.

In Præfectura *Hafniensi* exulat; in *Fridericoburgensi* numero Secale fere superat.

496. MELAMPYRUM *nemorosum* floribus secundis lateralibus, bracteis dentatis, cordato - lanceolatis: summis coloratis sterilibus: calycibus lanatis.

In prato nemoroso *Bagsværd*, & ad Molendina vulgaris.

497. MELAMPYRUM *pratense* floribus secundis lateralibus: conjugationibus remotis: corollis clausis.

In pratis.

498. MELAMPYRUM *sylvaticum* floribus secundis lateralibus: conjugationibus remotis: corollis hiantibus. Fl. Dan. t. 145.

In nemore.

228. LATHRÆA. 743.

299. † LATHRÆA *Squamaria* caule simplicissimo: corollis pendulis: labio inferiore trifido.

Fl. Dan. t. 134.

Dan. Tand - Urt.

Kylling quidem olim *Fridrichsdal* locum huic natalem indicit; jam vero mihi assiduissime quærenti legere non contigit.

229. PEDICULARIS. 746.

500. PEDICULARIS *palustris* caule ramoso: calycibus calloso-punctatis: corollis labio obliquis.

Dan. Luus - Urt; Eng - Troid.

In pratis paludosis frequens ad lacus *Lyngbye & Bugsværd.*

501. PEDICULARIS *sylvatica* caule ramoso: calycibus oblongis angulatis lævibus: corollis labio cordato.

Fl. Dan. t. 225.

Cum præcedente passim.

230. ANTIRRHINUM. 750.

502. ANTIRRHINUM *Linaria* foliis lanceolato-linearibus confertis: caule erecto: spicis terminalibus sessilibus: floribus imbricatis.

P. Fl. Oec. 22.

Dan. Vild - Hör: Torske - Mund.

Pharmac. LINARIÆ herba.

In ruderalis ac in ipsa area Villæ.

503. ANTIRRHINUM *minus* foliis plerisque alternis lanceolatis obtusis: caule ramosissimo diffuso.

In campis rarior.

504. † ANTIRRHINUM *Orontium* corollis ecaudatis floribus subspicatis: calycibus digitatis corolla longioribus.

Pharmac. ANTIRRHINI herba.

231. SCROPHULARIA. 756.

505. SCROPHULARIA *nodosa* foliis cordatis trinerviis: caule acutangulo.

Dan. Orme - Urt; *Bruun - Roed.*

Pharmac. SCROPHULARIÆ radix.

Ad maceries ac in foveis haud infrequens.

506. † SCROPHULARIA *aquatica* foliis cordatis petiolatis decurrentibus obtusis: caule membranis angulato: racemis terminalibus.

232. LIMOSELLA. 776.

507. LIMOSELLA *aquatica*.

Fl. Dan. t. 69.

In ripa amnis ad molendinum, ac in piscina sylva frequens.



Classis XV.

ETRADYNAMIA.

SILICULOSÆ.

Silicula integra nec apice emarginata.

- 1. DRABA. *Silicula* integra, ovali-oblonga: *Valvulis* planiusculis, dissepimento parallelis. *Stylus* nullus.
- 2. LUNARIA. *Silicula* integra, elliptica, compresso-plana: *Valvis* dissepimento æqualibus, parallelis, planis. *Stylus* exsertus.
- 3. SUBULARIA. *Silicula* integra, ovata, *Valvis* ovatis, concavis, dissepimento contrariis. *Stylus* silicula brevior.
- 4. MYAGRUM. *Silicula* stylo conico terminata persistente: *Valvulis* concavis.

Silicula apice emarginata.

- 5. IBERIS. *Silicula* polysperma, emarginata. *Cor.* irregularis: petalis 2 exterioribus majoribus.
- 6. COCHLEARIA. *Silicula* emarginata, turgida, scabra: *Valvulis* gibbis, obtusis.
- 7. LEPIDIUM. *Silicula* emarginata cordata, polysperma: *Valvulis* carinatis, contrariis.
- 8. THLASPI. *Silicula* emarginata, obcordata, polysperma: *Valvulis* navicularibus, marginato-carinatis.

SILIUOSÆ.

*Calyx clausus: Foliolis longitudinaliter
conniventibus.*

249. RAPHANUS. *Cal.* clausus. *Siliqua* torosa & articulata, teres. *Glandula* melliferæ 2 interamina breviora & pistillum, totidem inter staminibus longiora & calycem.
244. ERYSIMUM. *Cal.* clausus. *Siliqua* columnaris exacte tetraëdra.
245. ARABIS. *Glandula* nectariferæ 4, singulæ inter calycis foliola squamæ instar reflexæ. *Stigma* simplex.
247. BRASSICA. *Cal.* erectus connivens. *Sem.* globosa. *Glandula* inter stamina breviora & pistillum interque longiora & calycem.
246. TURRITIS. *Cal.* erectus, connivens. *Cor.* erecta. *Siliqua* longissima, angulata.
241. DENTARIA. *Cal.* longitudinaliter connivens. *Stigma* emarginatum. *Siliqua* elastice diffiliens. *Valvulis* revolutis.

Calyx hians: Foliolis superne distantibus.

252. CRAMBE. *Siliqua* decidua, globosa, sicco-baccata. *Filamenta* 4, apice bifurca: altero antherifero.
251. ISATIS. *Siliqua* lanceolata, 1-locularis, 1-sperma, decidua, bivalvis. *Valvulis* navicularibus.
250. BUNIAS. *Siliqua* decidua, tetraëdra, angulata inæqualibus acuminatis muricata.
242. CARDAMINE. *Siliqua* elastice diffiliens, *Valvulis* revolutis. *Stigma* integrum. *Cal.* subhians.
248. SINAPIS. *Siliqua* dehiscens. *Cal.* patens. *Corolla* unguis rectæ. *Glandula* inter stamina breviora & pistillum interque longiora & calycem.
243. SISYMBRIUM. *Siliqua* dehiscens valvulis ætiusculis. *Calyx* patens. *Corolla* patula.

SILICULOSA.

233. MYAGRUM. 796.

08. MYAGRUM *sativum* filiculis obovatis pedunculatis, polyspermis.

Dan. *Gier*; *Vild - Hör*.

In agris.

09. MYAGRUM *paniculatum* filiculis lentiformibus orbiculatis punctato-rugosis. *Fl. Dan. t. 204.*

In agris haud infrequens.

234. SUBULARIA. 799.

10. † SUBULARIA *aquatica*. *Fl. Dan. t. 35.*

In lacustribus fabulosis.

235. DRABA. 800.

* *Caule nudo.*

11. DRABA *verna* scapis nudis: foliis lanceolatis subincisis.

Dan. *Gæslinge - Blomster*.

In apricis aridis ubique.

236. LEPIDIUM. 801.

12. LEPIDIUM *sativum* floribus tetradynamiis: foliis oblongis multifidis.

Pharmac. NASTURTII semen.

Ad hortum.

13. LEPIDIUM *ruderales* floribus diandris apetalis: foliis radicalibus dentato-pinnatis: ramiferis linearibus integerrimis. *Fl. Dan. t. 184.*

In ruderales.

237. THLASPI. 802.

- § 14. THLASPI *arvense* filiculis orbiculatis: foliis oblongis, dentatis, glabris. *P. Fl. Oec.* 142.

Dan. *Penge-Urt.*

Pharmac. THLASPEOS semina.

In agris.

- § 15. THLASPI *campestre* filiculis subrotundis: foliis sagittatis dentatis incanis.

In arvis apricis.

- § 16. THLASPI *Bursa pastoris* filiculis obcordatis: foliis radicalibus pinnatifidis.

Dan. *Hyrde-Taske.*

Pharmac. BURSÆ PASTORIS herba.

In cultis ubique.

238. COCHLEARIA. 803.

- § 17. COCHLEARIA *Danica* foliis hastatis angulatis.

Fl. Dan. t. 100.

Ad littora maris.

- § 18. † COCHLEARIA *officinalis* foliis radicalibus subrotundis; caulinis oblongis sublinuatis.

Fl. Dan. t. 135. *P. Fl. Oec.* 53.

Dan. *Cochlearie*; *Skiörbugs-Urt.*

Pharmac. COCHLEARIÆ herba.

- § 19. † COCHLEARIA *Coronopus* foliis pinnatifidis.

Fl. Dan. t. 202.

- § 20. † COCHLEARIA *Armoracia* foliis radicalibus lanceolatis crenatis: caulinis incisís.

Dan. *Vild Peber-Roed.*

Pharmac. ARMORACIÆ radix.

239. IBERIS. 804.

- § 21. IBERIS *nudicaulis* herbacea foliis sinuatis: caule nudo simplici.

Dan. *Steenkarse.*

In arenosis.

240. LUNARIA. 809.

522. † LUNARIA *annua* foliis oppositis.

SILIKUOSA.

241. DENTARIA. 811.

523. DENTARIA *bulbifera* foliis inferioribus pinnatis: summis simplicibus.

In nemore; raro. In præcipitiis promontorii Insulæ *Mona* vulgaris.

242. CARDAMINE. 812.

524. CARDAMINE *impatiens* foliis pinnatis incisis stipulatis: floribus apetalis.

Morif. Hist. III. p. 221. t. 4. f. 1.

In umbrosis ad ripam, in Sylva *Nassët*.

525. CARDAMINE *hirsuta* foliis pinnatis: floribus terandris. *Fl. Dan. t. 148.*

In nemore ad ripam hortum versus, & alibi passim.

526. CARDAMINE *pratensis* foliis pinnatis: foliolis radicalibus subrotundis: caulinis lanceolatis.

Dan. Enge - Karse.

In pascuis aquosis ubique.

527. CARDAMINE *amara* foliis pinnatis: axillis stoloniferis. *P. Fl. Oec. 44.*

Dan. Vand - Karse.

In aquosis umbrosis passim.

243. SISYMBRIUM. 813.

* *Siliquis declinatis brevibus.*

528. SISYMBRIUM *Nasturtium aquat.* siliquis declinatis: foliis pinnatis: foliolis subcordatis.

Dan. Vand - Karse.

Pharmac. NASTURTHI AQUATICI herba.

Minus frequens.

529. *SISYMBRIUM sylvestre* filiquis declinatis: foliis pinnatis: foliolis lanceolatis ferratis.

In humidis agrorum.

530. *SISYMBRIUM amphibium* filiquis declinatis oblongo-ovatis: foliis pinnatifidis ferratis.

Dan. *Vand - Peberroed*,

In palustribus.

* * Caule nudo.

531. † *SISYMBRIUM arenosum* caule subfolioso ramo: foliis lyratis rectangulo-dentatis hispidis: pilis ramosis.

* * * Foliis pinnatis.

532. *SISYMBRIUM Sophia* petalis calyce minoribus: foliis decomposito-pinnatis.

Pharmac. SOPHIÆ femina.

In maceris passim.

244. ERYSIMUM. 814.

533. † *ERYSIMUM officinale* filiquis spicæ appressis: foliis runcinatis.

Dan. *Vild Senip*.

Pharmac. ERYSIMI herba.

534. *ERYSIMUM Barbaræa* foliis lyratis: extimo subrotundo. Dan. *Vinter - Karse*.

Ad ripas raro.

535. *ERYSIMUM Alliaria* foliis cordatis.

Dan. *Hvidlögs - Urt*.

Pharmac. ALLIARIÆ herba.

In umbrosis frequens.

536. † *ERYSIMUM Cheiranthoides* foliis lanceolatis integerrimis.

245. A R A B I S. 818.

537. † ARABIS *Alpina* foliis amplexicaulibus dentatis.
Fl. Dan. t. 62.

538. ARABIS *Thaliana* foliis petiolatis lanceolatis integerrimis.
In aridis.

246. T U R R I T I S. 819.

539. TURRITIS *glabra* foliis radicalibus dentatis hispidis: caulinis integerrimis amplexicaulibus glabris.

Dan. *Vild Kaal.*

In siccis glareosis.

540. TURRITIS *hirsuta* foliis omnibus hispidis, caulinis amplexicaulibus.

In pascuis sylvaticis.

247. B R A S S I C A. 820.

541. BRASSICA *campestris* radice cauleque tenui: foliis caulinis uniformibus cordatis sessilibus.

Dan. *Vild Kaal.*

Inter Segetes.

248. S I N A P I S. 821.

542. SINAPIS *arvensis* filiquis multangulis torosoturgidis rostro longioribus. P. Fl. Oec. 136.

Dan. *Ager - Kaal.*

In agris.

543. SINAPIS *nigra* filiquis glabris apice tetragonis.

Dan. *Senip.*

Pharmac. SINAPIS femina.

In agris.

249. R A P H A N U S. 822.

544. RAPHANUS *Raphanistrum* filiquis teretibus articulatis lævibus unilocularibus. *P. Fl. Oec.* 121.

Dan. *Ager - Kaal*.

Inter hordeum.

250. B U N I A S. 823.

545. † BUNIAS *Cakile* filiculis ovatis lævibus ancipitibus.

Dan. *Strand - Karse*.

In sabuloso maritimo ad villas *Fionie Lundsgaard & Hindsgavl*.

251. I S A T I S. 824.

546. † ISATIS *tinctoria* foliis radicalibus crenatis: caulinis sagittatis: filiculis oblongis.

P. Fl. Oec. 86.

252. C R A M B E. 825.

547. CRAMBE *maritima* foliis cauleque glabris.

Dan. *Strand - Kaal*.

In littore maris.



Classis XVI.


MONADELPHIA.

DECANDRIA.

253. GERANIUM. *Monogyna. Stigmata 5. Caps.*
rostrata, 5 - cocca.

POLYANDRIA.

255. MALVA. *Calyx duplex : exterior 3 - phyllus.*
Caps. loculis 1 - spermis, verticillatis.
254. ALTHÆA. *Calyx duplex: exterior 9 - fidus.*
Caps. loculis 1 - spermis, verticillatis.



DECANDRIA.

253. GERANIUM. 832.

* *Staminibus quinque fertilibus.*

548. GERANIUM *cicutarium* pedunculis multifloris:
floribus pentandris: foliis pinnatis incisus obtusis.

Dan. *Storke-Nab.*

In nemore & ad aggeres.

* * *Staminibus decem fertilibus.*

α) Perennia.

549. GERANIUM *macrorrhizum* pedunculis bifloris:
calycibus inflatis: pistillo longissimo.

550. GERANIUM *sylvaticum* pedunculis bifloris: foliis
subpeltatis quinquelobis inciso-ferratis: caule ere-
cto: petalis emarginatis.

*Fl Dan. t. 124.*Dan. *Trane-Hals.*

In Nemore.

551. GERANIUM *palustre* pedunculis bifloris longissi-
mis declinatis, foliis quinquelobis incisus: petalis
integris.

552. † GERANIUM *pratense* pedunculis bifloris: foliis
subpeltatis multipartitis, pinnato-laciniatis rugo-
sis acutis: petalis integris.

β) Annua.

553. GERANIUM *Robertianum* pedunculis bifloris: ca-
lycibus pilosis decem angulatis. *P. Fl. Oec. 76.*

Dan. *Roberts-Urt.**Pharmac. GERANII ROBERTIANI herba.*

In ruderatis & ad muros.

554. † GERANIUM *lucidum* pedunculis bifloris: calycibus pyramidalis angulatis elevato - rugosis: foliis quinquelobis rotundatis. *Fl. Dan. t. 218.*

555. * GERANIUM *molle* pedunculis bifloris foliisque floralibus alternis: caule ramoso erectiusculo: calycibus muticis: arillis lævibus.

Vaill. Bot. Paris. t. 15. f. 3.

In nemore.

556. GERANIUM *dissectum* pedunculis bifloris: foliis quinquepartito - trifidis: petalis emarginatis longitudine calycis: arillis villosis.

Vaill. Bot. Paris. t. 15. f. 2.

Ad sepimenta.

557. * GERANIUM *columbinum* pedunculis bifloris folio longioribus: foliis quinquepartito - multifidis: laciniis acutis: arillis glabris: calycibus aristatis.

Vaill. Bot. Paris. t. 15. f. 4.

In arvis.

558. GERANIUM *retundifolium* pedunculis bifloris: petalis integris obtusissimis longitudine calycis: caule prostrato: foliis reniformibus incis.

Vaill. Bot. Paris. t. 15. f. 1.

In cultis.

* * * Pedunculis *unifloris*.

559. GERANIUM *sanguineum* pedunculis unifloris: foliis quinquepartitis trifidis orbiculatis.

In dumetis ficcis collis *Brankebakke* frequens.

POLYANDRIA.

254. ALTHÆA. 839.

560. † ALTHÆA *officinalis* foliis simplicibus tomentosis.
Dan. Althee.

255. M A L V A. 841.

561. *MALVA rotundifolia* caule prostrato: foliis cordato-orbiculatis, obsolete quinquelobis: pedunculis fructiferis declinatis.

Dan. *Kat - Ost.*

In ruderatis & ad pagos.

562. *MALVA sylvestris* caule erecto herbaceo: foliis septemlobatis acutis: pedunculis petiolisque pilosis.

Pharmac. MALVÆ VULGARIS radix, herba, semina.
Ad vias.

563. † *MALVA Alcea* caule erecto: foliis multipartitis scabriusculis.

Dan. *Cimons - Urt.*



Classis XVII.

DIADELPHIA.

HEXANDRIA.

256. FUMARIA. *Cal.* 2-phyllus. *Cor.* ringens, basi gibbosa, nectarifera. *Filamenta* 2 membranacea, singula antheris 3.

OCTANDRIA.

257. POLYGALA. *Cal.* 5-phyllus: foliolis duobus alariorum, coloratis. *Cor.* vexillum cylindricum. *Stamina* connexa. *Legumen* obcordatum, biloculare.

DECANDRIA.

Stamina omnia connexa.

258. SPARTIUM. *Stigma* longitudinale, supra villosum. *Filamenta* carinae adhærentia. *Cal.* deorsum productus.
259. GENISTA. *Pistillum* deprimens carinam. *Stigma* involutum. *Cal.* 2-labiatus: $\frac{2}{7}$. *Vexillum* oblongum, a pistillo staminibusque deorsum reflexum.
262. ANTHYLLIS. *Cal.* ventricosus, turgidus, filicquam includens. *Legumen* subrotundum, tectum.
261. ONONIS. *Cal.* 5-partitus: laciniis linearibus. *Vexillum* striatum. *Legumen* turgidum, rhombeum, fetile.

Stigma pubescens, (nec priorum notæ.)

264. OROBUS. *Stylus* linearis, teretiusculus, supra villosus. *Cal.* basi obtusus: *Lacinis* superioribus profundioribus, brevioribus.
263. PISUM. *Stylus* triangulus supra carinatus villosusque. *Calycis* lacinie superiores 2 breviores.
265. LATHYRUS. *Stylus* planus, supra villosus, superne lator. *Calycis* lacinie superiores 2 breviores.
266. VICIA. *Stylus* sub stigmate barbatus.

Legumen subbiloculare (nec priorum.)

270. ASTRAGALUS. *Legumen* biloculare, rotundatum, gibbum.

Legumina submonosperma (nec priorum.)

271. TRIFOLIUM. *Legumen* vix calyce longius, 1 - seu 2 - spermum, non dehiscens, deciduum. *Flores* capitati.

Legumen subarticulatum.

269. HEDYSARUM. *Legumen* articulis subrotundis, compressis, 1 - spermis. *Cor.* carina obtusissima.
260. CORONILLA. *Legumen* isthmis interceptum, rectum. *Cal.* 2 - labiatus: $\frac{2}{3}$. dentibus superioribus connatis. *Vexillum* vix alis longius.
268. ORNITHOPUS. *Legumen* articulatum, teres, arcuatum.
273. MEDICAGO. *Legumen* spirale, membranaceo-compressum. *Carina* corollæ a vexillo deflectens.

Legumen uniloculare, polyspermum (nec priorum.)

267. ERVUM. *Calyx* 5 - partitus, subæqualis longitudine fere corollæ.
272. LOTUS. *Legumen* cylindricum, strictum, factum seminibus cylindricis. Alæ sursum longitudinaliter conniventes. *Cal.* tubulosus.

HEXANDRIA.

HEXANDRIA.

256. FUMARIA. 849.

64. FUMARIA *bulbosa* caule simplici: bracteis longitudine florum.

α) Fumaria bulbosa radice cava, major.

β) Fumaria bulbosa radice non cava, minor.

Morif. Hist. p. 261, t. 12. f. 8.

Dan. Hjul-Roed; Röd Hane-Kumm.

Pharmac. ARISTOLOCHIAE FABACEAE radix.

α) Frequens in nemorosis ad lacum Werelöse.

β) Frequens in Vivario Friderichsdalensi.

65. FUMARIA *officinalis* pericarpis monospermis racemosis: caule diffuso.

Dan. Jordrög.

Pharmac. FUMARIAE herba, semina.

In arvis & saxis frequens.

OCTANDRIA.

257. POLYGALA. 851.

66. POLYGALA *vulgaris* floribus cristatis racemosis: caulibus herbaceis simplicibus procumbentibus: foliis lineari-lanceolatis.

Vaill. Bot. Paris. t. 32. f. 2.

Dan. Kaars-Blomster.

In ficcis vulgaris.

β) POLYGALA violacea.

γ) POLYGALA carnea.

δ) POLYGALA alba.

ε) POLYGALA major. Vaill. Bot. Paris. t. 32. f. 2.

DECANDRIA.

258. SPARTIUM. 88.

567. SPARTIUM *scoparium* foliis ternatis solitariisque
ramis inermibus angulatis.

In arenosis fylvarum.

259. GENISTA. 859.

* *Inermes.*

568. † GENISTA *tinctoria* foliis lanceolatis glabris:
ramis striatis teretibus erectis.

Pharmac. GENISTÆ herba, flores, femina.

569. † GENISTA *pilosa* foliis lanceolatis obtusis: caulibus
tuberculato decumbente.

In Ducatibus obvia.

* * *Spinosa.*

570. † GENISTA *Germanica* spinis compositis: ramis
floriferis inermibus: foliis lanceolatis.

In ericetis *Holfatie* passim.

260. CORONILLA. 883.

571. † CORONILLA *varia* herbacea, leguminibus erectis
teretibus torosis numerosis: foliolis plurimis
glabris.

261. ONONIS. 863.

572. ONONIS *spinosa* floribus racemosis geminis, foliis
ternatis: superioribus solitariis: ramis inermibus
subvillosis.

Dan. Krage-Torn.

Pharmac. ONONIDIS radix, herba.

In campestri frequens.

262. ANTHYLLIS. 864.

3. ANTHYLLIS *vulneraria* herbacea: foliis quaternis - pinnatis: floribus lateralibus.

Dan. *Katte - Klever.*

In sterilibus.

263. PISUM. 870.

4. PISUM *maritimum* petiolis supra planiusculis: caule angulato: stipulis sagittatis: pedunculis multifloris.

Moriz. Hist. t. I. f. 5.

In maritimis.

264. OROBUS. 871.

5. OROBUS *vernus* foliis pinnatis ovatis: stipulis semisagittatis integerrimis: caule simplici.

Dan. *Muus - Erter.*

Pharmac. OROBI farina.

In montosis nemorum.

6. * OROBUS *tuberosus* foliis pinnatis lanceolatis: stipulis semisagittatis integerrimis: caule simplici.

In Sylvis.

7. * OROBUS *niger* caule ramoso: foliis sexjugis ovato - oblongis.

In montosis sylvaticis.

265. LATHYRUS. 872.

* Pedunculis *unifloris.*

78. † LATHYRUS *Aphaca* pedunculis unifloris: circumsistis aphyllis: stipulis sagittato - cordatis.

Dan. *Vikker.*

* * *Pedunculis multifloris.*

579. † *LATHYRUS tuberosus* pedunculis multifloris
cirrhis diphyllis: foliolis ovalibus: internodiis
nudis. *P. Fl. Oec.* 89.

Dan. *Jord - Rödder; Heede - Nödder.*

580. *LATHYRUS pratensis* pedunculis multifloris: cirrhis
diphyllis simplicissimis: foliolis lanceolatis.

Dan. *Vilde Erter.*

In dumetis.

581. *LATHYRUS sylvestris* pedunculis multifloris: cirrhis
diphyllis: foliolis ensiformibus: internodiis
membranaceis.

Extra hortum ac in declivi *Brankebakke.*

582. *LATHYRUS latifolius* pedunculis multifloris: cirrhis
diphyllis: foliolis lanceolatis: internodiis membranaceis.

583. * *LATHYRUS palustris* pedunculis multifloris
cirrhis polyphyllis: stipulis lanceolatis.

In pascuis humidis.

266. *V I C I A.* 873.* *Pedunculis elongatis.*

584. *VICIA dumetorum* pedunculis multifloris: foliolis
reflexis ovatis mucronatis: stipulis subdentatis.

Dan. *Vikker.*

585. † *VICIA sylvatica* pedunculis multifloris: foliolis
ovalibus: stipitulis denticulatis.

586. *VICIA Cracca* pedunculis multifloris: floribus imbricatis:
foliolis lanceolatis pubescentibus: stipulis integris.

Dan. *Muus - Erter.*

Inter segetes, ac in nemore versus amnem.

* * Floribus axillaribus subsessilibus.

7. VICIA *sativa* leguminibus sessilibus subbinatis erectis: foliolis retusis: stipulis notatis.

[inter segetes.

8. VICIA *Lathyroides* leguminibus sessilibus solitariis, erectis, glabris: foliolis senis: inferioribus obcordatis. *Fl. Dan. t. 58.*

In aridis montosis, ac in agris passim.

9. VICIA *sepium* leguminibus pedicellatis subquaternis erectis: foliolis ovatis integerrimis: exterioribus decescentibus.

In umbrosis.

267. ERVUM. 874.

90. ERVUM *tetraspermum* pedunculis subbifloris: feminibus globosis quaternis. *Fl. Dan. t. 95.*

In oleraceis frequens.

91. ERVUM *hirsutum* pedunculis multifloris: feminibus globosis binis.

In arvis, ac in horto frequens.

268. ORNITHOPUS. 884.

92. † ORNITHOPUS *perpusillus* foliis pinnatis: leguminibus subarcuatis.

269. HEDYSARUM. 887.

93. † HEDYSARUM *Onobrychis* foliis pinnatis: leguminibus monospermis aculeatis: corollarum alis calyce brevioribus: caule elongato.

* * Pedunculis multifloris in capitulum.

606. *LOTUS corniculatus* capitulis depressis: caulibus decumbentibus: leguminibus cylindricis patentibus. *P. Fl. Oec.* 95.

Dan. *Kierlinge - Tender; Krage - Kloer.*

In pratibus ac collibus.

273. M E D I C A G O. 899.

607. *MEDICAGO falcata* pedunculis racemosis: leguminibus lunatis: caule prostrato.

Fl. Dan. t. 233. *P. Fl. Oec.* 97.

Dan. *Gaul - Klever.*

In Icone legumina desunt, caracter plantæ specificus.

In agris ad vias.

608. † *MEDICAGO lupulina* spicis ovalibus: leguminibus reniformibus monospermis: caule procumbente.

In pratibus & ad vias.

609. † *MEDICAGO polymorpha* minima leguminibus cochleatis aculeis uncinatis alternis externe donatis: stipulis integris. *Fl. Dan. t.* 211.



Classis XVIII.

POLYADELPHIA.

POLYANDRIA.

274. **HYPERICUM.** *Cal.* 5 - partitus. *Cor.* 5 - petala.
Filamenta multa in 5 phalangibus. *Styli* 1, 3, 1. 5.
Caps. locularis.



POLYANDRIA.

274. HYPERICUM. 902.

610. HYPERICUM *quadrangulum* floribus trigynis: caule quadrato herbaceo.

Dan. *Piricum*.

In pratis.

611. HYPERICUM *perforatum* floribus trigynis: caule ancipiti, foliis obtusis pellucido - punctatis,

P. Fl. Oec. 85.

Dan. *Hypericon*; *Jord-Humle*; *Blod-Blomster*.

Pharmac. HYPERICI herba, flores, semina.

612. HYPERICUM *humifusum* floribus trigynis axillaribus solitariis: caulibus ancipitibus prostratis filiformibus: foliis glabris. *Fl. Dan. t.* 141.

In arvis haud infrequens.

613. HYPERICUM *hirsutum* floribus trigynis: calycibus ferrato-glandulosis: caule tereti erecto: foliis ovatis subpubescentibus.

Morif. Hist. III. t. 6. f. 11.

In montosis minus frequens.

614. * HYPERICUM *montanum* floribus trigynis: calycibus ferraturis glandulosis: caule tereti erecto glabro: foliis ovatis. *Fl. Dan. t.* 173.

Passim: ad molendina haud infrequens.

615. † HYPERICUM *pulchrum* floribus trigynis: calycibus ferrato-glandulosis: caule tereti: foliis perfoliatis, cordatis, glabris. *Fl. Dan. t.* 75.

In insula *Thorlinge* legi.



Classis XIX.

SYNGENESIA.

POLYGAMIA ÆQUALIS.

Semiflosculosi: Corollis ligulatis omnibus.

- 287. CICHORIUM. *Receptaculum* paleaceum. *Pappus* 5-dentatus. *Calyx* calyculatus.
- 285. HYPOCHÆRIS. *Recept.* paleaceum. *Pappus* subplumosus. *Calyx* subimbricatus.
- 275. TRAGOPOGON. *Recept.* nudum. *Pappus* plumosus. *Cal.* simplex polyphyllus.
- 277. PICRIS. *Recept.* nudum. *Pappus* plumosus. *Cal.* calyculatus. *Semina* transversim sulcata.
- 281. LEONTODON. *Recept.* nudum. *Pappus* plumosus. *Cal.* imbricatus. *Squamis* laxis.
- 278. SONCHUS. *Recept.* nudum. *Pappus* plumosus. *Cal.* imbricatus, basi gibbus.
- 276. SCORZONERA. *Recept.* nudum. *Pappus* plumosus. *Cal.* imbricatus. *Squamis* margine membranaceis.
- 283. CREPIS. *Recept.* nudum. *Pappus* pilosus, stipitatus. *Cal.* calyculatus. *Squamis* deciduis.
- 280. PRENANTHES. *Recept.* nudum. *Pappus* pilosus, simplex, sessilis. *Cal.* calyculatus sub-5-florus.
- 279. LACTUCA. *Recept.* nudum. *Pappus* pilosus, simplex, stipitatus. *Cal.* imbricatus, cylindricus.
- 282. HIERACIUM. *Recept.* nudum. *Pappus* pilosus, simplex, sessilis. *Cal.* imbricatus, ovatus.
- 286. LAPSANA. *Recept.* nudum. *Pappus* nonnullus nullus. *Cal.* calyculatus. *Squamis* singulis interioribus canaliculatis.

DECANDRIA.

258. SPARTIUM. 88.

567. SPARTIUM *scoparium* foliis ternatis solitariisque:
ramis inermibus angulatis.

In arenosis fylvarum.

259. GENISTA. 859.

* *Inermes.*

568. † GENISTA *tinctoria* foliis lanceolatis glabris:
ramis striatis teretibus erectis.

Pharmac. GENISTÆ herba, flores, semina.

569. † GENISTA *pilosa* foliis lanceolatis obtusis : caule
tuberculato decumbente.

In Ducatibus obvia.

* * *Spinosa.*

570. † GENISTA *Germanica* spinis compositis : ramis
floriferis inermibus : foliis lanceolatis.

In ericetis *Holsatiæ* passim.

260. CORONILLA. 883.

571. † CORONILLA *varia* herbacea, leguminibus ere-
ctis teretibus torosis numerosis : foliolis plurimis
glabris.

261. ONONIS. 863.

572. ONONIS *spinosa* floribus racemosis geminis, fo-
liis ternatis : superioribus solitariis : ramis inermi-
bus subvillosis.

Dan. *Krage-Torn.*

Pharmac. ONONIDIS radix, herba.

In campestri frequens.

262. ANTHYLLIS. 864.

73. ANTHYLLIS *vulneraria* herbacea: foliis quaternis-pinnatis: floribus lateralibus.

Dan. *Katte - Klever.*

In sterilibus.

263. PISUM. 870.

74. PISUM *maritimum* petiolis supra planiusculis: caule angulato: stipulis sagittatis: pedunculis multifloris.

Morif. Hist. t. I. f. 5.

In maritimis.

264. OROBUS. 871.

75. OROBUS *vernus* foliis pinnatis ovatis: stipulis semisagittatis integerrimis: caule simplici.

Dan. *Muus - Erter.*

Pharmac. OROBI farina.

In montosis nemorum.

76. * OROBUS *tuberosus* foliis pinnatis lanceolatis: stipulis semisagittatis integerrimis: caule simplici.

In Sylvis.

77. * OROBUS *niger* caule ramoso: foliis sexjugis ovato-oblongis.

In montosis sylvaticis.

265. LATHYRUS. 872.

* Pedunculis *unifloris*.

78. † LATHYRUS *Aphaca* pedunculis unifloris: circumsistis aphyllis: stipulis sagittato-cordatis.

Dan. *Vikker.*

* * Pedunculis multifloris.

579. † LATHYRUS *tuberosus* pedunculis multifloris : cirrhis diphyllis : foliolis ovalibus : internodiis nudis. *P. Fl. Oec.* 89.

Dan. *Jord - Rödder ; Heede - Nödder.*

580. LATHYRUS *pratensis* pedunculis multifloris : cirrhis diphyllis simplicissimis : foliolis lanceolatis.

Dan. *Vilde Erter.*

In dumetis.

581. LATHYRUS *sylvestris* pedunculis multifloris : cirrhis diphyllis : foliolis ensiformibus : internodiis membranaceis.

Extra hortum ac in declivi *Brankebakke.*

582. LATHYRUS *latifolius* pedunculis multifloris : cirrhis diphyllis : foliolis lanceolatis : internodiis membranaceis.

583. * LATHYRUS *palustris* pedunculis multifloris : cirrhis polyphyllis : stipulis lanceolatis.

In pascuis humidis.

266. V I C I A. 873.

* Pedunculis elongatis.

584. VICIA *dumetorum* pedunculis multifloris : foliolis reflexis ovatis mucronatis : stipulis subdentatis.

Dan. *Vikker.*

585. † VICIA *sylvatica* pedunculis multifloris : foliolis ovalibus : stipitulis denticulatis.

586. VICIA *Cracca* pedunculis multifloris : floribus imbricatis : foliolis lanceolatis pubescentibus : stipulis integris.

Dan. *Muus - Erter.*

Inter segetes, ac in nemore versus amnem.

* * Floribus axillaribus subsessilibus.

7. VICIA *fativa* leguminibus sessilibus subbinatis erectis: foliolis retusis: stipulis notatis.

Inter segetes.

38. VICIA *Lathyroides* leguminibus sessilibus solitariis, erectis, glabris: foliolis fenis: inferioribus obcordatis. *Fl. Dan. t. 58.*

In aridis montosis, ac in agris passim.

39. VICIA *sepium* leguminibus pedicellatis subquaternis erectis: foliolis ovatis integerrimis: exterioribus decescentibus.

In umbrosis.

267. ERVUM. 874.

90. ERVUM *tetraspermum* pedunculis subbifloris: seminibus globosis quaternis. *Fl. Dan. t. 95.*

In oleraceis frequens.

91. ERVUM *hirsutum* pedunculis multifloris: seminibus globosis binis.

In arvis, ac in horto frequens.

268. ORNITHOPUS. 884.

92. † ORNITHOPUS *perpusillus* foliis pinnatis: leguminibus subarcuatis.

269. HEDYSARUM. 887.

93. † HEDYSARUM *Onobrychis* foliis pinnatis: leguminibus monospermis aculeatis: corollarum alis calyce brevioribus: caule elongato.

270. ASTRAGALUS. 892.

- §94. ASTRAGALUS *glycyphyllus* caulescens prostratus leguminibus subtriquetris arcuatis: foliis ovalibus pedunculo longioribus. *Morif. Hist. III. t. 9. f. 2.*
Dan. *Vild Lakrits.*

In sylvatico *Deyligheden* frequens, & alibi.

- §95. † ASTRAGALUS *arenarius* subcaulescens, procumbens, floribus subracemosis erectis: foliis tomentosis.

Dan. *Bierg - Lakrits.*

271. TRIFOLIUM. 896.

* *Meliloti leguminibus nudis polyspermis.*

- §96. TRIFOLIUM *M. officinalis* leguminibus racemosis nudis, dispermis, rugosis, acutis: caule erecto.
Dan. *Amur; Steen - Klever.*

Pharmac. MELILOTI herba, flores, semina.

In *Fionia* vulgaris; extra *Haderslebiam* corollis omnibus nitide albis legi.

In campestribus, imprimis inter *Molendina*.

** *Lotoidea leguminibus testis polyspermis.*

- §97. † TRIFOLIUM *hybridum* capitulis umbellaribus: leguminibus tetraspermis: caule adscendente.
Caulis teres, subpilosus, non sulcatus.

In via ultra cauponam *Kiøge - Kroe* vulgo frequentissimam legi.

- §98. TRIFOLIUM *repens* capitulis umbellaribus: leguminibus tetraspermis: caule repente.

Dan. *Huid - Klever.*

Pharmac. TRIFOLII ALBI flores.

In pascuis.

* * * *Lagopoda calycibus villosis.*

599. *TRIFOLIUM pratense* spicis subvillosis cinctis: stipulis oppositis membranaceis: corollis monopetalis. *P. Fl. Oec.* 146.

Dan. *Sue - Kop; Röd - Klever; Sukker - Bröd.*

In pratis ubique.

600. *TRIFOLIUM arvense* spicis villosis ovalibus: dentibus calycinis setaceis æqualibus.

Dan. *Hare - Klever.*

In arvis.

* * * * *Vesicaria calycibus inflatis ventricosis.*

601. † *TRIFOLIUM spumosum* spicis ovatis: calycibus inflatis glabris quinque dentatis: involucris universalibus pentaphyllis.

602. *TRIFOLIUM fragiferum* capitulis subrotundis: calycibus inflatis bidentatis reflexis: caubus repentibus. *Vaill. Bot. Paris. t. 22. f. 2.*

* * * * * *Lupulina vexillis corollæ inflexis.*

603. † *TRIFOLIUM montanum* spicis subimbricatis subtribus: vexillis subulatis emarcescentibus: calycibus nudis: caule erecto.

604. † *TRIFOLIUM agrarium* spicis ovalibus imbricatis: vexillis deflexis persistentibus: calycibus nudis: caule erecto.

Dan. *Humble - Klever.*

272. *LOTUS*. — 897.

* *Leguminibus rarioribus, nec capitulum constituentibus.*

605. † *LOTUS maritimus* leguminibus solitariis membranaceo-quadrangulatis: foliis glabris bracteis lanceolatis.

Dan. *Steen - Klever.*

* * Pedunculis *multifloris in capitulum.*

606. *LOTUS corniculatus* capitulis depresso: caulis decumbentibus: leguminibus cylindricis patentibus. *P. Fl. Oec. 95.*

Dan. *Kierlinge - Tander; Krage - Kloer.*
In pratis ac collibus.

273. M E D I C A G O. 899.

607. *MEDICAGO falcata* pedunculis racemosis: leguminibus lunatis: caule prostrato.

Fl. Dan. t. 233. P. Fl. Oec. 97.

Dan. *Gaul - Klever.*

In Icone legumina defunt, caracter plantæ specificus.
In agris ad vias.

608. † *MEDICAGO lupulina* spicis ovalibus: leguminibus reniformibus monospermis: caule procumbente.

In pratis & ad vias.

609. † *MEDICAGO polymorpha* minima leguminibus cochleatis aculeis uncinatis alternis externe donatis: stipulis integris. *Fl. Dan. t. 211.*



Classis XVIII.

POLYADELPHIA.

POLYANDRIA.

174. HYPERICUM. *Cal.* 5 - partitus. *Cor.* 5 - petala.
Filamenta multa in 5 phalangibus. *Styli* 1, 3, 1 5.
Caps. locularis.



Stigma pubescens, (nec priorum notæ.)

264. OROBUS. *Stylus* linearis, teretiusculus, supra villosus. *Cal.* basi obtusus: *Lacinis* superioribus profundioribus, brevioribus.
263. PISUM. *Stylus* triangulus supra carinatus villosusque. *Calycis* lacinia superiores 2 breviores.
265. LATHYRUS. *Stylus* planus, supra villosus, superne lator. *Calycis* lacinia superiores 2 breviores.
266. VICIA. *Stylus* sub stigmate barbatus.

Legumen subbiloculare (nec priorum.)

270. ASTRAGALUS. *Legumen* biloculare, rotundatum, gibbum.

Legumina submonosperma (nec priorum.)

271. TRIFOLIUM. *Legumen* vix calyce longius, 1 - seu 2 - spermum, non dehiscens, deciduum. *Flores* capitati.

Legumen subarticulatum.

269. HEDYSARUM. *Legumen* articulis subrotundis, compressis, 1 - spermis. *Cor.* carina obtusissima.
260. CORONILLA. *Legumen* isthmis interceptum, rectum. *Cal.* 2-labiatus: $\frac{2}{3}$. dentibus superioribus connatis. *Vexillum* vix alis longius.
268. ORNITHOPUS. *Legumen* articulatum, teres, arcuatum.
273. MEDICAGO. *Legumen* spirale, membranaceo-compressum. *Carina* corollæ a vexillo deflectens.

Legumen uniloculare, polyspermum (nec priorum.)

267. ERVUM. *Calyx* 5-partitus, subæqualis longitudine fere corollæ.
272. LOTUS. *Legumen* cylindricum, strictum, factum seminibus cylindricis. Alæ sursum longitudinaliter conniventes. *Cal.* tubulosus.

HEXANDRIA.

HEXANDRIA.

256. FUMARIA. 849.

64. FUMARIA *bulbosa* caule simplici: bracteis longitudine florum.

- a) Fumaria bulbosa radice cava, major.
- β) Fumaria bulbosa radice non cava, minor.

Morif. Hist. p. 261. t. 12. f. 8.

Dan. *Huul-Roed; Röd Hane-Kamm.*

Pharmac. ARISTOLOCHIE FABACEÆ radix.

- a) Frequens in nemorosis ad lacum *Warelöse.*
- β) Frequens in Vivario *Friderichsdalensi.*

65. FUMARIA *officinalis* pericarpis monospermis racemosis: caule diffuso.

Dan. *Jordrög.*

Pharmac. FUMARIÆ herba, semina.

In arvis & fatis frequens.

OCTANDRIA.

257. POLYGALA. 851.

66. POLYGALA *vulgaris* floribus cristatis racemosis: caulibus herbaceis simplicibus procumbentibus: foliis lineari-lanceolatis.

Vaill. Bot. Paris. t. 32. f. 2.

Dan. *Kaars-Blomster.*

In ficcis vulgaris.

β) POLYGALA *violacea.*

γ) POLYGALA *carnea.*

δ) POLYGALA *alba.*

ε) POLYGALA *major.* *Vaill. Bot. Paris. t. 32. f. 2.*

529. *SISYMBRIUM sylvestre* filiquis declinatis: foliis pinnatis: foliolis lanceolatis ferratis.
In humidis agrorum.

530. *SISYMBRIUM amphibium* filiquis declinatis oblongo-ovatis: foliis pinnatifidis ferratis.
Dan. *Vand - Peberroed*.
In palustribus.

* * Caule nudo.

531. † *SISYMBRIUM arenosum* caule subfolioso ramofo: foliis lyratis rectangulo-dentatis hispidis: pilis ramolis.

* * * Foliis pinnatis.

532. *SISYMBRIUM Sophia* petalis calyce minoribus: foliis decomposito-pinnatis.
Pharmac. SOPHIÆ femina.
In maceris passim.

244. E R Y S I M U M. 814.

533. † *ERYSIMUM officinale* filiquis spicæ appressis: foliis runcinatis.

Dan. *Vild Senip*.

Pharmac. ERYSIMI herba.

534. *ERYSIMUM Barbaræa* foliis lyratis: extimo subrotundo. Dan. *Vinter - Karse*.

Ad ripas raro.

535. *ERYSIMUM Alliaria* foliis cordatis.

Dan. *Hvidlögs - Urt*.

Pharmac. ALLIARIÆ herba.

In umbrosis frequens.

536. † *ERYSIMUM Cheiranthoides* foliis lanceolatis integerrimis.

245. A R A B I S. 818.

537. † ARABIS *Alpina* foliis amplexicaulibus dentatis.
Fl. Dan. t. 62.

538. ARABIS *Thaliana* foliis petiolatis lanceolatis integerrimis.
In aridis.

246. T U R R I T I S. 819.

539. TURRITIS *glabra* foliis radicalibus dentatis hispidis: caulinis integerrimis amplexicaulibus glabris.

Dan. *Vild Kaal.*

In siccis glareosis.

540. TURRITIS *hirsuta* foliis omnibus hispidis, caulinis amplexicaulibus.

In pascuis sylvaticis.

247. B R A S S I C A. 820.

541. BRASSICA *campestris* radice cauleque tenui: foliis caulinis uniformibus cordatis sessilibus.

Dan. *Vild Kaal.*

Inter Segetes.

248. S I N A P I S. 821.

542. SINAPIS *arvensis* filiquis multangulis torosoturgidis rostro longioribus. P. Fl. Oec. 136.

Dan. *Ager - Kaal.*

In agris.

543. SINAPIS *nigra* filiquis glabris apice tetragonis.

Dan. *Senip.*

Pharmac. SINAPIS femina.

In agris.

249. R A P H A N U S. 822.

544. RAPHANUS *Raphanistrum* filiquis teretibus articulatis lævibus unilocularibus. *P. Fl. Oec.* 121.

Dan. *Ager - Kaal*.

Inter hordeum.

250. B U N I A S. 823.

545. † BUNIAS *Cakile* filiculis ovatis lævibus ancipitibus.

Dan. *Strand - Karse*.

In sabuloso maritimo ad villas *Fionie Lundsgaard* & *Hindsgavl*.

251. I S A T I S. 824.

546. † ISATIS *tinctoria* foliis radicalibus crenatis: caulinis sagittatis: filiculis oblongis.


P. Fl. Oec. 86.

252. C R A M B E. 825.

547. CRAMBE *maritima* foliis cauleque glabris.

Dan. *Strand - Kaal*.

In littore maris.




*Classis XVI.***MONADELPHIA.****DECANDRIA.**

- 253.** GERANIUM. *Monogyna. Stigmata 5. Caps.*
rostrata, 5-cocca.

POLYANDRIA.

- 255.** MALVA. *Calyx duplex: exterior 3-phyllus.*
Caps. loculis 1-spermis, verticillatis.
- 254.** ALTHÆA. *Calyx duplex: exterior 9-fidus.*
Caps. loculis 1-spermis, verticillatis.



DECANDRIA.

253. GERANIUM. 832.

* *Staminibus quinque fertilibus.*

548. GERANIUM *cicutarium* pedunculis multifloris:
floribus pentandris: foliis pinnatis incisis obtusis.

Dan. *Storke-Nab.*

In nemore & ad aggeres.

* * *Staminibus decem fertilibus.*

α) Perennia.

549. GERANIUM *macrorrhizum* pedunculis bifloris:
calycibus inflatis: pistillo longissimo.

550. GERANIUM *sylvaticum* pedunculis bifloris: foliis
subpeltatis quinquelobis inciso-ferratis: caule ere-
cto: petalis emarginatis.

*Fl. Dan. t. 124.*Dan. *Trane-Hals.*

In Nemore.

551. GERANIUM *palustre* pedunculis bifloris longissi-
mis declinatis, foliis quinquelobis incisis: petalis
integris.

552. † GERANIUM *pratense* pedunculis bifloris: foliis
subpeltatis multipartitis, pinnato-laciniatis rugo-
sis acutis: petalis integris.

β) Annua.

553. GERANIUM *Robertianum* pedunculis bifloris: ca-
lycibus pilosis decem angulatis. *P. Fl. Oec. 76.*

Dan. *Roberts-Urt.**Pharmac.* GERANII ROBERTIANI herba.

In ruderatis & ad muros.

554. † GERANIUM *lucidum* pedunculis bifloris: calycibus pyramidatis angulatis elevato-rugosis: foliis quinquelobis rotundatis. *Fl. Dan. t. 218.*

555. * GERANIUM *molle* pedunculis bifloris folisque floralibus alternis: caule ramoso erectiusculo: calycibus muticis: arillis lævibus.

Vaill. Bot. Paris. t. 15. f. 3.

In nemore.

556. GERANIUM *dissectum* pedunculis bifloris: foliis quinquepartito-trifidis: petalis emarginatis longitudine calycis: arillis villosis.

Vaill. Bot. Paris. t. 15. f. 2.

Ad sepimenta.

557. * GERANIUM *columbinum* pedunculis bifloris folio longioribus: foliis quinquepartito-multifidis: laciniis acutis: arillis glabris: calycibus aristatis.

Vaill. Bot. Paris. t. 15. f. 4.

In arvis.

558. GERANIUM *retundifolium* pedunculis bifloris: petalis integris obtusissimis longitudine calycis: caule prostrato: foliis reniformibus incis.

Vaill. Bot. Paris. t. 15. f. 1.

In cultis.

* * * Pedunculis *unifloris*.

559. GERANIUM *sanguineum* pedunculis unifloris: foliis quinquepartitis trifidis orbiculatis.

In dumetis siccis collis *Brankebakke* frequens.

POLYANDRIA.

254. A L T H Æ A. 839.

560. † ALTHÆA *officinalis* foliis simplicibus tomentosa.
Dan. *Althee*.

255. M A L V A. 841.

561. *MALVA rotundifolia* caule prostrato: foliis cordato-orbiculatis, obsolete quinquelobis: pedunculis fructiferis declinatis.

Dan. *Kat - Ost.*

In ruderatis & ad pagos.

562. *MALVA sylvestris* caule erecto herbaceo: foliis septemlobatis acutis: pedunculis petiolisque pilosis.

Pharmac. MALVÆ VULGARIS radix, herba, semina.
Ad vias.

563. † *MALVA Alcea* caule erecto: foliis multipartitis scabriusculis.

Dan. *Cimons - Urt.*



Classis XVII.

DIADELPHIA.

HEXANDRIA.

256. FUMARIA. *Cal.* 2-phyllus. *Cor.* ringens, basi gibbosa, nectarifera. *Filamenta* 2 membranacea, singula antheris 3.

OCTANDRIA.

257. POLYGALA. *Cal.* 5-phyllus: foliolis duobus alæformibus, coloratis. *Cor.* vexillum cylindricum. *Stamina* connexa. *Legumen* obcordatum, biloculare.

DECANDRIA.

Stamina omnia connexa.

258. SPARTIUM. *Stigma* longitudinale, supra villosum. *Filamenta* carinæ adhærentia. *Cal.* deorsum productus.
259. GENISTA. *Pistillum* deprimens carinam. *Stigma* involutum. *Cal.* 2-labiatus: $\frac{2}{3}$. *Vexillum* oblongum, a pistillo staminibusque deorsum reflexum.
262. ANTHYLLIS. *Cal.* ventricosus, turgidus, filiquam includens. *Legumen* subrotundum, tectum.
261. ONONIS. *Cal.* 5-partitus: laciniis linearibus. *Vexillum* striatum. *Legumen* turgidum, rhombeum, sessile.

Stigma pubescens, (nec priorum notæ.)

264. OROBUS. *Stylus* linearis, teretiusculus, supra villosus. *Cal.* basi obtusus: *Lacinis* superioribus profundioribus, brevioribus.
263. PISUM. *Stylus* triangulus supra carinatus villosusque. *Calycis* lacinix superiores 2 breviores.
265. LATHYRUS. *Stylus* planus, supra villosus, superne lator. *Calycis* lacinix superiores 2 breviores.
266. VICIA. *Stylus* sub stigmate barbatus.

Legumen subbiloculare (nec priorum.)

270. ASTRAGALUS. *Legumen* biloculare, rotundatum, gibbum.

Legumina submonosperma (nec priorum.)

271. TRIFOLIUM. *Legumen* vix calyce longius, 1 - seu 2 - spermum, non dehiscens, deciduum. *Flores* capitati.

Legumen subarticulatum.

269. HEDYSARUM. *Legumen* articulis subrotundis, compressis, 1 - spermis. *Cor.* carina obtusissima.
260. CORONILLA. *Legumen* isthmis interceptum, rectum. *Cal.* 2 - labiatus: $\frac{2}{3}$. dentibus superioribus connatis. *Vexillum* vix alis longius.
268. ORNITHOPUS. *Legumen* articulatum, teres, arcuatum.
273. MEDICAGO. *Legumen* spirale, membranaceo-compressum. *Carina* corollæ a vexillo deflectens.

Legumen uniloculare, polyspermum (nec priorum)

267. ERVUM. *Calyx* 5 - partitus, subæqualis longitudine fere corollæ.
272. LOTUS. *Legumen* cylindricum, strictum, factum feminibus cylindricis. Alæ sursum longitudinaliter conniventes. *Cal.* tubulosus.

HEXANDRIA

HEXANDRIA.

256. FUMARIA. 849.

64. FUMARIA *bulbosa* caule simplici: bracteis longitudine florum.

- a) Fumaria bulbosa radice cava, major.
- β) Fumaria bulbosa radice non cava, minor.

Morif. Hist. p. 261, t. 12. f. 8.

Dan. *Hvul-Roed; Röd Hane-Kamm.*

Pharmac. ARISTOLOCHIE FABACEE radix.

- a) Frequens in nemorosis ad lacum *Warelöse.*
- β) Frequens in Vivario *Friderichsdalensi.*

65. FUMARIA *officinalis* pericarpis monospermis racemosis: caule diffuso.

Dan. *Jordrög.*

Pharmac. FUMARIE herba, semina.

In arvis & fatis frequens.

OCTANDRIA.

257. POLYGALA. 851.

66. POLYGALA *vulgaris* floribus cristatis racemosis: caulibus herbaceis simplicibus procumbentibus: foliis lineari-lanceolatis.

Vaill. Bot. Paris. t. 32. f. 2.

Dan. *Kaars-Blomster.*

In ficcis vulgaris.

- β) POLYGALA *violacea.*
- γ) POLYGALA *carnea.*
- δ) POLYGALA *alba.*
- ε) POLYGALA *major.*

Vaill. Bot. Paris. t. 32. f. 2.

DECANDRIA.

258. SPARTIUM. 88.

567. SPARTIUM *scoparium* foliis ternatis solitariisque:
ramis inermibus angulatis.

In arenosis sylvarum.

259. GENISTA. 859.

* *Inermes.*

568. † GENISTA *tinctoria* foliis lanceolatis glabris:
ramis striatis teretibus erectis.

Pharmac. GENISTÆ herba, flores, semina.

569. † GENISTA *pilosa* foliis lanceolatis obtusis: caule
tuberculato decumbente.

In Ducatibus obvia.

* * *Spinosa.*

570. † GENISTA *Germanica* spinis compositis: ramis
floriferis inermibus: foliis lanceolatis.

In ericetis *Holfatiæ* passim.

260. CORONILLA. 883.

571. † CORONILLA *varia* herbacea, leguminibus ere-
ctis teretibus torosis numerosis: foliolis plurimis
glabris.

261. ONONIS. 863.

572. ONONIS *spinosa* floribus racemosis geminis, fo-
liis ternatis: superioribus solitariis: ramis inermi-
bus subvillosis.

Dan. *Krage-Torn.*

Pharmac. ONONIDIS radix, herba.

In campestri frequens.

262. ANTHYLLIS. 864.

73. ANTHYLLIS *vulneraria* herbacea: foliis quaternis pinnatis: floribus lateralibus.

Dan. *Katte - Klever.*

In sterilibus.

263. PISUM. 870.

74. PISUM *maritimum* petiolis supra planiusculis: caule angulato: stipulis sagittatis: pedunculis multifloris.

Morif. Hist. t. I. f. 5.

In maritimis.

264. OROBUS. 871.

75. OROBUS *vernus* foliis pinnatis ovatis: stipulis semisagittatis integerrimis: caule simplici.

Dan. *Muus - Erter.*

Pharmac. OROBI farina.

In montosis nemorum.

76. * OROBUS *tuberosus* foliis pinnatis lanceolatis: stipulis semisagittatis integerrimis: caule simplici.

In Sylvis.

77. * OROBUS *niger* caule ramoso: foliis sexjugis ovato-oblongis.

In montosis sylvaticis.

265. LATHYRUS. 872.

* Pedunculis *unifloris*.

78. † LATHYRUS *Aphaca* pedunculis unifloris: circa his aphyllis: stipulis sagittato-cordatis.

Dan. *Vikker.*

* * Pedunculis multifloris.

579. † LATHYRUS *tuberosus* pedunculis multifloris
cirrhis diphyllis: foliolis ovalibus: internodiis
nudis. *P. Fl. Oec.* 89.

Dan. *Jord - Rödder*; *Heede - Nödder*.

580. LATHYRUS *pratensis* pedunculis multifloris: cir-
rhis diphyllis simplicissimis: foliolis lanceolatis.

Dan. *Vilde Erter*.

In dumetis.

581. LATHYRUS *sylvestris* pedunculis multifloris: cir-
rhis diphyllis: foliolis ensiformibus: internodiis
membranaceis.

Extra hortum ac in declivi *Brankebakke*.

582. LATHYRUS *latifolius* pedunculis multifloris: cirrhis
diphyllis: foliolis lanceolatis: internodiis mem-
branaceis.

583. * LATHYRUS *palustris* pedunculis multifloris
cirrhis polyphyllis: stipulis lanceolatis.

In pascuis humidis.

266. V I C I A. 873.

* Pedunculis elongatis.

584. VICIA *dumetorum* pedunculis multifloris: foliolis
reflexis ovatis mucronatis: stipulis subdentatis.

Dan. *Vikker*.

585. † VICIA *sylvatica* pedunculis multifloris: foliolis
ovalibus: stipitulis denticulatis.

586. VICIA *Cracca* pedunculis multifloris: floribus im-
bricatis: foliolis lanceolatis pubescentibus: stipu-
lis integris.

Dan. *Muus - Erter*.

Inter segetes, ac in nemore versus amnem.

* * Floribus axillaribus subsessilibus.

7. *VICIA sativa* leguminibus sessilibus subbinatis erectis: foliolis retusis: stipulis notatis.

Inter segetes.

8. *VICIA Lathyroides* leguminibus sessilibus solitariis, erectis, glabris: foliolis senis: inferioribus obcordatis. *Fl. Dan. t. 58.*

In aridis montosis, ac in agris passim.

9. *VICIA sepium* leguminibus pedicellatis subquaternis erectis: foliolis ovatis integerrimis: exterioribus decrescentibus.

In umbrosis.

267. ERVUM. 874.

90. *ERVUM tetraspermum* pedunculis subbifloris: seminibus globosis quaternis. *Fl. Dan. t. 95.*

In oleraceis frequens.

91. *ERVUM hirsutum* pedunculis multifloris: seminibus globosis binis.

In arvis, ac in horto frequens.

268. ORNITHOPUS. 884.

92. † *ORNITHOPUS perpusillus* foliis pinnatis: leguminibus subarcuatis.

269. HEDYSARUM. 887.

93. † *HEDYSARUM Onobrychis* foliis pinnatis: leguminibus monospermis aculeatis: corollarum alis calyce brevioribus: caule elongato.

270. ASTRAGALUS. 892.

- §94. ASTRAGALUS *glycyphyllus* caulescens prostratus leguminibus subtriquetris arcuatis: foliis ovalibus pedunculo longioribus. *Morif. Hist. III. t. 9. f. 1.*
 Dan. *Vild Lakrits.*

In sylvatico *Deyligheden* frequens, & alibi.

- §95. † ASTRAGALUS *arenarius* subcaulescens, procumbens, floribus subracemosis erectis: foliis tomentosis.

Dan. *Bierg - Lakrits.*

271. TRIFOLIUM. 896.

* *Meliloti leguminibus nudis polyspermis.*

- §96. TRIFOLIUM *M. officinalis* leguminibus racemosis nudis, dispermis, rugosis, acutis: caule erecto.

Dan. *Amur; Steen - Klever.*

Pharmac. MELILOTI herba, flores, semina.

In *Fionia* vulgaris; extra *Haderslebiam* corollas omnibus nitide albis legi.

In campestribus, imprimis inter *Molendina*.

** *Lotoïdea leguminibus testis polyspermis.*

- §97. † TRIFOLIUM *hybridum* capitulis umbellaribus leguminibus tetraspermis: caule adscendente.

Caulis teres, subpilosus, non sulcatus.

In via ultra cauponam *Kiøge - Kroe* vulgo frequentissimam legi.

- §98. TRIFOLIUM *repens* capitulis umbellaribus leguminibus tetraspermis: caule repente.

Dan. *Huid - Klever.*

Pharmac. TRIFOLII ALBI flores.

In pascuis.

* * * *Lagopoda calycibus villosis.*

599. *TRIFOLIUM pratense* spicis subvillosis cinctis: stipulis oppositis membranaceis: corollis monope-
talis. *P. Fl. Oec.* 146.

Dan. *Sue - Kop; Rød - Klever; Sukker - Brød.*

In pratiis ubique.

600. *TRIFOLIUM arvense* spicis villosis ovalibus: den-
tibus calycinis fetaceis æqualibus.

Dan. *Hare - Klever.*

In arvis.

* * * * *Vesicaria calycibus inflatis ventricosis.*

601. † *TRIFOLIUM spumosum* spicis ovatis: calycibus
inflatis glabris quinque dentatis: involucris uni-
versalibus pentaphyllis.

602. *TRIFOLIUM fragiferum* capitulis subrotundis:
calycibus inflatis bidentatis reflexis: caulibus re-
pentibus. *Vaill. Bot. Paris. t. 22. f. 2.*

* * * * * *Lupulina vexillis corollæ inflexis.*

603. † *TRIFOLIUM montanum* spicis subimbricatis
subtribus: vexillis subulatis emarcescentibus: ca-
lycibus nudis: caule erecto.

604. † *TRIFOLIUM agrarium* spicis ovalibus imbricatis: vexillis deflexis persistentibus: calycibus
nudis: caule erecto.

Dan. *Humble - Klever.*

272 LOTUS 897.

* Leguminibus prioribus nec capitulum
conspicuum.

605. † *LOTUS maritimus* leguminibus solitariis mem-
branaceo-quadrangulatis: foliis glabris bracteis
lanceolatis.

Dan. *Steen - Klever.*

* * Pedunculis *multifloris in capitulum*.

606. *LOTUS corniculatus* capitulis depresso: caulibus decumbentibus; leguminibus cylindricis patentibus. *P. Fl. Oec.* 95.

Dan. *Kierlinge - Tander*; *Krage - Klover*.

In pratis ac collibus.

273. M E D I C A G O. 899.

607. *MEDICAGO falcata* pedunculis racemosis: leguminibus lunatis: caule prostrato.

Fl. Dan. t. 233. *P. Fl. Oec.* 97.

Dan. *Gnul - Klover*.

In Icone legumina desunt, caracter plantæ specificus.

In agris ad vias.

608. † *MEDICAGO lupulina* spicis ovalibus: leguminibus reniformibus monospermis: caule procumbente.

In pratis & ad vias.

609. † *MEDICAGO polymorpha* minima leguminibus cochleatis aculeis uncinatis alternis externe donatis: stipulis integris. *Fl. Dan. t.* 211.



Classis XVIII.

POLYADELPHIA.

POLYANDRIA.

274. HYPERICUM. *Cal.* 5 - partitus. *Cor.* 5 - petala.
Filamenta multa in 5 phalangibus. *Styli* 1, 2, 3, 4, 5.
Caps. locularis.



Stigma pubescens, (nec priorum notæ.)

264. OROBUS. *Stylus* linearis, teretiusculus, supra villosus. *Cal.* basi obtusus: *Lacinis* superioribus profundioribus, brevioribus.
265. PISUM. *Stylus* triangulus supra carinatus villosusque. *Calycis* lacinie superiores 2 breviores.
266. LATHYRUS. *Stylus* planus, supra villosus, superne lator. *Calycis* lacinie superiores 2 breviores.
266. VICIA. *Stylus* sub stigmate barbatus.

Legumen subbiloculare (nec priorum.)

270. ASTRAGALUS. *Legumen* biloculare, rotundatum, gibbum.

Legumina submonosperma (nec priorum.)

271. TRIFOLIUM. *Legumen* vix calyce longius, 1 - seu 2 - spermum, non dehiscens, deciduum. *Flores* capitati.

Legumen subarticulatum.

269. HEDYSARUM. *Legumen* articulis subrotundis, compressis, 1 - spermis. *Cor.* carina obtusissima.
260. CORONILLA. *Legumen* isthmis interceptum, rectum. *Cal.* 2-labiatus: $\frac{2}{3}$. dentibus superioribus connatis. *Vexillum* vix alis longius.
268. ORNITHOPUS. *Legumen* articulatum, teres, arcuatum.
273. MEDICAGO. *Legumen* spirale, membranaceo-compressum. *Carina* corollæ a vexillo deflectens.

Legumen uniloculare, polyspermum (nec priorum.)

267. ERVUM. *Calyx* 5-partitus, subæqualis longitudine fere corollæ.
272. LOTUS. *Legumen* cylindricum, strictum, factum seminibus cylindricis. Alæ sursum longitudinaliter conniventes. *Cal.* tubulosus.

HEXANDRIA.

HEXANDRIA.

256. FUMARIA. 849.

564. FUMARIA *bulbosa* caule simplici: bracteis longitudine florum.

α) Fumaria bulbosa radice cava, major.

β) Fumaria bulbosa radice non cava, minor.

Morif. Hist. p. 261, t. 12. f. 8.

Dan. *Hvul-Roed*; *Röd Hane-Kamm*.

Pharmac. ARISTOLOCHIÆ FABACEÆ radix.

α) Frequens in nemorosis ad lacum *Weselöfe*.

β) Frequens in Vivario *Friderichsdalensi*.

565. FUMARIA *officinalis* pericarpis monospermis racemosis: caule diffuso.

Dan. *Jordrög*.

Pharmac. FUMARIÆ herba, semina.

In arvis & fatis frequens.

OCTANDRIA.

257. POLYGALA. 851.

566. POLYGALA *vulgaris* floribus cristatis racemosis: caulibus herbaceis simplicibus procumbentibus: foliis lineari-lanceolatis.

Vaill. Bot. Paris. t. 32. f. 2.

Dan. *Kaars-Blomster*.

In siccis vulgaris.

β) POLYGALA *violacea*.

γ) POLYGALA *carnea*.

δ) POLYGALA *alba*.

ε) POLYGALA *major*. Vaill. Bot. Paris. t. 32. f. 2.

DECANDRIA.

258. SPARTIUM. 88.

567. SPARTIUM *scoparium* foliis ternatis solitariisque:
ramis inermibus angulatis.
In arenosis fylvarum.

259. GENISTA. 859.

* *Inermes.*

568. † GENISTA *tinctoria* foliis lanceolatis glabris:
ramis striatis teretibus erectis.
Pharmac. GENISTÆ herba, flores, femina.
569. † GENISTA *pilosa* foliis lanceolatis obtusis: caule
tuberculato decumbente.
In Ducatibus obvia.

* * *Spinosa.*

570. † GENISTA *Germanica* spinis compositis: ramis
floriferis inermibus: foliis lanceolatis.
In ericetis *Holsatiae* passim.

260. CORONILLA. 883.

571. † CORONILLA *varia* herbacea, leguminibus ere-
ctis teretibus torosis numerosis: foliolis plurimis
glabris.

261. ONONIS. 863.

572. ONONIS *spinosa* floribus racemosis geminis, fo-
liis ternatis: superioribus solitariis: ramis inermi-
bus subvillosis.
Dan. Krage-Torn.
Pharmac. ONONIDIS radix, herba.
In campestri frequens.

262. ANTHYLLIS. 864.

3. ANTHYLLIS *vulneraria* herbacea: foliis quaternis pinnatis: floribus lateralibus.

Dan. *Katte - Klever.*

In sterilibus.

263. PISUM. 870.

4. PISUM *maritimum* petiolis supra planiusculis: caule angulato: stipulis sagittatis: pedunculis multifloris.

Morif. Hist. t. 1. f. 5.

In maritimis.

264. OROBUS. 871.

5. OROBUS *vernus* foliis pinnatis ovatis: stipulis semisagittatis integerrimis: caule simplici.

Dan. *Muus - Erter.*

Pharmac. OROBI farina.

In montosis nemorum.

6. * OROBUS *tuberosus* foliis pinnatis lanceolatis: stipulis semisagittatis integerrimis: caule simplici.

In Sylvis.

7. * OROBUS *niger* caule ramoso: foliis sexjugis ovato-oblongis.

In montosis sylvaticis.

265. LATHYRUS. 872.

* Pedunculis *unifloris*.

8. † LATHYRUS *Aphaca* pedunculis unifloris: circumsistis aphyllis: stipulis sagittato-cordatis.

Dan. *Vikker.*

* * *Pedunculis multifloris.*

579. † *LATHYRUS tuberosus* pedunculis multifloris: cirrhis diphyllis: foliolis ovalibus: internodiis nudis. *P. Fl. Oec.* 89.

Dan. *Jord - Rødder*; *Heede - Nødder.*

580. *LATHYRUS pratensis* pedunculis multifloris: cirrhis diphyllis simplicissimis: foliolis lanceolatis.

Dan. *Vilde Erter.*

In dumetis.

581. *LATHYRUS sylvestris* pedunculis multifloris: cirrhis diphyllis: foliolis ensiformibus: internodiis membranaceis.

Extra hortum ac in declivi *Brankebakke.*

582. *LATHYRUS latifolius* pedunculis multifloris: cirrhis diphyllis: foliolis lanceolatis: internodiis membranaceis.

583. * *LATHYRUS palustris* pedunculis multifloris: cirrhis polyphyllis: stipulis lanceolatis.

In pascuis humidis.

266. V I C I A. 873.

* *Pedunculis elongatis.*

584. *VICIA dumetorum* pedunculis multifloris: foliolis reflexis ovatis mucronatis: stipulis subdentatis.

Dan. *Vikker.*

585. † *VICIA sylvatica* pedunculis multifloris: foliolis ovalibus: stipitulis denticulatis.

586. *VICIA Cracca* pedunculis multifloris: floribus imbricatis: foliolis lanceolatis pubescentibus: stipulis integris.

Dan. *Muus - Erter.*

Inter segetes, ac in nemore versus amnem.

* * Floribus axillaribus subsessilibus.

17. *VICIA sativa* leguminibus sessilibus subbinatis erectis: foliolis retusis: stipulis notatis.

Inter segetes.

38. *VICIA Lathyroides* leguminibus sessilibus solitariis, erectis, glabris: foliolis senis: inferioribus obcordatis. *Fl. Dan. t. 58.*

In aridis montosis, ac in agris passim.

89. *VICIA sepium* leguminibus pedicellatis subquaternis erectis: foliolis ovatis integerrimis: exterioribus decescentibus.

In umbrosis.

267. ERVUM. 874.

90. *ERVUM tetraspermum* pedunculis subbifloris: seminibus globosis quaternis. *Fl. Dan. t. 95.*

In oleraceis frequens.

91. *ERVUM hirsutum* pedunculis multifloris: seminibus globosis binis.

In arvis, ac in horto frequens.

268. ORNITHOPUS. 884.

92. † *ORNITHOPUS perpusillus* foliis pinnatis: leguminibus subarcuatis.

269. HEDYSARUM. 887.

93. † *HEDYSARUM Onobrychis* foliis pinnatis: leguminibus monospermis aculeatis: corollarum alis calyce brevioribus: caule elongato.

DECANDRIA.

258. SPARTIUM. 88.

567. SPARTIUM *scoparium* foliis ternatis solitariisque:
ramis inermibus angulatis.

In arenosis sylvarum.

259. GENISTA. 859.

* *Inermes.*

568. † GENISTA *tinctoria* foliis lanceolatis glabris:
ramis striatis teretibus erectis.

Pharmac. GENISTÆ herba, flores, semina.

569. † GENISTA *pilosa* foliis lanceolatis obtusis : caule
tuberculato decumbente.

In Ducatibus obvia.

* * *Spinosa.*

570. † GENISTA *Germanica* spinis compositis : ramis
floriferis inermibus : foliis lanceolatis.

In ericetis *Holsatiae* passim.

260. CORONILLA. 883.

571. † CORONILLA *varia* herbacea, leguminibus ere-
ctis teretibus torosis numerosis : foliolis plurimis
glabris.

261. ONONIS. 863.

572. ONONIS *spinosa* floribus racemosis geminis, fo-
liis ternatis : superioribus solitariis : ramis inermi-
bus subvillosis.

Dan. Krage-Torn.

Pharmac. ONONIDIS radix, herba.

In campestri frequens.

262. ANTHYLLIS. 864.

73. ANTHYLLIS *vulneraria* herbacea: foliis quaternis - pinnatis: floribus lateralibus.

Dan. *Katte - Klever.*

In sterilibus.

263. PISUM. 870.

74. PISUM *maritimum* petiolis supra planiusculis: caule angulato: stipulis sagittatis: pedunculis multifloris.

Morif. Hist. t. 1. f. 5.

In maritimis.

264. OROBUS. 871.

75. OROBUS *vernus* foliis pinnatis ovatis: stipulis semisagittatis integerrimis: caule simplici.

Dan. *Muus - Erter.*

Pharmac. OROBI farina.

In montosis nemorum.

76. * OROBUS *tuberosus* foliis pinnatis lanceolatis: stipulis semisagittatis integerrimis: caule simplici.

In Sylvis.

77. * OROBUS *niger* caule ramoso: foliis sexjugis ovato - oblongis.

In montosis sylvaticis.

265. LATHYRUS. 872.

* Pedunculis *unifloris.*

78. † LATHYRUS *Aphaca* pedunculis unifloris: circa his aphyllis: stipulis sagittato - cordatis.

Dan. *Vikker.*

* * *Pedunculis multifloris.*

579. † *LATHYRUS tuberosus* pedunculis multifloris: cirrhis diphyllis: foliolis ovalibus: internodiis nudis. *P. Fl. Oec.* 89.

Dan. *Jord - Rödder; Heede - Nödder.*

580. *LATHYRUS pratensis* pedunculis multifloris: cirrhis diphyllis simplicissimis: foliolis lanceolatis.

Dan. *Vilde Erter.*

In dumetis.

581. *LATHYRUS sylvestris* pedunculis multifloris: cirrhis diphyllis: foliolis ensiformibus: internodiis membranaceis.

Extra hortum ac in declivi *Brankebakke.*

582. *LATHYRUS latifolius* pedunculis multifloris: cirrhis diphyllis: foliolis lanceolatis: internodiis membranaceis.

583. * *LATHYRUS palustris* pedunculis multifloris: cirrhis polyphyllis: stipulis lanceolatis.

In pascuis humidis.

266. *V I C I A.* 873.* *Pedunculis elongatis.*

584. *VICIA dumetorum* pedunculis multifloris: foliolis reflexis ovatis mucronatis: stipulis subdentatis.

Dan. *Vikker.*

585. † *VICIA sylvatica* pedunculis multifloris: foliolis ovalibus: stipitulis denticulatis.

586. *VICIA Cracca* pedunculis multifloris: floribus imbricatis: foliolis lanceolatis pubescentibus: stipulis integris.

Dan. *Muus - Erter.*

Inter segetes, ac in nemore versus amnem.

* * Floribus axillaribus subsessilibus.

87. *VICIA sativa* leguminibus sessilibus subbinatis erectis: foliolis retusis: stipulis notatis.

Inter segetes.

88. *VICIA Lathyroides* leguminibus sessilibus solitariis, erectis, glabris: foliolis senis: inferioribus obcordatis. *Fl. Dan. t. 58.*

In aridīs montosis, ac in agris passim.

89. *VICIA sepium* leguminibus pedicellatis subquaternis erectis: foliolis ovatis integerrimis: exterioribus decescentibus.

In umbrosis.

267. ERVUM. 874.

90. *ERVUM tetraspermum* pedunculis subbifloris: seminibus globosis quaternis. *Fl. Dan. t. 95.*

In oleraceis frequens.

91. *ERVUM hirsutum* pedunculis multifloris: seminibus globosis binis.

In arvis, ac in horto frequens.

268. ORNITHOPUS. 884.

92. † *ORNITHOPUS perpusillus* foliis pinnatis: leguminibus subarcuatis.

269. HEDYSARUM. 887.

93. † *HEDYSARUM Onobrychis* foliis pinnatis: leguminibus monospermis aculeatis: corollarum alis calyce brevioribus: caule elongato.

DECANDRIA.

253. GERANIUM. 832.

* *Staminibus quinque fertilibus.*

548. GERANIUM *cicutarium* pedunculis multifloris :
floribus pentandris : foliis pinnatis incisus obtusis.

Dan. *Storke-Nab.*

In nemore & ad aggeres.

* * *Staminibus decem fertilibus.*

α) Perennia.

549. GERANIUM *macrorrhizum* pedunculis bifloris :
calycibus inflatis : pistillo longissimo.

550. GERANIUM *sylvaticum* pedunculis bifloris : foliis
subpeltatis quinquelobis inciso-ferratis : caule ere-
cto : petalis emarginatis.

*Fl Dan. t. 124.*Dan. *Trane-Hals.*

In Nemore.

551. GERANIUM *palustre* pedunculis bifloris longissi-
mis declinatis, foliis quinquelobis incisus : petalis
integris.

552. † GERANIUM *pratense* pedunculis bifloris : foliis
subpeltatis multipartitis, pinnato-laciniatis rugo-
sis acutis : petalis integris.

β) Annua.

553. GERANIUM *Robertianum* pedunculis bifloris : ca-
lycibus pilosis decem angulatis. *P. Fl. Oec. 76.*

Dan. *Roberts - Urt.**Pharmac.* GERANII ROBERTIANI herba.

In ruderatis & ad muros.

554. † GERANIUM *lucidum* pedunculis bifloris: calycibus pyramidalis angulatis elevato-rugosis: foliis quinquelobis rotundatis. *Fl. Dan. t. 218.*

555. * GERANIUM *molle* pedunculis bifloris foliisque floralibus alternis: caule ramoso erectiusculo: calycibus muticis: arillis lævibus.

Vaill. Bot. Paris. t. 15. f. 3.

In nemore.

556. GERANIUM *difsectum* pedunculis bifloris: foliis quinquepartito-trifidis: petalis emarginatis longitudine calycis: arillis villosis.

Vaill. Bot. Paris. t. 15. f. 2.

Ad sepimenta.

557. * GERANIUM *columbinum* pedunculis bifloris folio longioribus: foliis quinquepartito-multifidis: laciniis acutis: arillis glabris: calycibus aristatis.

Vaill. Bot. Paris. t. 15. f. 4.

In arvis.

558. GERANIUM *retundifolium* pedunculis bifloris: petalis integris obtusissimis longitudine calycis: caule prostrato: foliis reniformibus incisis.

Vaill. Bot. Paris. t. 15. f. 1.

In cultis.

* * * Pedunculis *unifloris*.

559. GERANIUM *sanguineum* pedunculis unifloris: foliis quinquepartitis trifidis orbiculatis.

In dumetis ficcis collis *Brankebakke* frequens.

POLYANDRIA.

254. A L T H Æ A. 839.

560. † ALTHÆA *officinalis* foliis simplicibus tomentosis.
Dan. Althee.

255. MALVA. 841.

561. MALVA *rotundifolia* caule prostrato: foliis cordato-orbiculatis, obsolete quinquelobis: pedunculis fructiferis declinatis.

Dan. *Kat - Ost.*

In ruderatis & ad pagos.

562. MALVA *sylvestris* caule erecto herbaceo: foliis septemlobatis acutis: pedunculis petiolisque pilosis.

Pharmac. MALVÆ VULGARIS radix, herba, semina.
Ad vias.

563. † MALVA *Alcea* caule erecto: foliis multipartitis scabriusculis.

Dan. *Cimons - Urt.*



Classis XVII.

DIADELPHIA.

HEXANDRIA.

256. FUMARIA. *Cal.* 2-phyllus. *Cor.* ringens, basi gibbosa, nectarifera. *Filamenta* 2 membranacea, singula antheris 3.

OCTANDRIA.

257. POLYGALA. *Cal.* 5-phyllus: foliolis duobus alæformibus, coloratis. *Cor.* vexillum cylindricum. *Stamina* connexa. *Legumen* obcordatum, biloculare.

DECANDRIA.

Stamina omnia connexa.

258. SPARTIUM. *Stigma* longitudinale, supra villosum. *Filamenta* carinæ adhærentia. *Cal.* deorsum productus.
259. GENISTA. *Pistillum* deprimens carinam. *Stigma* involutum. *Cal.* 2-labiatus: $\frac{2}{3}$. *Vexillum* oblongum, a pistillo staminibusque deorsum reflexum.
262. ANTHYLLIS. *Cal.* ventricosus, turgidus, siliquam includens. *Legumen* subrotundum, tectum.
261. ONONIS. *Cal.* 5-partitus: laciniis linearibus. *Vexillum* striatum. *Legumen* turgidum, rhombeum, sessile.

Stigma pubescens, (nec priorum notæ.)

264. OROBUS. *Stylus* linearis, teretiusculus, supra villosus. *Cal.* basi obtusus: *Laciniis* superioribus profundioribus, brevioribus.
263. PISUM. *Stylus* triangulus supra carinatus villosusque. *Calycis* lacinia superiores 2 breviores.
265. LATHYRUS. *Stylus* planus, supra villosus, superne lator. *Calycis* lacinia superiores 2 breviores.
266. VICIA. *Stylus* sub stigmate barbatus.

Legumen subbiloculare (nec priorum.)

270. ASTRAGALUS. *Legumen* biloculare, rotundatum, gibbum.

Legumina submonosperma (nec priorum.)

271. TRIFOLIUM. *Legumen* vix calyce longius, 1 - seu 2 - spermum, non dehiscens, deciduum. *Flores* capitati.

Legumen subarticulatum.

269. HEDYSARUM. *Legumen* articulis subrotundis, compressis, 1 - spermis. *Cor.* carina obtusissima.
260. CORONILLA. *Legumen* isthmis interceptum, rectum. *Cal.* 2-labiatus: $\frac{2}{3}$. dentibus superioribus connatis. *Vexillum* vix alis longius.
268. ORNITHOPUS. *Legumen* articulatum, teres, arcuatum.
273. MEDICAGO. *Legumen* spirale, membranaceo-compressum. *Carina* corollæ a vexillo deflectens.

Legumen uniloculare, polyspermum (nec priorum.)

267. ERVUM. *Calyx* 5-partitus, subæqualis longitudine fere corollæ.
272. LOTUS. *Legumen* cylindricum, strictum, factum seminibus cylindricis. Alæ sursum longitudinaliter conniventes. *Cal.* tubulosus.

HEXANDRIA.

HEXANDRIA.

256. FUMARIA. 849.

564. FUMARIA *bulbosa* caule simplici: bracteis longitudine florum.

a) Fumaria bulbosa radice cava, major.

β) Fumaria bulbosa radice non cava, minor.

Morif. Hist. p. 261. t. 12. f. 8.

Dan. *Huul-Roed; Röd Hane-Kamm.*

Pharmac. ARISTOLOCHIE FABACEÆ radix.

a) Frequens in nemorosis ad lacum *Warelöse*.

β) Frequens in Vivario *Friderichsdalensi*.

565. FUMARIA *officinalis* pericarpis monospermis racemosis: caule diffuso.

Dan. *Jordrög.*

Pharmac. FUMARIÆ herba, semina.

In arvis & fatis frequens.

OCTANDRIA.

257. POLYGALA. 851.

566. POLYGALA *vulgaris* floribus cristatis racemosis: caulibus herbaceis simplicibus procumbentibus: foliis lineari-lanceolatis.

Vaill. Bot. Paris. t. 32. f. 2.

Dan. *Kaars-Blomster.*

In ficcis vulgaris.

β) POLYGALA *violacea*.

γ) POLYGALA *carnea*.

δ) POLYGALA *alba*.

ε) POLYGALA *major*. Vaill. Bot. Paris. t. 32. f. 2.

DECANDRIA.

258. SPARTIUM. 88.

567. SPARTIUM *scoparium* foliis ternatis solitariisque:
ramis inermibus angulatis.

In arenosis fylvarum.

259. GENISTA. 859.

* *Inermes.*

568. † GENISTA *tinctoria* foliis lanceolatis glabris:
ramis striatis teretibus erectis.

Pharmac. GENISTÆ herba, flores, femina.

569. † GENISTA *pilosa* foliis lanceolatis obtusis : cau-
tuberculato decumbente.

In Ducatibus obvia.

* * *Spinosa.*

570. † GENISTA *Germanica* spinis compositis : ramis
floriferis inermibus : foliis lanceolatis.

In ericetis *Holsatiae* passim.

260. CORONILLA. 883.

571. † CORONILLA *varia* herbacea, leguminibus ere-
ctis teretibus torosis numerosis : foliolis plurimis
glabris.

261. ONONIS. 863.

572. ONONIS *spinosa* floribus racemosis geminis, fo-
liis ternatis : superioribus solitariis : ramis inermi-
bus subvillosis.

Dan. *Krage-Torn.*

Pharmac. ONONIDIS radix, herba.

In campestri frequens.

262. ANTHYLLIS. 864.

3. ANTHYLLIS *vulneraria* herbacea: foliis quaternis: pinnatis: floribus lateralibus.

Dan. *Katte - Klever.*

In sterilibus.

263. PISUM. 870.

4. PISUM *maritimum* petiolis supra planiusculis: caule angulato: stipulis sagittatis: pedunculis multifloris.

Morif. Hist. t. 1. f. 5.

In maritimis.

264. OROBUS. 871.

5. OROBUS *vernus* foliis pinnatis ovatis: stipulis semisagittatis integerrimis: caule simplici.

Dan. *Muus - Erter.*

Pharmac. OROBI farina.

In montosis nemorum.

6. * OROBUS *tuberosus* foliis pinnatis lanceolatis: stipulis semisagittatis integerrimis: caule simplici.

In Sylvis.

7. * OROBUS *niger* caule ramoso: foliis sexjugis ovato-oblongis.

In montosis sylvaticis.

265. LATHYRUS. 872.

* Pedunculis *unifloris*.

8. † LATHYRUS *Aphaca* pedunculis unifloris: circumsistis aphyllis: stipulis sagittato-cordatis.

Dan. *Vikker.*

* * Pedunculis multifloris.

579. † LATHYRUS *tuberosus* pedunculis multifloris: cirrhis diphyllis: foliolis ovalibus: internodiis nudis. *P. Fl. Oec.* 89.

Dan. *Jord - Rødder*; *Heede - Nødder*.

580. LATHYRUS *pratensis* pedunculis multifloris: cirrhis diphyllis simplicissimis: foliolis lanceolatis.

Dan. *Vilde Erter*.

In dumetis.

581. LATHYRUS *sylvestris* pedunculis multifloris: cirrhis diphyllis: foliolis ensiformibus: internodiis membranaceis.

Extra hortum ac in declivi *Brankebakke*.

582. LATHYRUS *latifolius* pedunculis multifloris: cirrhis diphyllis: foliolis lanceolatis: internodiis membranaceis.

583. * LATHYRUS *palustris* pedunculis multifloris: cirrhis polyphyllis: stipulis lanceolatis.

In pascuis humidis.

266. V I C I A. 873.

* Pedunculis elongatis.

584. VICIA *dumetorum* pedunculis multifloris: foliolis reflexis ovatis mucronatis: stipulis subdentatis.

Dan. *Vikker*.

585. † VICIA *sylvatica* pedunculis multifloris: foliolis ovalibus: stipitulis denticulatis.

586. VICIA *Cracca* pedunculis multifloris: floribus imbricatis: foliolis lanceolatis pubescentibus: stipulis integris.

Dan. *Muus - Erter*.

Inter segetes, ac in nemore versus amnem.

* * Floribus axillaribus subsessilibus.

7. *VICIA sativa* leguminibus sessilibus subbinatis erectis : foliolis retusis : stipulis notatis.

Inter segetes.

38. *VICIA Lathyroides* leguminibus sessilibus solitariis, erectis, glabris : foliolis senis : inferioribus obcordatis. *Fl. Dan. t. 58.*

In aridis montosis, ac in agris passim.

89. *VICIA sepium* leguminibus pedicellatis subquaternis erectis : foliolis ovatis integerrimis : exterioribus decreescentibus.

In umbrosis.

267. ERVUM. 874.

90. *ERVUM tetraspermum* pedunculis subbifloris : seminibus globosis quaternis. *Fl. Dan. t. 95.*

In oleraceis frequens.

91. *ERVUM hirsutum* pedunculis multifloris : seminibus globosis binis.

In arvis, ac in horto frequens.

268. ORNITHOPUS. 884.

92. † *ORNITHOPUS perpusillus* foliis pinnatis : leguminibus subarcuatis.

269. HEDYSARUM. 887.

93. † *HEDYSARUM Onobrychis* foliis pinnatis : leguminibus monospermis aculeatis : corollarum alis calyce brevioribus : caule elongato.

270. ASTRAGALUS. 892.

594. ASTRAGALUS *glycyphyllus* caulescens prostratus: leguminibus subtriquetris arcuatis: foliis ovalibus pedunculo longioribus. *Morif. Hist. III. t. 9. f. 8.*
 Dan. *Vild Lakrits.*

In sylvatico *Deyligheden* frequens, & alibi.

595. † ASTRAGALUS *arenarius* subcaulescens, procumbens, floribus subracemosis erectis: foliis tomentosis.

Dan. *Bierg - Lakrits.*

271. TRIFOLIUM. 896.

* *Meliloti leguminibus nudis polyspermis.*

596. TRIFOLIUM *M. officinalis* leguminibus racemosis nudis, dispermis, rugosis, acutis: caule erecto.

Dan. *Amur; Steen - Klever.*

Pharmac. MELILOTI herba, flores, femina.

In *Fionia* vulgaris; extra *Haderslebiam* corollis omnibus nitide albis legi.

In campestribus, imprimis inter *Molendina*.

** *Lotoïdea leguminibus testis polyspermis.*

597. † TRIFOLIUM *hybridum* capitulis umbellaribus: leguminibus tetraspermis: caule adscendente.

Caulis teres, subpilosus, non sulcatus.

In via ultra cauponam *Kiøge - Kroe* vulgo frequentissimam legi.

598. TRIFOLIUM *repens* capitulis umbellaribus: leguminibus tetraspermis: caule repente.

Dan. *Huid - Klever.*

Pharmac. TRIFOLII ALBI flores.

In pascuis.

* * * *Lagopoda calycibus villosis.*

599. *TRIFOLIUM pratense* spicis subvillosis cinctis: stipulis oppositis membranaceis: corollis monopetalis. *P. Fl. Oec.* 146.

Dan. *Sue - Kop; Rød - Klever; Sukker - Brød.*

In pratis ubique.

600. *TRIFOLIUM arvense* spicis villosis ovalibus: dentibus calycinis setaceis æqualibus.

Dan. *Hare - Klever.*

In arvis.

* * * * *Vesicaria calycibus inflatis ventricosis.*

601. † *TRIFOLIUM spumosum* spicis ovatis: calycibus inflatis glabris quinque dentatis: involucris universalibus pentaphyllis.

602. *TRIFOLIUM fragiferum* capitulis subrotundis: calycibus inflatis bidentatis reflexis: cauhibus repentibus. *Vaill. Bot. Paris. t. 22. f. 2.*

* * * * * *Lupulina vexillis corollæ inflexis.*

603. † *TRIFOLIUM montanum* spicis subimbricatis subtribus: vexillis subulatis emarcescentibus: calycibus nudis: caule erecto.

604. † *TRIFOLIUM agrarium* spicis ovalibus imbricatis: vexillis deflexis persistentibus: calycibus nudis: caule erecto.

Dan. *Humble - Klever.*

272. *LOTUS.* — 897.

* *Leguminibus rarioribus, nec capitulum constituentibus.*

605. † *LOTUS maritimus* leguminibus solitariis membranaceo-quadrangulatis: foliis glabris bracteis lanceolatis.

Dan. *Steen - Klever.*

* * Pedunculis *multifloris in capitulum.*

606. *LOTUS corniculatus* capitulis depressis: caulibus decumbentibus: leguminibus cylindricis patentibus. *P. Fl. Oec. 95.*

Dan. *Kierlinge - Tender; Krage - Klover.*
In pratibus ac collibus.

273. M E D I C A G O. 899.

607. *MEDICAGO falcata* pedunculis racemosis: leguminibus lunatis: caule prostrato.

Fl. Dan. t. 233. P. Fl. Oec. 97.

Dan. *Gaul - Klover.*

In Icone legumina desunt, character plantæ specificus.
In agris ad vias.

608. † *MEDICAGO lupulina* spicis ovalibus: leguminibus reniformibus monospermis: caule procumbente.

In pratibus & ad vias.

609. † *MEDICAGO polymorpha* minima leguminibus cochleatis aculeis uncinatis alternis externe donatis: stipulis integris. *Fl. Dan. t. 211.*



Classis XVIII.

POLYADELPHIA.

POLYANDRIA.

274. HYPERICUM. *Cal.* 5 - partitus. *Cor.* 5 - petala.
Filamenta multa in 5 phalangibus. *Styli* 1, 3, 5.
Caps. locularis.



POLYANDRIA.

274. HYPERICUM. 902.

610. HYPERICUM *quadrangulum* floribus trigynis :
caule quadrato herbaceo.

Dan. *Piricum*.

In pratis.

611. HYPERICUM *perforatum* floribus trigynis : caule
ancipiti, foliis obtusis pellucido - punctatis.

P. Fl. Occ. 85.

Dan. *Hypericon*; *Jord-Humle*; *Blod-Blømsker*.

Pharmac. HYPERICI herba, flores, semina.

612. HYPERICUM *humifusum* floribus trigynis axilla-
ribus solitariis : caulibus ancipitibus prostratis fili-
formibus : foliis glabris. Fl. Dan. t. 141.

In arvis haud infrequens.

613. HYPERICUM *hirsutum* floribus trigynis : caly-
bus ferrato - glandulosis : caule tereti erecto : fo-
liis ovatis subpubescentibus.

Morif. Hist. III. t. 6. f. 11.

In montosis minus frequens.

614. * HYPERICUM *montanum* floribus trigynis : ca-
lycibus ferraturis glandulosis : caule tereti erecto
glabro : foliis ovatis. Fl. Dan. t. 173.

Passim : ad molendina haud infrequens.

615. † HYPERICUM *pulchrum* floribus trigynis : caly-
cibus ferrato - glandulosis : caule tereti : foliis per-
foliatis, cordatis, glabris. Fl. Dan. t. 75.

In insula *Thorſinge* legi.



Classis XIX.

SYNGENESIA.

POLYGAMIA AQUALIS.

Semiflosculosi: Corollis ligulatis omnibus.

- 287. CICHORIUM. *Receptaculum paleaceum. Pappus 5-dentatus. Calyx calyculatus.*
- 285. HYPOCHERIS. *Recept. paleaceum. Pappus subplumosus. Calyx subimbricatus.*
- 275. TRAGOPOGON. *Recept. nudum. Pappus plumosus. Cal. simplex polyphyllus.*
- 277. PICRIS. *Recept. nudum. Pappus plumosus. Cal. calyculatus. Semina transversim fulcata.*
- 281. LEONTODON. *Recept. nudum. Pappus plumosus. Cal. imbricatus. Squamis laxis.*
- 278. SONCHUS. *Recept. nudum. Pappus plumosus. Cal. imbricatus, basi gibbus.*
- 276. SCORZONERA. *Recept. nudum. Pappus plumosus. Cal. imbricatus. Squamis margine membranaceis.*
- 283. CREPIS. *Recept. nudum. Pappus pilosus, stipitatus. Cal. calyculatus. Squamis deciduis.*
- 280. PRENANTHES. *Recept. nudum. Pappus pilosus, simplex, subsessilis. Cal. calyculatus sub-5-florus.*
- 279. LACTUCA. *Recept. nudum. Pappus pilosus, simplex, stipitatus. Cal. imbricatus, cylindricus.*
- 282. HIERACIUM. *Recept. nudum. Pappus pilosus, simplex, sessilis. Cal. imbricatus, ovatus.*
- 286. LAPSANA. *Recept. nudum. Pappus nonnullus nullus. Cal. calyculatus. Squamis singulis interioribus canaliculatis.*

DECANDRIA.

258. SPARTIUM. 88.

567. SPARTIUM *scoparium* foliis ternatis solitariisque:
ramis inermibus angulatis.

In arenosis fylvarum.

259. GENISTA. 859.

* *Inermes.*

568. † GENISTA *tinctoria* foliis lanceolatis glabris:
ramis striatis teretibus erectis.

Pharmac. GENISTÆ herba, flores, semina.

569. † GENISTA *pilosa* foliis lanceolatis obtusis: caule
tuberculato decumbente.

In Ducatibus obvia.

* * *Spinosa.*

570. † GENISTA *Germanica* spinis compositis: ramis
floriferis inermibus: foliis lanceolatis.

In ericetis *Holsatiæ* passim.

260. CORONILLA. 883.

571. † CORONILLA *varia* herbacea, leguminibus ere-
ctis teretibus torosis numerosis: foliolis plurimis
glabris.

261. ONONIS. 863.

572. ONONIS *spinosa* floribus racemosis geminis, fo-
liis ternatis: superioribus solitariis: ramis inermi-
bus subvillosis.

Dan. *Krage-Torn.*

Pharmac. ONONIDIS radix, herba.

In campestri frequens.

262. ANTHYLLIS. 864.

73. ANTHYLLIS *vulneraria* herbacea: foliis quaternis: pinnatis: floribus lateralibus.

Dan. *Katte - Klever.*

In sterilibus.

263. PISUM. 870.

74. PISUM *maritimum* petiolis supra planiusculis: caule angulato: stipulis sagittatis: pedunculis multifloris.

Morif. Hist. t. I. f. 5.

In maritimis.

264. OROBUS. 871.

75. OROBUS *vernus* foliis pinnatis ovatis: stipulis semisagittatis integerrimis: caule simplici.

Dan. *Muus - Erter.*

Pharmac. OROBI farina.

In montosis nemorum.

76. * OROBUS *tuberosus* foliis pinnatis lanceolatis: stipulis semisagittatis integerrimis: caule simplici.

In Sylvis.

77. * OROBUS *niger* caule ramoso: foliis sexjugis ovato-oblongis.

In montosis sylvaticis.

265. LATHYRUS. 872.

* Pedunculis *unifloris*.

78. † LATHYRUS *Aphaca* pedunculis unifloris: circumsistis aphyllis: stipulis sagittato-cordatis.

Dan. *Vikker.*

* * *Pedunculis multifloris.*

579. † *LATHYRUS tuberosus* pedunculis multifloris: cirrhis diphyllis: foliolis ovalibus: internodiis nudis. *P. Fl. Oec.* 89.

Dan. *Jord - Rødder*; *Heede - Nødder*.

580. *LATHYRUS pratensis* pedunculis multifloris: cirrhis diphyllis simplicissimis: foliolis lanceolatis.

Dan. *Vilde Erter*.

In dumetis.

581. *LATHYRUS sylvestris* pedunculis multifloris: cirrhis diphyllis: foliolis ensiformibus: internodiis membranaceis.

Extra hortum ac in declivi *Brankebakke*.

582. *LATHYRUS latifolius* pedunculis multifloris: cirrhis diphyllis: foliolis lanceolatis: internodiis membranaceis.

583. * *LATHYRUS palustris* pedunculis multifloris: cirrhis polyphyllis: stipulis lanceolatis.

In pascuis humidis.

266. *V I C I A.* 873.* *Pedunculis elongatis.*

584. *VICIA dumetorum* pedunculis multifloris: foliolis reflexis ovatis mucronatis: stipulis subdentatis.

Dan. *Vikker*.

585. † *VICIA sylvatica* pedunculis multifloris: foliolis ovalibus: stipitulis denticulatis.

586. *VICIA Cracca* pedunculis multifloris: floribus imbricatis: foliolis lanceolatis pubescentibus: stipulis integris.

Dan. *Muus - Erter*.

Inter segetes, ac in nemore versus amnem.

* * Floribus axillaribus subsessilibus.

87. *VICIA sativa* leguminibus sessilibus subbinatis erectis: foliolis retusis: stipulis notatis.

Inter segetes.

88. *VICIA Lathyroides* leguminibus sessilibus solitariis, erectis, glabris: foliolis senis: inferioribus obcordatis. *Fl. Dan. t. 58.*

In aridis montosis, ac in agris passim.

89. *VICIA sepium* leguminibus pedicellatis subquaternis erectis: foliolis ovatis integerrimis: exterioribus decreescentibus.

In umbrosis.

267. ERVUM. 874.

90. *ERVUM tetraspermum* pedunculis subbifloris: seminibus globosis quaternis. *Fl. Dan. t. 95.*

In oleraceis frequens.

91. *ERVUM hirsutum* pedunculis multifloris: seminibus globosis binis.

In arvis, ac in horto frequens.

268. ORNITHOPUS. 884.

92. † *ORNITHOPUS perpusillus* foliis pinnatis: leguminibus subarcuatis.

269. HEDYSARUM. 887.

93. † *HEDYSARUM Onobrychis* foliis pinnatis: leguminibus monospermis aculeatis: corollarum alis calyce brevioribus: caule elongato.

270. ASTRAGALUS. 892.

594. ASTRAGALUS *glycyphyllus* caulescens prostratus
leguminibus subtriquetris arcuatis: foliis ovalibus
pedunculo longioribus. *Morif. Hist. III. t. 9. f. 8.*

Dan. *Vild Lakrits.*

In sylvatico *Deyligheden* frequens, & alibi.

595. † ASTRAGALUS *arenarius* subcaulescens, pro-
cumbens, floribus subracemosis erectis: foliis co-
mentosis.

Dan. *Bierg - Lakrits.*

271. TRIFOLIUM. 896.

* *Meliloti leguminibus nudis polyspermis.*

596. TRIFOLIUM *M. officinalis* leguminibus racemosis
nudis, dispermis, rugosis, acutis: caule erecto.

Dan. *Amur; Steen - Klever.*

Pharmac. MELILOTI herba, flores, femina.

In *Fionia* vulgaris; extra *Haderslebiam* corollae
omnibus nitide albis legi.

In campestribus, imprimis inter *Molendina*.

** *Lotoidea leguminibus testis polyspermis.*

597. † TRIFOLIUM *hybridum* capitulis umbellaribus
leguminibus tetraspermis: caule adscendente.

Caulis teres, subpilosus, non sulcatus.

In via ultra cauponam *Kiøge - Kroe* vulgo frequen-
tissimam legi.

598. TRIFOLIUM *repens* capitulis umbellaribus: legu-
minibus tetraspermis: caule repente.

Dan. *Huid - Klev.*

Pharmac. TRIFOLIUM

In pascuis.

* * * *Lagopoda calycibus villosis.*

99. *TRIFOLIUM pratense* spicis subvillosis cinctis: stipulis oppositis membranaceis: corollis monopetalis. *P. Fl. Oec.* 146.

Dan. *Sue - Kop; Röd - Klever; Sukker - Bröd.*

In pratis ubique.

100. *TRIFOLIUM arvense* spicis villosis ovalibus: dentibus calycinis fetaceis æqualibus.

Dan. *Hare - Klever.*

In arvis.

* * * * *Vesicaria calycibus inflatis ventricosis.*

101. † *TRIFOLIUM spumosum* spicis ovatis: calycibus inflatis glabris quinque dentatis: involucris universalibus pentaphyllis.

102. *TRIFOLIUM fragiferum* capitulis subrotundis: calycibus inflatis bidentatis reflexis: caulibus repentibus. *Vaill. Bot. Paris. t. 22. f. 2.*

* * * * * *Lupulina vexillis corollæ inflexis.*

103. † *TRIFOLIUM montanum* spicis subimbricatis subtribus: vexillis subulatis emarcescentibus: calycibus nudis: caule erecto.

104. † *TRIFOLIUM agrarium* spicis ovalibus imbricatis: vexillis deflexis persistentibus: calycibus nudis: caule erecto.

Dan. *Humle - Klever.*

272. LOTUS. — 897.

* Leguminibus rarioribus, nec capitulum constituentibus.

105. † *LOTUS maritimus* leguminibus solitariis membranaceis quadrangulatis: foliis glabris bracteis

Klever.

* * Floribus axillaribus subsessilibus.

1. *Vicia sativa* leguminibus sessilibus subbinatis erectis: foliolis retusis: stipulis notatis. inter segetes.

2. *Vicia Lathyroides* leguminibus sessilibus solitariis, erectis, glabris: foliolis senis: inferioribus obcordatis. *Fl. Dan. t. 58.*
In aridis montosis, ac in agris passim.

3. *Vicia sepium* leguminibus pedicellatis subquaternis erectis: foliolis ovatis integerrimis: exterioribus decrescentibus.
In umbrosis.

267. ERVUM. 874.

10. *ERVUM tetraspermum* pedunculis subbifloris: feminibus globosis quaternis. *Fl. Dan. t. 95.*
In oleraceis frequens.

91. *ERVUM hirsutum* pedunculis multifloris: feminibus globosis binis.
In arvis, ac in horto frequens.

268. ORNITHOPUS. 884.

92. † *ORNITHOPUS perpusillus* foliis pinnatis: leguminibus subarcuatis.

269. HEDYSARUM. 887.

93. † *HEDYSARUM Onobrychis* foliis pinnatis: leguminibus monospermis aculeatis: corollarum alis calyce brevioribus: caule elongato.

DECANDRIA.

253. GERANIUM. 832.

* *Staminibus quinque fertilibus.*

548. GERANIUM *cicutarium* pedunculis multifloris :
floribus pentandris : foliis pinnatis incisus obtusis.

Dan. *Storke-Nab.*

In nemore & ad aggeres.

* * *Staminibus decem fertilibus.*

a) *Perennia.*

549. GERANIUM *macrorrhizum* pedunculis bifloris :
calycibus inflatis : pistillo longissimo.

550. GERANIUM *sylvaticum* pedunculis bifloris : foliis
subpeltatis quinquelobis inciso-ferratis : caule ere-
cto : petalis emarginatis.

Fl Dan. t. 124.

Dan. *Trane-Hals.*

In Nemore.

551. GERANIUM *palustre* pedunculis bifloris longissi-
mis declinatis, foliis quinquelobis incisus : petalis
integris.

552. † GERANIUM *pratense* pedunculis bifloris : foliis
subpeltatis multipartitis, pinnato-laciniatis rugo-
sis acutis : petalis integris.

β) *Annua.*

553. GERANIUM *Robertianum* pedunculis bifloris : ca-
lycibus pilosis decem angulatis. *P. Fl. Oec. 76.*

Dan. *Roberts-Urt.*

Pharmac. GERANII ROBERTIANI herba.

In ruderatis & ad muros.

554. † GERANIUM *lucidum* pedunculis bifloris: calycibus pyramidatis angulatis elevato-rugosis: foliis quinquelobis rotundatis. *Fl. Dan. t. 218.*

555. * GERANIUM *molle* pedunculis bifloris foliisque floralibus alternis: caule ramoso erectiusculo: calycibus muticis: arillis lævibus.

Vaill. Bot. Paris. t. 15. f. 3.

In nemore.

556. GERANIUM *dissectum* pedunculis bifloris: foliis quinquepartito-trifidis: petalis emarginatis longitudine calycis: arillis villosis.

Vaill. Bot. Paris. t. 15. f. 2.

Ad sepimenta.

557. * GERANIUM *columbinum* pedunculis bifloris folio longioribus: foliis quinquepartito-multifidis: laciniis acutis: arillis glabris: calycibus aristatis.

Vaill. Bot. Paris. t. 15. f. 4.

In arvis.

558. GERANIUM *retundifolium* pedunculis bifloris: petalis integris obtusissimis longitudine calycis: caule prostrato: foliis reniformibus incis.

Vaill. Bot. Paris. t. 15. f. 1.

In cultis.

* * * Pedunculis *unifloris*.

559. GERANIUM *sanguineum* pedunculis unifloris: foliis quinquepartitis trifidis orbiculatis.

In dumetis ficcis collis *Brankebakke* frequens.

POLYANDRIA.

254. A L T H Æ A. 839.

560. † ALTHÆA *officinalis* foliis simplicibus tomentosis.
Dan. *Althee*.

255. M A L V A. 841.

561. *MALVA rotundifolia* caule prostrato: foliis cordato-orbiculatis, obsolete quinquelobis: pedunculis fructiferis declinatis.

Dan. *Kat - Ost.*

In ruderatis & ad pagos.

562. *MALVA sylvestris* caule erecto herbaceo: foliis septemlobatis acutis: pedunculis petiolisque pilosis.

Pharmac. MALVÆ VULGARIS radix, herba, semina.
Ad vias.

563. † *MALVA Alcea* caule erecto: foliis multipartitis scabriusculis.

Dan. *Cimons - Urt.*



Classis XVII.

DIADELPHIA.

HEXANDRIA.

256. FUMARIA. *Cal.* 2-phyllus. *Cor.* ringens, basi gibbosa, nectarifera. *Filamenta* 2 membranacea, singula antheris 3.

OCTANDRIA.

257. POLYGALA. *Cal.* 5-phyllus: foliolis duobus ætèrormibus, coloratis. *Cor.* vexillum cylindricum. *Stamina* connexa. *Legumen* obcordatum, biloculare.

DECANDRIA.

Stamina omnia connexa.

258. SPARTIUM. *Stigma* longitudinale, supra villosum. *Filamenta* carinæ adhærentia. *Cal.* deorsum productus.
259. GENISTA. *Pistillum* deprimens carinam. *Stigma* involutum. *Cal.* 2-labiatus: $\frac{2}{3}$. *Vexillum* oblongum, a pistillo staminibusque deorsum reflexum.
262. ANTHYLLIS. *Cal.* ventricosus, turgidus, siliquam includens. *Legumen* subrotundum, tectum.
261. ONONIS. *Cal.* 5-partitus: laciniis linearibus. *Vexillum* striatum. *Legumen* turgidum, rhombeum, sessile.

Stigma pubescens, (nec priorum notæ.)

264. OROBUS. *Stylus* linearis, teretiusculus, supra villosus. *Cal.* basi obtusus: *Lacinis* superioribus profundioribus, brevioribus.
263. PISUM. *Stylus* triangulus supra carinatus villosusque. *Calycis* lacinia superiores 2 breviores.
265. LATHYRUS. *Stylus* planus, supra villosus, superne latior. *Calycis* lacinia superiores 2 breviores.
266. VICIA. *Stylus* sub stigmate barbatus.

Legumen subbiloculare (nec priorum.)

270. ASTRAGALUS. *Legumen* biloculare, rotundatum, gibbum.

Legumina submonosperma (nec priorum.)

271. TRIFOLIUM. *Legumen* vix calyce longius, 1 - seu 2 - spermum, non dehiscens, deciduum. *Flores* capitati.

Legumen subarticulatum.

269. HEDYSARUM. *Legumen* articulis subrotundis, compressis, 1 - spermis. *Cor.* carina obtusissima.
260. CORONILLA. *Legumen* isthmis interceptum, rectum. *Cal.* 2-labiatus: $\frac{2}{3}$. dentibus superioribus connatis. *Vexillum* vix alis longius.
268. ORNITHOPUS. *Legumen* articulatum, teres, arcuatum.
273. MEDICAGO. *Legumen* spirale, membranaceo-compressum. *Carina* corollæ a vexillo deflectens.

Legumen uniloculare, polyspermum (nec priorum.)

267. ERVUM. *Calyx* 5-partitus, subæqualis longitudine fere corollæ.
272. LOTUS. *Legumen* cylindricum, strictum, factum feminibus cylindricis. Alæ sursum longitudinaliter conniventes. *Cal.* tubulosus.

HEXANDRIA.

HEXANDRIA.

256. FUMARIA. 849.

564. FUMARIA *bulbosa* caule simplici: bracteis longitudine florum.

- a) Fumaria bulbosa radice cava, major.
- β) Fumaria bulbosa radice non cava, minor.

Morif. Hist. p. 261. t. 12. f. 8.

Dan. *Hvul-Roed; Röd Hane-Kamm.*

Pharmac. ARISTOLOCHIE FABACEÆ radix.

- a) Frequens in nemorosis ad lacum *Wareslöse.*
- β) Frequens in Vivario *Friderichsdalensi.*

565. FUMARIA *officinalis* pericarpis monospermis racemosis: caule diffuso.

Dan. *Jordrög.*

Pharmac. FUMARIÆ herba, semina.

In arvis & fatis frequens.

OCTANDRIA.

257. POLYGALA. 851.

566. POLYGALA *vulgaris* floribus cristatis racemosis: caulibus herbaceis simplicibus procumbentibus: foliis lineari-lanceolatis.

Vaill. Bot. Paris. t. 32. f. 2.

Dan. *Kaars-Blomster.*

In siccis vulgaris.

β) POLYGALA *violacea.*

γ) POLYGALA *carnea.*

δ) POLYGALA *alba.*

ε) POLYGALA *major.* Vaill. Bot. Paris. t. 32. f. 2.

DECANDRIA.

258. SPARTIUM. 88.

567. SPARTIUM *scoparium* foliis ternatis solitariisque:
ramis inermibus angulatis.

In arenosis sylvarum.

259. GENISTA. 859.

* *Inermes.*

568. † GENISTA *tinctoria* foliis lanceolatis glabris:
ramis striatis teretibus erectis.

Pharmac. GENISTÆ herba, flores, semina.

569. † GENISTA *pilosa* foliis lanceolatis obtusis : caule
tuberculato decumbente.

In Ducatibus obvia.

* * *Spinosa.*

570. † GENISTA *Germanica* spinis compositis : ramis
floriferis inermibus : foliis lanceolatis.

In ericetis *Holsatiae* passim.

260. CORONILLA. 883.

571. † CORONILLA *varia* herbacea, leguminibus ere-
ctis teretibus torosis numerosis : foliolis plurimis
glabris.

261. ONONIS. 863.

572. ONONIS *spinosa* floribus racemosis geminis, fo-
liis ternatis : superioribus solitariis : ramis inermi-
bus subvillosis.

Dan. *Krage-Torn.*

Pharmac. ONONIDIS radix, herba.

In campestri frequens.

262. ANTHYLLIS. 864.

73. ANTHYLLIS *vulneraria* herbacea: foliis quaternis-pinnatis: floribus lateralibus.

Dan. *Katte - Klever.*

In sterilibus.

263. PISUM. 870.

74. PISUM *maritimum* petiolis supra planiusculis: caule angulato: stipulis sagittatis: pedunculis multifloris.

Morif. Hist. t. 1. f. 5.

In maritimis.

264. OROBUS. 871.

75. OROBUS *vernus* foliis pinnatis ovatis: stipulis semisagittatis integerrimis: caule simplici.

Dan. *Muus - Erter.*

Pharmac. OROBI farina.

In montosis nemorum.

76. * OROBUS *tuberosus* foliis pinnatis lanceolatis: stipulis semisagittatis integerrimis: caule simplici.

In Sylvis.

77. * OROBUS *niger* caule ramoso: foliis sexjugis ovato-oblongis.

In montosis sylvaticis.

265. LATHYRUS. 872.

* Pedunculis *unifloris*.

78. † LATHYRUS *Aphaca* pedunculis unifloris: circa his aphyllis: stipulis sagittato-cordatis.

Dan. *Vikker.*

* * Pedunculis multifloris.

579. † LATHYRUS *tuberosus* pedunculis multifloris: cirrhis diphyllis: foliolis ovalibus: internodiis nudis. *P. Fl. Oec.* 89.

Dan. *Jord - Rødder*; *Heede - Nødder*.

580. LATHYRUS *pratensis* pedunculis multifloris: cirrhis diphyllis simplicissimis: foliolis lanceolatis.

Dan. *Vilde Erter*.

In dumetis.

581. LATHYRUS *sylvestris* pedunculis multifloris: cirrhis diphyllis: foliolis ensiformibus: internodiis membranaceis.

Extra hortum ac in declivi *Brankebakke*.

582. LATHYRUS *latifolius* pedunculis multifloris: cirrhis diphyllis: foliolis lanceolatis: internodiis membranaceis.

583. * LATHYRUS *palustris* pedunculis multifloris: cirrhis polyphyllis: stipulis lanceolatis.

In pascuis humidis.

266. V I C I A. 873.

* Pedunculis elongatis.

584. VICIA *dumetorum* pedunculis multifloris: foliolis reflexis ovatis mucronatis: stipulis subdentatis.

Dan. *Vikker*.

585. † VICIA *sylvatica* pedunculis multifloris: foliolis ovalibus: stipitulis denticulatis.

586. VICIA *Cracca* pedunculis multifloris: floribus imbricatis: foliolis lanceolatis pubescentibus: stipulis integris.

Dan. *Muus - Erter*.

Inter segetes, ac in nemore versus amnem.

* * Floribus axillaribus subsessilibus.

87. *VICIA sativa* leguminibus sessilibus subbinatis erectis: foliolis retusis: stipulis notatis.

Inter segetes.

88. *VICIA Lathyroides* leguminibus sessilibus solitariis, erectis, glabris: foliolis senis: inferioribus obcordatis. *Fl. Dan. t. 58.*

In aridis montosis, ac in agris passim.

89. *VICIA sepium* leguminibus pedicellatis subquaternis erectis: foliolis ovatis integerrimis: exterioribus decreascentibus.

In umbrosis.

267. ERVUM. 874.

90. *ERVUM tetraspermum* pedunculis subbifloris: seminibus globosis quaternis. *Fl. Dan. t. 95.*

In oleraceis frequens.

91. *ERVUM hirsutum* pedunculis multifloris: seminibus globosis binis.

In arvis, ac in horto frequens.

268. ORNITHOPUS. 884.

92. † *ORNITHOPUS perpusillus* foliis pinnatis: leguminibus subarcuatis.

269. HEDYSARUM. 887.

93. † *HEDYSARUM Onobrychis* foliis pinnatis: leguminibus monospermis aculeatis: corollarum alis calyce brevioribus: caule elongato.

270. ASTRAGALUS. 892.

- §94. ASTRAGALUS *glycyphyllus* caulescens prostratus: leguminibus subtriquetris arcuatis: foliis ovalibus pedunculo longioribus. *Morif. Hist. III. t. 9. f. 1.*
 Dan. *Vild Lakrits.*

In sylvatico *Deyligheden* frequens, & alibi.

- §95. † ASTRAGALUS *arenarius* subcaulescens, procumbens, floribus subracemosis erectis: foliis tomentosis.

Dan. *Bierg - Lakrits.*

271. TRIFOLIUM. 896.

* *Meliloti leguminibus nudis polyspermis.*

- §96. TRIFOLIUM *M. officinalis* leguminibus racemosis nudis, dispermis, rugosis, acutis: caule erecto.
 Dan. *Amur; Steen - Klever.*

Pharmac. MELILOTI herba, flores, semina.

In *Fionia* vulgaris; extra *Haderslebiam* corollis omnibus nitide albis legi.

In campestribus, imprimis inter *Molendina*.

** *Lotoidea leguminibus testis polyspermis.*

- §97. † TRIFOLIUM *hybridum* capitulis umbellaribus: leguminibus tetraspermis: caule adscendente.
 Caulis teres, subpilosus, non sulcatus.

In via ultra cauponam *Kiøge - Kroe* vulgo frequentissimam legi.

- §98. TRIFOLIUM *repens* capitulis umbellaribus: leguminibus tetraspermis: caule repente.

Dan. *Huid - Klever.*

Pharmac. TRIFOLII ALBI flores.

In pascuis.

* * * *Lagopoda calycibus villosis.*

599. *TRIFOLIUM pratense* spicis subvillosis cinctis: stipulis oppositis membranaceis: corollis monopetalis. *P. Fl. Oec.* 146.

Dan. *Sue - Kop*; *Röd - Klever*; *Sukker - Bröd.*

In pratiis ubique.

600. *TRIFOLIUM arvense* spicis villosis ovalibus: dentibus calycinis setaceis æqualibus.

Dan. *Hare - Klever.*

In arvis.

* * * * *Vesicaria calycibus inflatis ventricosis.*

601. † *TRIFOLIUM spumosum* spicis ovatis: calycibus inflatis glabris quinq̃ue dentatis: involucris universalibus pentaphyllis.

602. *TRIFOLIUM fragiferum* capitulis subrotundis: calycibus inflatis bidentatis reflexis: caulibus repentibus. *Vaill. Bot. Paris.* t. 22. f. 2.

* * * * * *Lupulina vexillis corollæ inflexis.*

603. † *TRIFOLIUM montanum* spicis subimbricatis subtribus: vexillis subulatis emarcescentibus: calycibus nudis: caule erecto.

604. † *TRIFOLIUM agrarium* spicis ovalibus imbricatis: vexillis deflexis persistentibus: calycibus nudis: caule erecto.

Dan. *Humle - Klever.*

272 **LOTUS** 897.

* *Leguminibus prioribus nec capitulum conspicuis.*

605. † *LOTUS maritimus* leguminibus solitariis membranaceo - quadrangulatis: foliis glabris bracteis lanceolatis.

Dan. *Steen - Klever.*

* * Pedunculis *multifloris in capitulum.*

606. LOTUS *corniculatus* capitulis depresso: caulibus decumbentibus; leguminibus cylindricis patentibus. *P. Fl. Oec. 95.*

Dan. *Kierlinge - Tander; Krage - Kloer.*

In pratis ac collibus.

273. M E D I C A G O. 899.

607. MEDICAGO *falcata* pedunculis racemosis: leguminibus lunatis: caule prostrato.

Fl. Dan. t. 233. P. Fl. Oec. 97.

Dan. *Gnul - Klover.*

In Icone legumina defunt, caracter plantæ specificus.

In agris ad vias.

608. † MEDICAGO *lupulina* spicis ovalibus: leguminibus reniformibus monospermis: caule procumbente.

In pratis & ad vias.

609. † MEDICAGO *polymorpha* minima leguminibus cochleatis aculeis uncinatis alternis externe domatis: stipulis integris. *Fl. Dan. t. 211.*



Classis XVIII.

POLYADELPHIA.

POLYANDRIA.

274. HYPERICUM. Cal. 5 - partitus. Cor. 5 - petala.
Filamenta multa in 5 phalangibus. Styli 1, 3, 5.
Caps. locularis.



255. MALVA. 841.

561. MALVA *rotundifolia* caule prostrato: foliis cordato-orbiculatis, obsolete quinquelobis: pedunculis fructiferis declinatis.

Dan. *Kat - Ost.*

In ruderatis & ad pagos.

562. MALVA *sylvestris* caule erecto herbaceo: foliis septemlobatis acutis: pedunculis petiolisque pilosis.

Pharmac. MALVÆ VULGARIS radix, herba, semina.
Ad vias.

563. † MALVA *Alcea* caule erecto: foliis multipartitis scabriusculis.

Dan. *Cimons - Urt.*



Classis XVII.

DIADELPHIA.

HEXANDRIA.

256. FUMARIA. *Cal.* 2-phyllus. *Cor.* ringens, basi gibbosa, nectarifera. *Filamenta* 2 membranacea, singula antheris 3.

OCTANDRIA.

257. POLYGALA. *Cal.* 5-phyllus: foliolis duobus ætèrormibus, coloratis. *Cor.* vexillum cylindricum. *Stamina* connexa. *Legumen* obcordatum, biloculare.

DECANDRIA.

Stamina omnia connexa.

258. SPARTIUM. *Stigma* longitudinale, supra villosum. *Filamenta* carinæ adhærentia. *Cal.* deorsum productus.
259. GENISTA. *Pistillum* deprimens carinam. *Stigma* involutum. *Cal.* 2-labiatus: $\frac{2}{3}$. *Vexillum* oblongum, a pistillo staminibusque deorsum reflexum.
262. ANTHYLLIS. *Cal.* ventricosus, turgidus, filiquam includens. *Legumen* subrotundum, tectum.
261. ONONIS. *Cal.* 5-partitus: laciniis linearibus. *Vexillum* striatum. *Legumen* turgidum, rhombeum, setifere.

255. MALVA. 841.

561. MALVA *rotundifolia* caule prostrato: foliis cordato-orbiculatis, obsolete quinquelobis: pedunculis fructiferis declinatis.

Dan. *Kat - Ost.*

In ruderatis & ad pagos.

562. MALVA *sylvestris* caule erecto herbaceo: foliis septemlobatis acutis: pedunculis petiolisque pilosis.

Pharmac. MALVÆ VULGARIS radix, herba, semina.
Ad vias.

563. † MALVA *Alcea* caule erecto: foliis multipartitis scabriusculis.

Dan. *Cimons - Urt.*



Classis XVII.

DIADELPHIA.

HEXANDRIA.

256. FUMARIA. *Cal.* 2-phyllus. *Cor.* ringens, basi gibbosa, nectarifera. *Filamenta* 2 membranacea, singula antheris 3.

OCTANDRIA.

257. POLYGALA. *Cal.* 5-phyllus: foliolis duobus æqualioribus, coloratis. *Cor.* vexillum cylindricum. *Stamina* connexa. *Legumen* obcordatum, biloculare.

DECANDRIA.

Stamina omnia connexa.

258. SPARTIUM. *Stigma* longitudinale, supra villosum. *Filamenta* carinæ adhærentia. *Cal.* deorsum productus.
259. GENISTA. *Pistillum* deprimens carinam. *Stigma* involutum. *Cal.* 2-labiatus: $\frac{2}{3}$. *Vexillum* oblongum, a pistillo staminibusque deorsum reflexum.
262. ANTHYLLIS. *Cal.* ventricosus, turgidus, filiquam includens. *Legumen* subrotundum, tectum.
261. ONONIS. *Cal.* 5-partitus: laciniis linearibus. *Vexillum* striatum. *Legumen* turgidum, rhombeum, fetile.

Stigma pubescens, (nec priorum notæ.)

264. OROBUS. *Stylus* linearis, teretiusculus, supra villosus. *Cal.* basi obtusus: *Lacinii* superioribus profundioribus, brevioribus.
265. PISUM. *Stylus* triangulus supra carinatus villosusque. *Calycis* lacinie superiores 2 breviores.
265. LATHYRUS. *Stylus* planus, supra villosus, superne latior. *Calycis* lacinie superiores 2 breviores.
266. VICIA. *Stylus* sub stigmate barbatus.

Legumen subbiloculare (nec priorum.)

270. ASTRAGALUS. *Legumen* biloculare, rotundatum, gibbum.

Legumina submonosperma (nec priorum.)

271. TRIFOLIUM. *Legumen* vix calyce longius, 1 - seu 2 - spermum, non dehiscens, deciduum. *Flores* capitati.

Legumen subarticulatum.

269. HEDYSARUM. *Legumen* articulis subrotundis, compressis, 1 - spermis. *Cor.* carina obtusissima.
260. CORONILLA. *Legumen* isthmis interceptum, rectum. *Cal.* 2-labiatus: $\frac{2}{3}$. dentibus superioribus connatis. *Vexillum* vix alis longius.
268. ORNITHOPUS. *Legumen* articulatum, teres, arcuatum.
273. MEDICAGO. *Legumen* spirale, membranaceo-compressum. *Carina* corollæ a vexillo deflectens.

Legumen uniloculare, polyspermum (nec priorum.)

267. ERVUM. *Calyx* 5-partitus, subæqualis longitudine fere corollæ.
272. LOTUS. *Legumen* cylindricum, strictum, factum seminibus cylindricis. Alæ sursum longitudinaliter conniventes. *Cal.* tubulosus.

HEXANDRIA.

HEXANDRIA.

256. FUMARIA. 849.

564. *FUMARIA bulbosa* caule simplici: bracteis longitudine florum.

a) *Fumaria bulbosa* radice cava, major.

β) *Fumaria bulbosa* radice non cava, minor.

Morif. Hist. p. 261. t. 12. f. 8.

Dan. *Huul-Roed; Röd Hane-Kamm.*

Pharmac. ARISTOLOCHIE FABACEÆ radix.

a) Frequens in nemorosis ad lacum *Warelöse.*

β) Frequens in Vivario *Friderichsdalensi.*

565. *FUMARIA officinalis* pericarpis monospermis racemosis: caule diffuso.

Dan. *Jordrög.*

Pharmac. FUMARIÆ herba, semina.

In arvis & fatis frequens.

OCTANDRIA.

257. POLYGALA. 851.

566. *POLYGALA vulgaris* floribus cristatis racemosis: caulibus herbaceis simplicibus procumbentibus: foliis lineari-lanceolatis.

Vaill. Bot. Paris. t. 32. f. 2.

Dan. *Kaars-Blomster.*

In siccis vulgaris.

β) *POLYGALA violacea.*

γ) *POLYGALA carnea.*

δ) *POLYGALA alba.*

ε) *POLYGALA major. Vaill. Bot. Paris. t. 32. f. 2.*

DECANDRIA.

258. SPARTIUM. 88.

567. SPARTIUM *scoparium* foliis ternatis solitariisque:
ramis inermibus angulatis.

In arenosis fylvarum.

259. GENISTA. 859.

* *Inermes.*

568. † GENISTA *tinctoria* foliis lanceolatis glabris:
ramis striatis teretibus erectis.

Pharmac. GENISTÆ herba, flores, femina.

569. † GENISTA *pilosa* foliis lanceolatis obtusis : caule
tuberculato decumbente.

In Ducatibus obvia.

* * *Spinosa.*

570. † GENISTA *Germanica* spinis compositis : ramis
floriferis inermibus : foliis lanceolatis.

In ericetis *Holsatie* passim.

260. CORONILLA. 883.

571. † CORONILLA *varia* herbacea, leguminibus ere-
ctis teretibus torosis numerosis : foliolis plurimis
glabris.

261. ONONIS. 863.

572. ONONIS *spinosa* floribus racemosis geminis, fo-
liis ternatis : superioribus solitariis : ramis inermi-
bus subvillosis.

Dan. *Krage-Torn.*

Pharmac. ONONIDIS radix, herba.

In campestri frequens.

262. ANTHYLLIS. 864.

73. ANTHYLLIS *vulneraria* herbacea: foliis quaternio-pinnatis: floribus lateralibus.

Dan. *Katte - Klever.*

In sterilibus.

263. PISUM. 870.

74. PISUM *maritimum* petiolis supra planiusculis: caule angulato: stipulis sagittatis: pedunculis multifloris.

Morif. Hist. t. I. f. 5.

In maritimis.

264. OROBUS. 871.

75. OROBUS *vernus* foliis pinnatis ovatis: stipulis semisagittatis integerrimis: caule simplici.

Dan. *Muus - Erter.*

Pharmac. OROBI farina.

In montosis nemorum.

76. * OROBUS *tuberosus* foliis pinnatis lanceolatis: stipulis semisagittatis integerrimis: caule simplici.

In Sylvis.

77. * OROBUS *niger* caule ramoso: foliis sexjugis ovato-oblongis.

In montosis fylvaticis.

265. LATHYRUS. 872.

* Pedunculis *unifloris*.

78. † LATHYRUS *Aphaca* pedunculis unifloris: circumsistis aphyllis: stipulis sagittato-cordatis.

Dan. *Vikker.*

* * Pedunculis multifloris.

579. † LATHYRUS *tuberosus* pedunculis multifloris: cirrhis diphyllis: foliolis ovalibus: internodiis nudis. *P. Fl. Oec.* 89.

Dan. *Jord - Rødder*; *Heede - Nødder*.

580. LATHYRUS *pratensis* pedunculis multifloris: cirrhis diphyllis simplicissimis: foliolis lanceolatis.

Dan. *Vilde Erter*.

In dumetis.

581. LATHYRUS *sylvestris* pedunculis multifloris: cirrhis diphyllis: foliolis ensiformibus: internodiis membranaceis.

Extra hortum ac in declivi *Brankebakke*.

582. LATHYRUS *latifolius* pedunculis multifloris: cirrhis diphyllis: foliolis lanceolatis: internodiis membranaceis.

583. * LATHYRUS *palustris* pedunculis multifloris: cirrhis polyphyllis: stipulis lanceolatis.

In pascuis humidis.

266. V I C I A. 873.

* Pedunculis elongatis.

584. VICIA *dumetorum* pedunculis multifloris: foliolis reflexis ovatis mucronatis: stipulis subdentatis.

Dan. *Vikker*.

585. † VICIA *sylvatica* pedunculis multifloris: foliolis ovalibus: stipitulis denticulatis.

586. VICIA *Cracca* pedunculis multifloris: floribus imbricatis: foliolis lanceolatis pubescentibus: stipulis integris.

Dan. *Muus - Erter*.

Inter segetes, ac in nemore versus amnem.

* * Floribus axillaribus subsessilibus.

87. *VICIA sativa* leguminibus sessilibus subbinatis erectis : foliolis retusis : stipulis notatis.

Inter segetes.

88. *VICIA Lathyroides* leguminibus sessilibus solitariis, erectis, glabris : foliolis fenis : inferioribus obcordatis. *Fl. Dan. t. 58.*

In aridis montosis, ac in agris passim.

89. *VICIA sepium* leguminibus pedicellatis subquaternis erectis : foliolis ovatis integerrimis : exterioribus deoresentibus.

In umbrosis.

267. ERVUM. 874.

90. *ERVUM tetraspermum* pedunculis subbifloris : seminibus globosis quaternis. *Fl. Dan. t. 95.*

In oleraceis frequens.

91. *ERVUM hirsutum* pedunculis multifloris : seminibus globosis binis.

In arvis, ac in horto frequens.

268. ORNITHOPUS. 884.

92. † *ORNITHOPUS perpusillus* foliis pinnatis : leguminibus subarcuatis.

269. HEDYSARUM. 887.

93. † *HEDYSARUM Onobrychis* foliis pinnatis : leguminibus monospermis aculeatis : corollarum alis calyce brevioribus : caule elongato.

270. ASTRAGALUS. 892.

594. ASTRAGALUS *glycyphyllus* caulescens prostratus: leguminibus subtriquetris arcuatis: foliis ovalibus pedunculo longioribus. *Morif. Hist. III. t. 9. f. 8.*
 Dan. *Vild Lakrits.*

In sylvatico *Deyligheden* frequens, & alibi.

595. † ASTRAGALUS *arenarius* subcaulescens, procumbens, floribus subracemosis erectis: foliis tomentosis.

Dan. *Bierg - Lakrits.*

271. TRIFOLIUM. 896.

* *Meliloti leguminibus nudis polyspermis.*

596. TRIFOLIUM *M. officinalis* leguminibus racemosis nudis, dispermis, rugosis, acutis: caule erecto.

Dan. *Amur; Steen - Klever.*

Pharmac. MELILOTI herba, flores, semina.

In *Fionia* vulgaris; extra *Haderslebiam* corollis omnibus nitide albis legi.

In campestribus, imprimis inter *Molendina*.

** *Lotoidea leguminibus tectis polyspermis.*

597. † TRIFOLIUM *hybridum* capitulis umbellaribus: leguminibus tetraspermis: caule ascendente.

Caulis teres, subpilosus, non sulcatus.

In via ultra cauponam *Kiøge - Kroe* vulgo frequentissimam legi.

598. TRIFOLIUM *repens* capitulis umbellaribus: leguminibus tetraspermis: caule repente.

Dan. *Huid - Klever.*

Pharmac. TRIFOLII ALBI flores.

In pascuis.

* * * *Lagopoda calycibus villosis.*

599. *TRIFOLIUM pratense* spicis subvillosis cinctis: stipulis oppositis membranaceis: corollis monopetalis. *P. Fl. Oec.* 146.

Dan. *Sue - Kop; Röd - Klever; Sukker - Bröd.*

In pratiis ubique.

600. *TRIFOLIUM arvense* spicis villosis ovalibus: dentibus calycinis setaceis æqualibus.

Dan. *Hare - Klever.*

In arvis.

* * * * *Vesicaria calycibus inflatis ventricosis.*

601. † *TRIFOLIUM spumosum* spicis ovatis: calycibus inflatis glabris quinque dentatis: involucris universalibus pentaphyllis.

602. *TRIFOLIUM fragiferum* capitulis subrotundis: calycibus inflatis bidentatis reflexis: caubus repentibus. *Vaill. Bot. Paris. t. 22. f. 2.*

* * * * * *Lupulina vexillis corollæ inflexis.*

603. † *TRIFOLIUM montanum* spicis subimbricatis subtribus: vexillis subulatis emarcescentibus: calycibus nudis: caule erecto.

604. † *TRIFOLIUM agrarium* spicis ovalibus imbricatis: vexillis deflexis persistentibus: calycibus nudis: caule erecto.

Dan. *Humble - Klever.*

2725 **LOTUS** 897.

* Leguminibus parvis nec capitulum
constituentibus.

605. † *LOTUS maritimus* leguminibus solitariis membranaceo-quadrangulatis: foliis glabris bracteis lanceolatis.

Dan. *Steen - Klever.*

* * Pedunculis *multifloris in capitulum*.

606. *LOTUS corniculatus* capitulis depressis: caulibus decumbentibus: leguminibus cylindricis patentibus. *P. Fl. Oec. 95.*

Dan. *Kierlinge - Tender; Krage - Klier.*
In pratis ac collibus.

273. M E D I C A G O. 899.

607. *MEDICAGO falcata* pedunculis racemosis: leguminibus lunatis: caule prostrato.

Fl. Dan. t. 233. P. Fl. Oec. 97.

Dan. *Gnul - Klee.*

In Icone legumina defunt, caracter plantæ specificus.

In agris ad vias.

608. † *MEDICAGO lupulina* spicis ovalibus: leguminibus reniformibus monospermis: caule procumbente.

In pratis & ad vias.

609. † *MEDICAGO polymorpha* minima leguminibus cochleatis aculeis uncinatis alternis externe domatis stipulis integris. *Fl. Dan. t. 211.*



Classis XVIII.

POLYADELPHIA.

POLYANDRIA.

- 274.** HYPERICUM. *Cal.* 5 - partitus. *Cor.* 5 - petala.
Filamenta multa in 5 phalangibus. *Styli* 1, 3, 5.
Caps. locularis.



Stigma pubescens, (nec priorum notæ.)

264. OROBUS. *Stylus* linearis, teretiusculus, supra villosus. *Cal.* basi obtusus: *Lacinii* superioribus profundioribus, brevioribus.
265. PISUM. *Stylus* triangulus supra carinatus villosusque. *Calycis* laciniae superiores 2 breviores.
266. LATHYRUS. *Stylus* planus, supra villosus, superne latior. *Calycis* laciniae superiores 2 breviores.
266. VICIA. *Stylus* sub stigmate barbatus.

Legumen subbiloculare (nec priorum.)

270. ASTRAGALUS. *Legumen* biloculare, rotundatum, gibbum.

Legumina submonosperma (nec priorum.)

271. TRIFOLIUM. *Legumen* vix calyce longius, 1 - seu 2 - spermum, non dehiscens, deciduum. *Flores* capitati.

Legumen subarticulatum.

269. HEDYSARUM. *Legumen* articulis subrotundis, compressis, 1 - spermis. *Cor.* carina obtusissima.
260. CORONILLA. *Legumen* isthmis interceptum, rectum. *Cal.* 2-labiatus: $\frac{2}{3}$. dentibus superioribus connatis. *Vexillum* vix alis longius.
268. ORNITHOPUS. *Legumen* articulatum, teres, arcuatum.
273. MEDICAGO. *Legumen* spirale, membranaceo-compressum. *Carina* corollae a vexillo deflectens.

Legumen uniloculare, polyspermum (nec priorum.)

267. ERVUM. *Calyx* 5-partitus, subaequalis longitudine fere corollae.
272. LOTUS. *Legumen* cylindricum, strictum, factum seminibus cylindricis. Alae sursum longitudinaliter conniventes. *Cal.* tubulosus.

HEXANDRIA.

HEXANDRIA.

256. FUMARIA. 849.

564. FUMARIA *bulbosa* caule simplici: bracteis longitudine florum.

α) Fumaria bulbosa radice cava, major.

β) Fumaria bulbosa radice non cava, minor.

Morif. Hist. p. 261, t. 12, f. 8.

Dan. *Hul-Roed; Röd Hane-Kumm.*

Pharmac. ARISTOLOCHIE FABACEE radix.

α) Frequens in nemorosis ad lacum *Wæreløse*.

β) Frequens in Vivario *Friderichsdalensi*.

565. FUMARIA *officinalis* pericarpis monospermis racemosis: caule diffuso.

Dan. *Jordrög.*

Pharmac. FUMARIÆ herba, semina.

In arvis & satis frequens.

OCTANDRIA.

257. POLYGALA. 851.

566. POLYGALA *vulgaris* floribus cristatis racemosis: caulibus herbaceis simplicibus procumbentibus: foliis lineari-lanceolatis.

Vaill. Bot. Paris. t. 32, f. 2.

Dan. *Kaars-Blomster.*

In ficcis vulgaris.

β) POLYGALA *violacea*.

γ) POLYGALA *carnea*.

δ) POLYGALA *alba*.

ε) POLYGALA *major*. *Vaill. Bot. Paris. t. 32, f. 2.*

DECANDRIA.

258. SPARTIUM. 88.

567. SPARTIUM *scoparium* foliis ternatis solitariisque:
ramis inermibus angulatis.

In arenosis fylvarum.

259. GENISTA. 859.

* *Inermes.*

568. † GENISTA *tinctoria* foliis lanceolatis glabris:
ramis striatis teretibus erectis.

Pharmac. GENISTÆ herba, flores, femina.

569. † GENISTA *pilosa* foliis lanceolatis obtusis : caule
tuberculato decumbente.

In Ducatibus obvia.

* * *Spinosa.*

570. † GENISTA *Germanica* spinis compositis : ramis
floriferis inermibus : foliis lanceolatis.

In ericetis *Holsatiæ* passim.

260. CORONILLA. 883.

571. † CORONILLA *varia* herbacea, leguminibus ere-
ctis teretibus torosis numerosis : foliolis plurimis
glabris.

261. ONONIS. 863.

572. ONONIS *spinosa* floribus racemosis geminis, fo-
liis ternatis : superioribus solitariis : ramis inermi-
bus subvillosis.

Dan. *Krage-Torn.*

Pharmac. ONONIDIS radix, herba.

In campestri frequens.

262. ANTHYLLIS. 864.

73. ANTHYLLIS *vulneraria* herbacea: foliis quaternis: pinnatis: floribus lateralibus.

Dan. *Katte - Klever.*

In sterilibus.

263. PISUM. 870.

74. PISUM *maritimum* petiolis supra planiusculis: caule angulato: stipulis sagittatis: pedunculis multifloris.

Morif. Hist. t. 1. f. 5.

In maritimis.

264. OROBUS. 871.

75. OROBUS *vernus* foliis pinnatis ovatis: stipulis semisagittatis integerrimis: caule simplici.

Dan. *Muus - Erter.*

Pharmac. OROBI farina.

In montosis nemorum.

76. * OROBUS *tuberosus* foliis pinnatis lanceolatis: stipulis semisagittatis integerrimis: caule simplici.

In Sylvis.

77. * OROBUS *niger* caule ramoso: foliis sexjugis ovato-oblongis.

In montosis sylvaticis.

265. LATHYRUS. 872.

* Pedunculis *unifloris.*

78. † LATHYRUS *Aphaca* pedunculis unifloris: circa his aphyllis: stipulis sagittato-cordatis.

Dan. *Vikker.*

* * Pedunculis multifloris.

579. † LATHYRUS *tuberosus* pedunculis multifloris: cirrhis diphyllis: foliolis ovalibus: internodiis nudis. *P. Fl. Oec.* 89.

Dan. *Jord-Rödder*; *Heede-Nödder*.

580. LATHYRUS *pratensis* pedunculis multifloris: cirrhis diphyllis simplicissimis: foliolis lanceolatis.

Dan. *Vilde Erter*.

In dumetis.

581. LATHYRUS *sylvestris* pedunculis multifloris: cirrhis diphyllis: foliolis ensiformibus: internodiis membranaceis.

Extra hortum ac in declivi *Brankebakke*.

582. LATHYRUS *latifolius* pedunculis multifloris: cirrhis diphyllis: foliolis lanceolatis: internodiis membranaceis.

583. * LATHYRUS *palustris* pedunculis multifloris: cirrhis polyphyllis: stipulis lanceolatis.

In pascuis humidis.

266. V I C I A. 873.

* Pedunculis elongatis.

584. VICIA *dumetorum* pedunculis multifloris: foliolis reflexis ovatis mucronatis: stipulis subdentatis.

Dan. *Vikker*.

585. † VICIA *sylvatica* pedunculis multifloris: foliolis ovalibus: stipitulis denticulatis.

586. VICIA *Cracca* pedunculis multifloris: floribus imbricatis: foliolis lanceolatis pubescentibus: stipulis integris.

Dan. *Muus-Erter*.

Inter segetes, ac in nemore versus amnem.

* * Floribus axillaribus subsessilibus.

87. *VICIA sativa* leguminibus sessilibus subbinatis erectis: foliolis retusis: stipulis notatis.

Inter segetes.

88. *VICIA Lathyroides* leguminibus sessilibus solitariis, erectis, glabris: foliolis senis: inferioribus obcordatis. *Fl. Dan. t. 58.*

In aridis montosis, ac in agris passim.

89. *VICIA sepium* leguminibus pedicellatis subquaternis erectis: foliolis ovatis integerrimis: exterioribus decescentibus.

In umbrosis.

267. ERVUM. 874.

90. *ERVUM tetraspermum* pedunculis subbifloris: seminibus globosis quaternis. *Fl. Dan. t. 95.*

In oleraceis frequens.

91. *ERVUM hirsutum* pedunculis multifloris: seminibus globosis binis.

In arvis, ac in horto frequens.

268. ORNITHOPUS. 884.

92. † *ORNITHOPUS perpusillus* foliis pinnatis: leguminibus subarcuatis.

269. HEDYSARUM. 887.

93. † *HEDYSARUM Onobrychis* foliis pinnatis: leguminibus monospermis aculeatis: corollarum alis calyce brevioribus: caule elongato.

270. ASTRAGALUS. 892.

- §94. ASTRAGALUS *glycyphyllus* caulescens prostratus: leguminibus subtriquetris arcuatis: foliis ovalibus pedunculo longioribus. *Morif. Hist. III. t. 9. f. 8.*

Dan. *Vild Lakrits.*

In sylvatico *Deyligheden* frequens, & alibi.

- §95. † ASTRAGALUS *arenarius* subcaulescens, procumbens, floribus subracemosis erectis: foliis tomentosis.

Dan. *Bierg - Lakrits.*

271. TRIFOLIUM. 896.

* *Meliloti leguminibus nudis polyspermis.*

- §96. TRIFOLIUM *M. officinalis* leguminibus racemosis nudis, dispermis, rugosis, acutis: caule erecto.

Dan. *Amur; Steen - Klever.*

Pharmac. MELILOTI herba, flores, semina.

In *Fionia* vulgaris; extra *Haderslebiam* corollis omnibus nitide albis legi.

In campestribus, imprimis inter *Molendina*.

** *Lotoidea leguminibus testis polyspermis.*

- §97. † TRIFOLIUM *hybridum* capitulis umbellaribus: leguminibus tetraspermis: caule adscendente.

Caulis teres, subpilosus, non sulcatus.

In via ultra cauponam *Kiøge - Kroe* vulgo frequentissimam legi.

- §98. TRIFOLIUM *repens* capitulis umbellaribus: leguminibus tetraspermis: caule repente.

Dan. *Huid - Klever.*

Pharmac. TRIFOLII ALBI flores.

In pascuis.

* * * *Lagopoda calycibus villosis.*

99. *TRIFOLIUM pratense* spicis subvillosis cinctis: stipulis oppositis membranaceis: corollis monope-
talis. *P. Fl. Oec.* 146.

Dan. *Sue - Kop; Rød - Klever; Sukker - Brød.*

In pratis ubique.

600. *TRIFOLIUM arvense* spicis villosis ovalibus: den-
tibus calycinis fetaceis æqualibus.

Dan. *Hare - Klever.*

In arvis.

* * * * *Vesicaria calycibus inflatis ventricosis.*

601. † *TRIFOLIUM spumosum* spicis ovatis: calycibus
inflatis glabris quinque dentatis: involucris uni-
versalibus pentaphyllis.

602. *TRIFOLIUM fragiferum* capitulis subrotundis:
calycibus inflatis bidentatis reflexis: caulibus re-
pentibus. *Vaill. Bot. Paris. t. 22. f. 2.*

* * * * * *Lupulina vexillis corollæ inflexis.*

603. † *TRIFOLIUM montanum* spicis subimbricatis
subtribus: vexillis subulatis emarcescentibus: ca-
lycibus nudis: caule erecto.

604. † *TRIFOLIUM agrarium* spicis ovalibus imbricatis: vexillis deflexis persistentibus: calycibus
nudis: caule erecto.

Dan. *Humble - Klever.*

272. LOTUS. — 897.

* Leguminibus rarioribus, nec capitulum
constituentibus.

605. † *LOTUS maritimus* leguminibus solitariis mem-
branaceo-quadrangulatis: foliis glabris bracteis
lanceolatis.

Dan. *Steen - Klever.*

* * Pedunculis *multifloris in capitulum*.

606. *LOTUS corniculatus* capitulis depressis: caulibus decumbentibus: leguminibus cylindricis patentibus. *P. Fl. Oec. 95.*

Dan. *Kierlinge - Tander; Krage - Kloer.*

In pratis ac collibus.

273. M E D I C A G O. 899.

607. *MEDICAGO falcata* pedunculis racemosis: leguminibus lunatis: caule prostrato.

Fl. Dan. t. 233. P. Fl. Oec. 97.

Dan. *Gnul - Klover.*

In Icone legumina defunt, caracter plantæ specificus.

In agris ad vias.

608. † *MEDICAGO lupulina* spicis ovalibus: leguminibus reniformibus monospermis: caule procumbente.

In pratis & ad vias.

609. † *MEDICAGO polymorpha* minima leguminibus cochleatis aculeis uncinatis alternis externe donatis: stipulis integris. *Fl. Dan. t. 211.*



Classis XVIII.

POLYADELPHIA.

POLYANDRIA.

274. HYPERICUM. *Cal.* 5 - partitus. *Cor.* 5 - petala.
Filamenta multa in 5 phalangibus. *Styli* 1, 3, 5.
Caps. locularis.



* * Pedunculis multifloris in capitulum.

606. LOTUS *corniculatus* capitulis depresso: caulibus decumbentibus; leguminibus cylindricis patentibus. *P. Fl. Oec.* 95.

Dan. *Kierlinge - Tander; Krage - Kloer.*

In pratis ac collibus.

273. M E D I C A G O. 899.

607. MEDICAGO *falcata* pedunculis racemosis: leguminibus lunatis: caule prostrato.

Fl. Dan. t. 233. *P. Fl. Oec.* 97.

Dan. *Gnul - Klever.*

In Icone legumina desunt, caracter plantæ specificus.

In agris ad vias.

608. † MEDICAGO *lupulina* spicis ovalibus: leguminibus reniformibus monospermis: caule procumbente.

In pratis & ad vias.

609. † MEDICAGO *polymorpha* minima leguminibus cochleatis aculeis uncinatis alternis externe donatis: stipulis integris. *Fl. Dan. t.* 211.



Classis XVIII.

POLYADELPHIA.

POLYANDRIA.

274. HYPERICUM. *Cal.* 5 - partitus. *Cor.* 5 - petala.
Filamenta multa in 5 phalangibus. *Styli* 1, 3, 1. 5.
Caps. locularis.



POLYANDRIA.

274. HYPERICUM M. 902.

610. *HYPERICUM quadrangulum* floribus trigynis: caule quadrato herbaceo.

Dan. *Piricum*.

In pratis.

611. *HYPERICUM perforatum* floribus trigynis: caule ancipiti, foliis obtusis pellucido-punctatis.

P. Fl. Oec. 85.

Dan. *Hypericon*; *Jord-Humle*; *Blod-Blomster*.

Pharmac. HYPERICI herba, flores, semina.

612. *HYPERICUM humifusum* floribus trigynis axillaribus solitariis: caulibus ancipitibus prostratis filiformibus: foliis glabris. *Fl. Dan.* t. 141.

In arvis haud infrequens.

613. *HYPERICUM hirsutum* floribus trigynis: calycibus ferrato-glandulosis: caule tereti erecto: foliis ovatis subpubescentibus.

Morif. Hist. III. t. 6. f. 11.

In montosis minus frequens.

614. * *HYPERICUM montanum* floribus trigynis: calycibus ferraturis glandulosis: caule tereti erecto glabro: foliis ovatis. *Fl. Dan.* t. 173.

Passim: ad molendina haud infrequens.

615. † *HYPERICUM pulchrum* floribus trigynis: calycibus ferrato-glandulosis: caule tereti: foliis perfoliatis, cordatis, glabris. *Fl. Dan.* t. 75.

In insula *Thorsinge* legi.



Classis XIX.

SYNGENESIA.

POLYGAMIA ÆQUALIS.

Semiflosculosi: Corollis ligulatis omnibus.

- 287. CICHORIUM. *Receptaculum* paleaceum. *Pappus* 5-dentatus. *Calyx* calyculatus.
- 285. HYPOCHÆRIS. *Recept.* paleaceum. *Pappus* subplumosus. *Calyx* subimbricatus.
- 275. TRAGOPOGON. *Recept.* nudum. *Pappus* plumosus. *Cal.* simplex polyphyllus.
- 277. PICRIS. *Recept.* nudum. *Pappus* plumosus. *Cal.* calyculatus. *Semina* transversim fulcata.
- 281. LEONTODON. *Recept.* nudum. *Pappus* plumosus. *Cal.* imbricatus. *Squamis* laxis.
- 278. SONCHUS. *Recept.* nudum. *Pappus* plumosus. *Cal.* imbricatus, basi gibbus.
- 276. SCORZONERA. *Recept.* nudum. *Pappus* plumosus. *Cal.* imbricatus. *Squamis* margine membranaceis.
- 283. CREPIS. *Recept.* nudum. *Pappus* pilosus, stipitatus. *Cal.* calyculatus. *Squamis* deciduis.
- 280. PRENANTHES. *Recept.* nudum. *Pappus* pilosus, simplex, sessilis. *Cal.* calyculatus sub-5-florus.
- 279. LACTUCA. *Recept.* nudum. *Pappus* pilosus, simplex, stipitatus. *Cal.* imbricatus, cylindricus.
- 282. HIERACIUM. *Recept.* nudum. *Pappus* pilosus, simplex, sessilis. *Cal.* imbricatus, ovatus.
- 286. LAPSANA. *Recept.* nudum. *Pappus* nonnullus nullus. *Cal.* calyculatus. *Squamis* singulis interioribus canaliculatis.



284. HYOSERIS. *Recept.* nudum. *Pappus* non pilosus vel obsoletus. *Cal.* subæqualis.

Capitati.

293. CARLINA. *Calyx* radiatus. *Squamis* marginalibus longis, coloratis.
291. CNICUS. *Cal.* ovatus, imbricatus. *Squamis* ramoso-spinosis, obvallatus bracteis. *Corollule* æquales.
288. ARCTIUM. *Cal.* globosus. *Squamis* apice hamis inflexis.
290. CARDUUS. *Cal.* ovatus. *Squamis* ventricosus spinosis. *Recept.* pilosum.
292. ONOPORDON. *Cal.* *Squamis* ventricosus spinosis. *Recept.* nudum.
289. SERRATULA. *Cal.* subcylindricus, subuniflorus, imbricatus, muticus.

Discoidei.

295. EUPATORIUM. *Recept.* nudum. *Pappus* plumosus. *Cal.* imbricatus oblongus. *Stylus* semibifidus, longus.
294. BIDENS. *Recept.* paleaceum. *Pappus* aristis erectis, scabris. *Cal.* imbricatus. *Cor.* rarius flosculo uno alterno radiante instruitur.

POLYGAMIA SUPERFLUA.

Discoidei.

297. ARTEMISIA. *Recept.* subnudum vel subvillosum. *Pappus* nullus. *Cor.* radii nulli. *Cal.* imbricatus. *Squamis* rotundatis conniventibus.

296. TANACETUM. *Recept.* nudum. *Pappus* submarginatus. *Cor.* radii obsoletæ, 3 - fidæ.
299. CONYZA. *Recept.* nudum. *Pappus* pilosus. *Cor.* radii 3 - fidæ.
298. GNAPHALIUM. *Recept.* nudum. *Pappus* plumosus. *Cal.* imbricatus. *Squamis* marginalibus rotundatis, membranaceis, coloratis.

Radiati.

308. BELLIS. *Recept.* nudum, conicum. *Pappus* nullus. *Cal.* hemisphæricus: *Squamis* æqualibus. *Sem.* obovata.
310. MATRICARIA. *Recept.* nudum. *Pappus* nullus. *Cal.* hemisphæricus, imbricatus. *Squamis* marginalibus solidis, acutiusculis.
309. CHRYSANTHEMUM. *Recept.* nudum. *Pappus* nullus. *Cal.* hemisphæricus, imbricatus: *Squamis* marginalibus membranaceis.
307. ARNICA. *Recept.* nudum. *Pappus* simplex, pilosus. *Corollula* radii filamentis 5 absque antheris.
306. INULA. *Recept.* nudum. *Pappus* simplex, pilosus. *Anthera* basi in setas duas desinentes.
300. ERIGERON. *Recept.* nudum. *Pappus* pilosus. *Cor.* radii lineares, angustissimi.
304. SOLIDAGO. *Recept.* nudum. *Pappus* pilosus, simplex. *Cor.* radii circiter 3. *Cal.* Squamæ imbricatæ, clausæ.
305. CINERARIA. *Recept.* nudum. *Pappus* pilosus, copiosus. *Calyx* simplex. *Anthera* apice quinquefidæ.
302. SENECIO. *Recept.* nudum. *Pappus* pilosus, simplex. *Cal.* cylindricus, calyculatus: *Squamis* apice sphacelatis.
301. TUSSILAGO. *Recept.* nudum. *Pappus* pilosus, simplex. *Cal.* squamæ æquales, discum æquant, submembranaceæ.

303. ASTER. *Recept.* nudum. *Pappus* pilosus, simplex. *Cor.* radii plures decem. *Cal.* imbricati Squamæ inferiores patulæ.
311. ANTHEMIS. *Recept.* paleaceum. *Pappus* nullus. *Cal.* hemisphæricus subæqualis. *Flosculi* radii plures quam 5.
312. ACHILLEA. *Recept.* paleaceum. *Pappus* nullus. *Cal.* ovatus imbricatus. *Flosculi* radii circiter 5.

POLYGAMIA FRUSTRANEA.

314. CENTAUREA. *Recept.* fetosum. *Pappus* pilosus, simplex. *Cor.* radii infundibuliformes, longiores, irregulares.
313. COREOPSIS. *Recept.* paleaceum. *Pappus* aristatus. *Cal.* erectus, polyphyllus, basi radiis patentibus cinctus.

POLYGAMIA NECESSARIA.

315. FILAGO. *Recept.* nudum. *Pappus* nullus. *Cal.* imbricatus. *Flosculi* feminei inter squamas calycis.

MONOGAMIA.

316. JASIONE. *Cal.* communis, 40-phyllus. *Cor.* 5-petala, regularis. *Capsf.* infera, 2-locularis.
317. VIOLA. *Cal.* 5-phyllus. *Cor.* 5-petala, irregularis, pone cornuta. *Capsf.* supera, 3-valvis, 1-locularis.
318. IMPATIENS. *Cal.* 2-phyllus. *Cor.* 5-petala, irregularis, *Nectario* cucullato. *Capsf.* supera 5-valvis.



POLYGAMIA ÆQUALIS.

* *Semiflosculosi*: Corollis ligulatis omnibus.

275. TRAGOPOGON. 905.

616. TRAGOPOGON *pratense* calycibus corollæ radium æquantibus: foliis integris strictis. *P. Fl. Oec.* 145.

Dan. *Giæde - Skiæg.*

In campis montosis, ac in horto frequentissima.

Vere variat calyce corollæ radio æquali, majore ac minore, post Solstitium calyces omnium radium corollæ superant.

276. SCORZONERA. 906.

617. SCORZONERA *humilis* caule subnudo unifloro: foliis lato - lanceolatis nervosis planis.

Dan. *Slange - Myrder.*

In campo sterili versus pagum *Virum* ac in apricis prati *Jonstrup.*

618. SCORZONERA *purpurea* foliis lineari - subulatis integris planis: pedunculis cylindricis.

In sterili versus *Virum.*

277. PICRIS. 907.

619. PICRIS *Hieracioides* perianthiis laxis: foliis integris.

In agris.

278. SONCHUS. 908.

620. SONCHUS *arvensis* pedunculis calycibusque hispidis subumbellatis: foliis runcinatis basi cordatis.

In argillofis.

621. *SONCHUS oleraceus* pedunculis tomentosis, calycibus glabris. *P. Fl. Oec.* 138.

Dan. *Svinedild*; *Svine-Tidsej*.

In hortis.

α) *SONCHUS levis* laciniatus latifolius. *Baub.*

β) *SONCHUS levis* minor, paucioribus laciniis. *Baub.*

γ) *SONCHUS asper* laciniatus angustifolius. *Læf.*

Obs. Pedunculi modo ante florescentiam, floribus adhuc confertis, tomentosi, mox nudi.

*) * *SONCHUS* pedunculis glabris subumbellatis: calycibus aculeatis.

279. LACTUCA. 909.

622. *LACTUCA virofa* foliis horizontalibus carina aculeatis, dentatis.

Dan. *Stinkende Lactuc.*

Raro.

280. PRENANTHES. 911.

623. *PRENANTHES muralis* flosculis quinis: foliis runcinatis.

In sylvaticis.

281. LEONTODON. 912.

624. *LEONTODON Taraxacum* calyce inferne reflexo: foliis runcinatis, denticulatis levibus.

P. Fl. Oec. 91.

Dan. *Lövetand*; *Præste-Kone*; *Munke-Hoved.*

Pharmac. TARAXACI radix, herba.

In pascuis ubique.

625. *LEONTODON autumnale* caule ramoso: pedunculis squamosis: foliis lanceolatis dentatis integerrimis glabris.

In pratis frequens.

626. * *LEON.*

626. * LEONTODON *hispidum* calyce toto erecto: foliis dentatis integerrimis hispidis: pilis furcatis.
In pratis.

282. HIERACIUM. 913.

* Scapo nudo unifloro.

627. HIERACIUM *Pilosella* foliis integerrimis ovatis; subtus tomentosis: caule repente: scapo unifloro.

P. Fl. Oec. 80.

Dan. *Muus-öre*.

Pharmac. AURICULÆ MURIS *herba*.

In pascuis siccis.

* * Scapo nudo multifloro.

628. HIERACIUM *dubium* foliis integris ovato-oblongis: stolonibus repentibus: scapo nudo multifloro.
Raro.

629. HIERACIUM *Auricula* foliis integerrimis lanceolatis: scapo nudo multifloro: stolonibus reptantibus.

In pratis, ac in horto frequens.

630. † HIERACIUM *cymosum* foliis lanceolatis integris pilosis: scapo subnudo basi piloso: floribus subumbellatis.

631. HIERACIUM *præmorsum* foliis ovatis subdentatis: scapo nudo racemoso: floribus superioribus præmorsis.

Dan. *Högs-Urt*.

In pratis apricis.

* * * Caute folioso.

632. HIERACIUM *murorum* caule ramoso: foliis radicalibus ovatis dentatis: caulino minori.

Norv. *Guld-Sorl-öye*.

In sylvis; frequens in nemore.

a) folio pilosissimo.

b) minus pilosum.

c) HIERACIUM *myopherum*.

Ibidem.

K

633. * *HIERACIUM paludosum* caule paniculato, foliis amplexicaulibus dentatis glabris: calycibus hispidis

In alneto paludoso ad lacum *Bagsværd*.

634. *HIERACIUM Sabaudum* caule erecto multifloro: foliis ovato-lanceolatis, dentatis, semi-amplexicaulibus.

In nemore.

635. *HIERACIUM umbellatum* foliis linearibus subdentatis sparsis: floribus subumbellatis. *P. Fl. Oec.* 81.

Dan. *Smalblad*; *Högs-Urt.*

In pascuis siccis, imprimis ad molendina.

283. CREPIS. 914.

636. *CREPIS tectorum* foliis lanceolato-runcinatis sessilibus lævibus: inferioribus dentatis: caule glabro.

In aridis frequens.

637. † *CREPIS biennis* foliis runcinato-pinnatifidis scabris: basi superne dentatis: calycibus muricatis.

284. HYOSERIS. 916.

638. † *HYOSERIS minima* caule diviso nudo: pedunculis incrassatis. *Fl. Dan.* t. 201.

In campis fabulosis ad oppidum *Slesvici Rii* copiose legi; Icone duplo minorem; ac in agris Secalinis ad urbem *Slesvicum* vulgarem, uti passim *Rensburgum* usque.

285. HYPOCHÆRIS. 918.

639. *HYPOCHÆRIS maculata* caule subnudo: ramo solitario: foliis ovato-oblongis integris dentatis. *Fl. Dan.* t. 149.

In montosis pratis pagi *Bagsværd*.

640. HYPOCHÆRIS *glabra* calycibus oblongis imbricatis: caule ramoso nudo: foliis dentato-sinuatis.

Inter Secale & alibi passim.

641. HYPOCHÆRIS *radicata* foliis runcinatis obtusis scabris: caule ramoso nudo lævi: pedunculis squamosis. *Fl. Dan. t. 150.*

In pascuis ac in horto vulgaris.

286. L A P S A N A. 919.

642. LAPSANA *communis* calycibus fructus angulatis: pedunculis tenuibus ramosissimis. *P. Fl. Oec. 88.*

Dan. Hare-Kaal.

In cultis frequens, ac in nemore.

287. C I C H O R I U M. 921.

643. CICHORIUM *Intybus* floribus geminis sessilibus: foliis runcinatis.

Pharmac. CICHOREI radix, herba, flores.

Ad margines agrorum passim.

* * *Capitati.*

288. A R C T I U M. 923.

644. ARCTIUM *Lappa* foliis cordatis inermibus petiolatis.

Dan. Stor-Skreppe.

Pharmac. BARDANÆ radix, herba, femina.

289. S E R R A T U L A. 924.

645. SERRATULA *tinctoria* foliis lyrato-pinnatifidis: pinna terminali maxima: flosculis conformibus.

P. Fl. Oec. 135.

Dan. Enge-Skær.

In pratis.

646. *SERRATULA arvensis* foliis dentatis spinosis.

Dan. *Tidsel*.

In cultis frequens.

290. C A R D U U S. 925.

* *Foliis decurrentibus.*

647. *CARDUUS lanceolatus* foliis decurrentibus pinnatifidis hispidis: laciniis divaricatis: calycibus ovatis spinosis villosis: caule piloso.

Dan. *Tidsel*.

Ad margines viarum.

648. *CARDUUS acanthoides* foliis decurrentibus sinuatis margine spinosis: calycibus pedunculatis foliariis erectis villosis.

Dan. *Engel-Tidsel*.

Passim ad vias.

649. *CARDUUS crispus* foliis decurrentibus sinuatis margine spinosis: floribus aggregatis terminalibus inermibus.

Norv. *Juttel-Tröst*.

In agris.

650. *CARDUUS palustris* foliis decurrentibus dentatis, margine spinosis: floribus racemosis erectis: pedunculis inermibus.

In paludosis frequens.

* * *Foliis sessilibus.*

651. † *CARDUUS marianus* foliis amplexicaulibus hastato-pinnatis spinosis: calycibus aphyllis: spinis canaliculatis duplicato-spinosis.

652. † *CARDUUS heterophyllus* foliis amplexicaulibus lanceolatis ciliatis integris laciniatisque: caule subunifloro: calyce inermi.

653. *CARDUUS acaulis* calyce glabro.

In pascuis aridis frequens.

291. C N I C U S. 926.

654. CNICUS *oleraceus* foliis pinnatifidis carinatis nudis, bracteis subcoloratis integris concavis.
In nemoroso prato.

292. O N O P O R D U M. 927.

655. † ONOPORDUM *Acanthium* calycibus squarrosis: spinis subulatis: foliis ovato-oblongis sinuatis.

293. C A R L I N A. 929.

656. CARLINA *vulgaris* caule multifloro corymbofo: floribus terminalibus: calycibus radio albis.
In monticulis aridis ac campis passim.

294. B I D E N S. 932.

657. BIDENS *tripartita* foliis trifidis: calycibus subfoliosis: feminibus erectis. *P. Fl. Oec.* 39.
Dan. *Brönsel*.
Ad amnem vulgarissima.
658. BIDENS *cernua* foliis lanceolatis amplexicaulibus: floribus cernuis: feminibus erectis.
Ad fossas rarior.

295. E U P A T O R I U M. 935.

659. EUPATORIUM *cannabinum* foliis digitatis.
Pharmac. EUPATORII herba.
Ad aquas.

POLYGAMIA SUPERFLUA.

296. TANACETUM. 944.

660. TANACETUM *vulgare*. foliis bipinnatis incisfis serratis.

Dan. *Reinfand*.

Pharmac. TANACETI herba, flores, femina.

Ad hortum Molitoris.

297. ARTEMISIA. 945.

* *Procumbentes ante florescentiam.*

661. ARTEMISIA *campestris* foliis multifidis linearibus: caulibus procumbentibus virgatis.

Dan. *Vild Abrod*; *Bynke*.

In campis aridis.

662. † ARTEMISIA *maritima* foliis multipartitis tomentosis: racemis cernuis: flosculis femineis ternis.

Dan. *Vild Cypres*.

In maritimis sabulosis viæ *Coagensis* legi, ac insulæ *Thorsinge*.

* * *Erectæ herbaceæ, foliis compositis.*

663. ARTEMISIA *Absinthium* foliis compositis multifidis: floribus subglobosis pendulis: receptaculo villoso. *P. Fl. Oec.* 25.

Dan. *Mal - Urt*.

Pharmac. ABSINTHII VULGARIS herba.

In ruderatis.

664. ARTEMISIA *vulgaris* foliis pinnatifidis planis incisfis: subtus tomentosis: racemis simplicibus: floribus ovatis: radio quinquefloro. *P. Fl. Oec.* 26.

Dan. *Bynke*; *Graabone*.

Pharmac. ARTEMISIE RUBRÆ & ALBÆ herba.

In arvis.

* * * *Foliis simplicibus.*

665. † ARTEMISIA *cærulescens* foliis caulinis lanceolatis integris: radicalibus multifidis: flosculis femineis ternis.

298. GNAPHALIUM. 946.

* *Chrysocoma.*

666. GNAPHALIUM *arenarium* herbaceum: foliis lanceolatis: inferioribus obtusis: caule simplicissimo: corymbo composito.

Dan. *Evigheds - Blomster.*

Pharmac. STOECHADIS CITRINÆ flores.

In arvis arenosis.

* * *Argyrocoma.*

667. GNAPHALIUM *dioicum* sarmentis procumbentibus: caule simplicissimo: corymbo simplici terminali: floribus dioicis.

Dan. *Hare - Fod.*

Pharmac. GNAPHALII flores.

In campis aridis.

* * * *Filaginoïdea.*

668. GNAPHALIUM *sylvaticum* caule simplicissimo: floribus sparsis.

Dan. *Muse - Urt.*

In arenoso Nemoris frequens.

669. GNAPHALIUM *uliginosum* herbaceum: caule ramoso diffuso: floribus confertis terminalibus.

In paludosis.

299. CONYZA. 950.

670. † CONYZA *squarrosa* foliis lanceolatis acutis:
caule annuo corymboso.

Dan. *Trold-Urt*; *Torden-Urt*.

In aridis.

Intra patriam nunquam legi.

In rimis petrae vulgo *Winterkastan Hassia* copiose.

300. ERIGERON. 951.

671. ERIGERON *acre* pedunculis alternis unifloris.

Dan. *Blaa Trold-Urt*.

In montosis exaridis.

301. TUSSILAGO. 952.

672. TUSSILAGO *Farfara* scapo imbricato unifloro:
foliis subcordatis angulatis denticulatis.

P. Fl. Oec. 148.

Dan. *Heste-Hov*; *Fölle-Fod*.

Pharmac. TUSSILAGINIS s. FARFARÆ radix, herba.

In argillosis ubique vulgarissima.

673. † TUSSILAGO *Petasites* thyrsis ovato: flosculis
omnibus hermaphroditis.

Pharmac. PETASITIDIS radix, flores.

302. SENECIO. 953.

* Floribus *flosculosis*.

674. SENECIO *vulgaris* corollis nudis: foliis pinnato-
sinuatis amplexicaulibus: floribus sparsis.

Dan. *Kaars-Urt*.

In cultis, viisque ubique.

* * Floribus *radiatis* : radio *revoluto*.

675. † *SENECIO viscosus* corollis revolutis : foliis pinnatifidis viscidis : squamis calycinis laxis longitudine perianthii.

676. * *SENECIO sylvaticus* corollis revolutis : foliis pinnatifidis denticulatis : caule corymbofo erecto.
In pratis sylvaticis.

*** Floribus *radiatis* : radio *patente* : foliis *pinnatifidis*.

677. *SENECIO Jacobæa* corollis radiantibus : foliis pinnato-lyratis : laciniis lacinulatis : caule erecto.

P. Fl. Oec. 134.

Dan. St. Jacobs - Urt.

In nemore.

**** Floribus *radiatis* : radio *patente* : foliis *indivisis*.

678. † *SENECIO paludosus* corollis radiantibus : foliis ensiformibus acute ferratis subtus subvillosis : caule stricto.

303. A S T E R. 954.

679. † *ASTER Tripolium* foliis lanceolatis integerrimis carnosis glabris : ramis inæquatis : floribus corymbolis.

Dan. Strand - Stierne - Urt.

In sabulosis maritimis ad Villas *Fionia Lundsgaard & Hindsgaul*.

304. S O L I D A G O. 955.

680. *SOLIDAGO Virgaurea* caule subflexuoso angulato : racemis paniculatis erectis confertis.

Dan. Gylden Vund - Urt.

Pharmac. VIRGAUREÆ herba.

In nemore raro. In luco *Kalven* ad *Dronninggaard* copiose.

305. CINERARIA. 957.

681. CINERARIA *palustris* foliis lato-lanceolatis, dentato-sinuatis, caule villoso.

OTHONNA *palustris* Specier. edit. prior.

Ad lacum Lyngbye frequens.

306. INULA. 956.

682. INULA *Helenium* foliis amplexicaulibus ovatis rugosis: subtus tomentosis: calycum squamis ovatis.

Dan. *St. Ellens* - Roed.

Pharmac. ENULÆ radix.

Passim ad maceries.

683. † INULA *Oculus Christi* foliis amplexicaulibus oblongis integerrimis hirsutis: caule piloso corymboso.

684. † INULA *Britannica* foliis amplexicaulibus lanceolatis distinctis ferratis subtus villosis: caule ramoso villoso erecto.

685. † INULA *dysenterica* foliis amplexicaulibus cordato-oblongis subtomentosis: caule villoso paniculato: squamis calycinis fetaceis.

686. † INULA *pulicaria* foliis amplexicaulibus undulatis: caule prostrato: floribus subglobosis.

687. † INULA *salicina* foliis sessilibus lanceolatis recurvis ferrato-scabris: floribus inferioribus altioribus: ramis subangulatis glabris.

307. ARNICA. 958.

688. † ARNICA *montana* foliis ovatis integris: caulinis geminis oppositis. *Fl. Dan.* t. 63. *P. Fl. Oec.* 24.

Dan. *Volverley*; *Guldblomme*.

Pharmac. ARNICÆ herba, flores.

308. BELLIS. 962.

308. BELLIS *perennis* scapo nudo. P. Fl. Oec. 35.
 Dan. *Gaasc-Urt*; *Tusind-Fryd*; *Tusind-Dyder*.
Pharmac. BELLIDIS MINORIS herba.
 Ad vias ubique.

309. CHRYSANTHEMUM. 966.

309. CHRYSANTHEMUM *Leucanthemum* foliis amplexicaulibus oblongis: superne ferratis: inferne dentatis.
 Dan. *Oxe-øye*.
Pharmac. BELLIDIS PRATENSIS herba.
 In pratis ac arvis frequens.
310. CHRYSANTHEMUM *segetum* foliis amplexicaulibus: superne laciniatis: inferne dentato-ferratis.
 P. Fl. Oec. 51.
 Dan. *Onde-Urter*; *Aager-Urt*; *Brandeborger*; *Fattigmands-Trudsel*.
 In agris sat frequens.

310. MATRICARIA. 967.

310. † MATRICARIA *Parthenium* foliis compositis planis: foliolis ovatis incisis: pedunculis ramosis.
Pharmac. MATRICARIÆ herba.
 In maceriis pagi circa *Rensburgum* legi.
311. MATRICARIA *Chamomilla* receptaculis conicis: radiis patentibus: seminibus nudis: squamis calycinis margine æqualibus.
 Dan. *Kameel-Blomster*.
Pharmac. CHAMOMILLE NOSTR. herba, flores.
 In ruderatis passim.

694. *MATRICARIA inodora* receptaculis hemisphæricis: radiis patentibus: feminibus coronato-marginatis: squamis calycinis margine exfoletis.

Juxta vias.

Hanc *Linnaeus* in altera *Specierum* editione omisit, etiam si in *Fl. Suec.* differentiam a *Cotula* & *Chamomilla* tradiderat.

695. † *MATRICARIA maritima* receptaculis hemisphæricis: foliis bipinnatis subcarnosis: supra convexis, subtus carinatis.

311. ANTHEMIS. 970.

696. *ANTHEMIS arvensis* receptaculis conicis: paleis fetaceis: feminibus coronato-marginatis.

In agris.

697. *ANTHEMIS Cotula* receptaculis conicis: paleis fetaceis: feminibus nudis. *Pauli Fl. Oec.* 20.

Dan. *Gaafe - Urt*; *Hunde - Urt*.

Ad pagum *Virum* passim.

312. ACHILLEA. 971.

698. *ACHILLEA Ptarmica* foliis lanceolatis acuminatis argute ferratis.

Dan. *Hvid Regn - Fan*.

Pharmac. PTARMICÆ herba.

In ripa paludosa nemoris.

699. *ACHILLEA Millefolium* foliis bipinnatis nudis: laciniis linearibus dentatis: caulibus fulcatis.

Pauli Fl. Oec. 2.

Dan. *Röllike*.

Pharmac. MILLEFOLII herba, flores.

In horto frequens, & alibi passim.

POLYGAMIA FRUSTRANEA.

313. COREOPSIS. 981.

700. † COREOPSIS *bidens* foliis lanceolatis ferratis oppositis amplexicaulibus.

314. CENTAUREA. 984.

* Calycinis squamis *ferrato-ciliatis*.

701. † CENTAUREA *Phrygia* calycibus recurvato-plumosis: foliis indivisis.

Dan. *Knop-Urt.*

Extra urbem *Colding* meridiem versus copiose.

702. CENTAUREA *Cyanus* calycibus ferratis: foliis linearibus integerrimis: infimis dentatis.

P. Fl. Oec. 47.

Dan. *Korn-Blomster*; *Blaa-Urt.*

Pharmac. CYANI flores.

Inter segetes.

703. CENTAUREA *scabiosa* calycibus ciliatis: foliis pinnatifidis: pinnis lanceolatis.

In pratis passim.

* * Calycinis squamis *membranaceis, aridis, integerrimis, muticis.*

704. CENTAUREA *Jacea* calycibus scariosis laceris: foliis lanceolatis radicalibus sinuato-dentatis: ramis angulatis. *P. Fl. Oec.* 48.

Dan. *Knop-Urt.*

In agris.

POLYGAMIA NECESSARIA.

315. FILAGO. 995.

705. FILAGO *pyramidata* caule dichotomo: floribus

758 *Syngenesia. Polygamia necessaria. Monogamia.*

pyramidalis pentagonis axillaribus: flosculis feminis feriatis.

Dan. *Muse - Urt.*

706. *FILAGO montana* caule erecto diviso: floribus conicis lateralibus axillaribus.

In aridis fabulosis.

MONOGAMIA.

316. *JASIONE*. 1005.

707. *JASIONE montana*.

In campis aridis.

317. *VIOLA*. 1007.

* *Acaules.*

708. *VIOLA hirta* acaulis: foliis cordatis pilosis-hispidis.

In horto vetere.

709. *VIOLA palustris* acaulis: foliis reniformibus.

Fl. Dan. t. 83.

In paludosis haud frequens.

710. *VIOLA odorata* acaulis: foliis cordatis: stolonibus reptantibus. *P. Fl. Oec. 156.*

Dan. *Marts - Violer.*

Pharmac. VIOLE flores.

In horto ac nemore.

* * *Caulescentes.*

711. *VIOLA canina* caule aultiore adscendente: foliis oblongo-cordatis.

In sylvestribus frequens.

712. *VIOLA tricolor* caule triquetro diffuso: foliis oblongis incis: stipulis dentatis.

Ad margines in horto.


713. † VIOLA *mirabilis* caule triquetro: foliis reniformi-cordatis: floribus caulinis apetalis.

318. I M P A T I E N S. 1008.

714. IMPATIENS *noli tangere* pedunculis multifloris solitariis: foliis ovatis: geniculis caulinis tumens.

Dan. *Spring-Urt*; *Uroe*.

In fylva Næffet ad ripam ac ad maceriem fylvæ Kollæ kolle.



Classis XX.

GYNANDRIA.

DIANDRIA.

319. ORCHIS. *Nectarium* corniforme pone florem.
 320. SATYRIUM. *Nect.* scrotiforme s. inflato - didymum, pone florem.
 321. OPHRYS. *Nect.* subtus subcarinatum.
 322. SERAPIAS. *Nect.* ovatum, subtus gibbum: labio ovato.
 323. CYPRIPIEDUM. *Nect.* ventricosum, inflatum cavum.

POLYANDRIA.

325. CALLA. *Spatha* plana. *Spadix* tectus flosculis. *Cal.* o. *Cor.* o. *Stam.* mixta pistillis. *Bacca* polyspermæ.
 324. ARUM. *Spatha* monophylla, cucullata. *Spadix* supra nudus inferne femineus, medio stamineus. *Cal.* o. *Cor.* o. *Stam.* supra pistilla.
 326. ZOSTERA. *Spadix* linearis altero latere fructifero. *Cal.* o. *Cor.* o. *Stam.* alterna. *Sem.* solitaria, alterna.



DIANDRIA.

DIANDRIA.

319. ORCHIS. 1009.

* Bulbis indivisis.

715. ORCHIS *bifolia* bulbis indivisis: nectarium labio lanceolato integerrimo: cornu longissimo: petalis patentibus.

Fl. Dan. t. 235. Vaill. Bot. Paris. t. 30. f. 7.

Dan. Gögs-Urt.

Norv. Rams, Valmosöster.

In pascuis sylvaticis haud infrequens; in prato nemoroso ad Hummeltofte.

716. ORCHIS *Morio* bulbis indivisis: nectarium labio quadrifido crenulato: cornu obtuso: petalis omnibus conniventibus. Vaill. Bot. Paris. t. 31. f. 13. 14.

Pharmac. SATYRII radix.

Passim, ut in subhumido umbroso ad lacum Verelöse.

717. ORCHIS *mascula* bulbis indivisis: nectarium labio quadrilobo crenulato: cornu obtuso: petalis dorsalibus reflexis. Vaill. Bot. Paris. t. 31. f. 12.

In luco ad lacum Verelöse.

718. † ORCHIS *militaris* bulbis indivisis: nectarium labio quinquefido, punctis scabro: cornu obtuso: petalis confluentibus.

Vaill. Bot. Paris. t. 31. f. 21. 27. 28.

Dan. Hunde-Nosser.

In præcipitiis promontorii insulæ Mona passim legitur speciosissimam plantam.

719. ORCHIS *ustulata* bulbis indivisis: nectarium labio quadrifido, punctis scabro: cornu obtuso: petalis distinctis. Vaill. Bot. Paris. t. 31. f. 35. 36.

* * Bulbis palmatis.

720. † ORCHIS *latifolia* bulbis subpalmatis rectis: neu

L

nectarii cornu conico: labio trilobo lateribus reflexo: bracteis flore longioribus.

Vaill. Bot. Paris. t. 31. f. 1-5.

Dan. *Guds-Haand.*

In prato humido ad radicem promontorii *Mona* legi.

721. * ORCHIS *incarnata* bulbis palmatis: nectarii cornu conico: labio obscure trilobo serrato: petalis dorsalibus reflexis.

In pratis subhumidis passim.

722. ORCHIS *maculata* bulbis palmatis patentibus: nectarii cornu germinibus brevior: labio plano: petalis dorsalibus patulis.

Vaill. Bot. Paris. t. 31. f. 9. 10.

Norv. *Mari-Gras; Orme-Gras.*

In pratis.

723. ORCHIS *conopsea* bulbis palmatis: nectarii cornu fetaceo germinibus longior: labio trifido: petalis duobus patentissimis.

Fl. Dan. t. 224. Vaill. Bot. Paris. t. 30. f. 88.

Semel modo legi loci haud memor.

320. SATYRIUM. 1010.

724. † SATYRIUM *viride* bulbis palmatis: foliis oblongis obtusis: nectarii labio lineari trifido: lacinia intermedia obsoleta. *Fl. Dan. t. 77.*

Dan. *Guds-Haand.*

725. † SATYRIUM *albidum* bulbis fasciculatis: foliis lanceolatis: nectarii labio trifido acuto: lacinia intermedia obtusa. *Fl. Dan. t. 115.*

321. OPHRYS. 1011.

* Bulbis *ramosis.*

726. OPHRYS *Nidus avis* bulbis fibroso fasciculatis:

caule vaginato: nectarii labio bifido. *Fl. Dan. t. 181.*

In quovis coryleto prati *Jonstrup*, ac sub corylis ad tuguria *Ravnehusene* legi potest, planta alibi rara.

In nemore sub fagis viginti duas in plano duarum ulnarum numèravi. Loco priori vel vicies vidi diversis locis initio Junii florentem.

727. *OPHRYS ovata* bulbo fibroso: caule folio: foliis ovatis: nectarii labio bifido. *Fl. Dan. t. 137.*

Dan. *Toblad.*

In pratis sylvaticis.

* * *Bulbis rotundis.*

728. *OPHRYS Monarchis* bulbo globoso: scapo nudo: nectarii labio trifido cruciato. *Fl. Dan. t. 102.*

Dan. *Sygs - Knap.*

In prato humido ad lacum *Bagsværd.*

729. † *OPHRYS anthropophora* bulbis subrotundis: scapo folioso: nectarii labio lineari tripartito: medio elongato bifido.

Fl. Dan. t. 103. Vaill. Bot. Paris. t. 31. f. 19. 20.

322. SERAPIAS. 1012.

730. *SERAPIAS Helleborine* bulbis fibrosis: floribus erectis bractea brevioribus.

Dan. *Vild Nyffe - Ræd.*

In nemore ad paludem minimam.

731. *SERAPIAS longifolia* bulbis fibrosis: floribus erectis bractea longioribus.

Helleborine floribus obsolete carneis: labello obtuso. *Haller. Helv. 275; Scopoli Carr. 256.*

In paludosis lacus *Lyngby.*

323. CYPRIPIEDUM. 1015.

732. † CYPRIPIEDUM *Calceolus* radicibus fibrosis: foliis ovato-lanceolatis caulinis.

In præcipitiis promontorii insulæ *Mona*, vulgo *Klint*, legi rarissimam plantam.

POLYANDRIA.

324. ARUM. 1028.

733. † ARUM *maculatum* acaule: foliis hastatis integerrimis: spadice clavato. *P. Fl. Oec.* 27.

Dan. *Arum*; *Dansk - Ingfer*.

In sylva ad villam *Lundsgaard Fionia* legi.

325. CALLA. 1030.

734. CALLA *palustris* foliis cordatis: spatha plana: spadice undique hermaphrodito. *P. Fl. Oec.* 41.

Dan. *Vand - Ingfer*.

In paludoso amnis frequens.

326. ZOSTERA. 1032.

735. ZOSTERA *marina*.

Fl. Dan. t. 15. *P. Fl. Oec.* 158.

Dan. *Tang*; *Strand - Gras*.

In mari Balthico.



Classis XXI.

MONOECIA.

MONANDRIA.

327. ZANNICHELLIA. *Masc. Cal. 0. Cor. 0.*
Fœminei. Cal. 1 - phyllus. Cor. 0. Germina cir-
citer 4. Sem. totidem.
 † *Callitriche verna.*

DIANDRIA.

328. LEMNA. *Masc. Cal. 1 - phyllus. Cor. 0.*
Fœm. Cal. 1 - phyllus. Cor. 0. Styl. 1. Caps. 1 - lo-
cularis.

TRIANDRIA.

331. CAREX. *Masc. Amentum imbricatum. Cal. 1 -*
phyllus. Cor. 0.
Fœm. Amentum imbricatum. Cal. 1 - phyllus. Cor. 0.
Nectarium inflatum, 3 - dentatum. Stigm. 3.
Sem. triquetrum intra Nectarium.
 330. SPARGANIUM. *Masc. Amentum subrotundum.*
Cal. 3 - phyllus. Cor. 0.
Fœm. Amentum subrotundum. Cal. 3 - phyllus.
Cor. 0. Stigm. bifidum. Drupa exsucca, 2 - sperma.
 329. TYPHA. *Masc. Amentum cylindricum. Cal. 3 -*
phyllus. Cor. 0.
Fœm. Amentum cylindricum infra masculos. Cal. ca-
pillo villoso. Cor. 0. Sem. 1. insidens pappo ca-
pillari.



TETRANDRIA.

333. URTICA. *Masc.* Cal. 4-phyllus. Cor. o. *Nectarium* centrale, cyathiforme.
Fœm. Cal. 2-valvis. Cor. o. *Stigm.* villosum. *Sem.* 1, ovatum, nitidum.
332. BETULA. *Masc.* *Amentum* 3-florum. Cor. 4-partita. Cal. 1-phyllus, 3-fidus, 3-florus.
Fœm. *Amentum* 2-florum. Cor. o. *Styl.* 2. Cal. 1-phyllus sub 3-fidus, 2-florus. *Strobilus* cylindricus. *Sem.* 1, ovatum, utrinque membrana alata.

PENTANDRIA.

334. XANTHIUM. *Masc.* Cal. communis, polyphyllus, imbricatus. Cor. 1-petalæ, 5-fidæ, infundibulif. *Filam.* connexa. *Recept.* paleaceum.
Fœm. Cal. involucrum, 2-phyllum, 2-florum. Cor. o. *Styl.* 2. *Drupe* sicca, muricata, 2-fida. *Nucleus* bilocularis.

HEXANDRIA.

† *Rumex spinosus.*

POLYANDRIA.

337. SAGITTARIA. *Masc.* Cal. 3-phyllus. Cor. 3-petala. *Stamina* circiter 24.
Fœm. Cal. 3-phyllus. Cor. 3-petala. *Germ.* plurima. *Sem.* numerosa, nuda.
336. MYRIOPHYLLUM. *Masc.* Cal. 4-phyllus. Cor. o. *Stam.* 8.
Fœm. Cal. 4-phyllus. Cor. o. *Pist.* 4. *Styl.* o. *Sem.* 4, nuda.

MONANDRIA.

327. ZANNICHELLIA. 1034.

736. ZANNICHELLIA *palustris*.

Mich. Gen. t. 34. f. 1. 2. Fl. Dan. t. 67.

In fossis.

DIANDRIA.

328. LEMNA. 1038.

737. LEMNA *trifulca* foliis petiolatis, lanceolatis.

Mich. Gen. t. 11. f. 5.

In fossis.

738. LEMNA *polyrbiza* foliis sessilibus: radicalibus confertis. *Mich. Gen. t. 11. f. 1. P. Fl. Oec. 90.*

Dan. Andemad.

In fossis; copiosissima ad paludosa amnis in sylva *Næfset*.

739. LEMNA *minor* foliis sessilibus utrinque planiusculis: radicibus solitariis. *Mich. Gen. t. 11. f. 2.*

In aquis quietis ripæ lacus *Lyngbye*.

TRIANDRIA.

329. TYPHA. 1040.

740. TYPHA *latifolia* foliis subensiformibus: spica mascula foemineaque approximatis.

Dan. Donnemors - Kõlle.

In paludosis passim.

741. † TYPHA *angustifolia* foliis semicylindricis: spica mascula foemineaque remotis.

330. SPARGANIUM. 1041.

742. SPARGANIUM *erectum* foliis erectis triquetris.
 Dan. *Pindsvin - Knoppe*.
 In paludoso ad hortum ac in lacu *Huulsøe* frequens.
743. SPARGANIUM *nataus* foliis decumbentibus planis.
 In paludibus passim.

331. CAREX. 1046.

* Spica *unica simplici*.

744. CAREX *dioica*, spica simplici dioica.
Mich. Gen. t. 32 f. 1.
 In pratis humidis.
745. CAREX *pulicaris* spica simplici androgyna: superne mascula: capsulis divaricatis retro flexis.
Fl. Dan. t. 166.
746. * CAREX *capitata* spica simplici androgyna ovata: superne mascula: capsulis imbricato-patulis.
 In insula paludosa lacus *Bagsværd*.

* * Spicis *androgynis*.

747. * CAREX *arenaria* spica composita, spiculis androgynis: inferioribus remotioribus foliolo longiori instructis: culmo triquetro.
Mich. Gen. t. 33. f. 4. Scheuchz. Agrost. t. 11. f. 6.
 Ad lacum *Lyngby - Søe* legi ac in maritimo sabuloso ad diversorium vulgo *Kiøge - Kro* copiose.
748. CAREX *leporina* spica composita: spiculis ovatis sessilibus, approximatis alternis androgynis nudis.
Scheuchz. Agrost. t. 10. f. 15.
 In pratis humidis.
749. * CAREX *brizoides* spica composita disticha nuda: spiculis androgynis oblongis contiguis: culmo nudo.
 In paludoso nemoris.

L 5



750. * CAREX *uliginosa* spica composita spiculis androgynis: inferioribus remotioribus foliolo longiori instructis: culmo tereti.

Ubi legerim haud jam succurrit.

Spica in mea spiculis ad 20 usque composita.

751. * CAREX *vulpina* spica supradecomposita inferne laxiore: spiculis androgynis ovatis sessilibus glomeratis: superne masculis. *Mich. Gen. t. 33. f. 13. 14.*

752. * CAREX *muricata* spiculis subovatis sessilibus distinctis androgynis: capsulis acutis divergentibus spinosis. *Mich. Gen. t. 33. f. 11.*

In subhumido nemoris.

753. * CAREX *loliacea* spicis subovatis sessilibus remotis androgynis: capsulis ovatis teretiusculis muticis divaricatis.

In humentibus nemoris.

754. CAREX *axillaris* spiculis ovatis sessilibus remotis androgynis: bracteis culmum æquantibus.

Morif. Hist. III. t. 12. f. 17.

Passim in ripa paludosa annis.

755. * CAREX *canescens* spiculis subrotundis remotis sessilibus obtusis androgynis: capsulis ovatis obtusiusculis.

In sylvis passim.

756. * CAREX *remota* spiculis oblongis sessilibus remotis androgynis: foliis caulinis culmum æquantibus.

In paludoso sub alnis ad viam versus lacum *Varelöse*.

757. * CAREX *elongata* spiculis oblongis sessilibus remotis androgynis: capsulis ovatis acutis.

Scheuchz. Agrost. t. 11. f. 4.

In paludosis ad *Valhalla* haud vulgaris.

*** Spicis sexu distinctis: foemineis sessilibus.

758. CAREX *flava* spicis confertis subsessilibus subrotundis: mascula lineari: capsulis acutis recurvis. In paludibus.

759. * *CAREX digitata* spicis linearibus erectis: mascula brevior inferiorique: bracteis aphyllis: capsulis distantibus. *Mich. Gen. t. 32. f. 9.*

In præcipitiis cretaceis promontorii insulæ *Monæxopio* legi, ac in præcipitiis ad lacum *Fuursoë*.

760 * *CAREX montana* spicis femineis sessilibus foliariis ovatis masculæ approximatis: culmo nudo: feminibus pubescentibus.

In locis elatioribus.

761. * *CAREX globularis* spica mascula oblonga: feminea sessili ovata foliolo florali breviori approximata. *Mich. Gen. t. 32. f. 3.*

In campis ac collibus apricis frequens.

762. * *CAREX filiformis* spica mascula oblonga: femineis sessilibus oblongis: inferiore foliolo proprio brevior. *Scheuchz. Agrost. t. 10. f. 11.*

763. * *CAREX pilulifera* spicis terminalibus confertis subrotundis: masculina oblonga.

Raro.

*** Spicis sexu distinctis: femineis pedunculatis.

764. * *CAREX limosa* spicis ovatis pendulis: mascula longiore, erectiore: radice repente.

Scheuchz. Agrost. t. 10. f. 15.

In paludibus sylvaticis.

765. * *CAREX capillaris* spicis pendulis: mascula erecta: femineis oblongis distichis: capsulis nudis acuminatis. *Fl. Dan. t. 168. g.*

In subhumidis sylvestribus; ad paludem minorem.

766. *CAREX pallescens* spicis pendulis: mascula erecta: femineis ovatis imbricatis: capsulis confertis obtusis. *Mich. Gen. t. 39. f. 12.*

In paludibus.

767. * CAREX *panicea* spicis pedunculatis erectis remotis: femineis linearibus: capsulis obtusiusculis inflatis. *Mich. Gen. t. 32. f. 11.*

In uliginosis.

768. CAREX *pseudo-Cyperus* spicis pendulis: pedunculis geminatis.

Dan. *Cyper-Græs*, nomen pluribus commune.

In paludoso lacus *Lyngby*.

769. * CAREX *caspitosa* spicis erectis cylindricis ternis subsessilibus: mascula terminali: culmo triquetro.

In paludibus turfosis.

770. * CAREX *distans* spicis remotissimis subsessilibus bractea vaginante: capsulis angulatis mucronatis.

In paludibus.

***** Spicis sexu distinctis: masculis pluribus.

771. CAREX *acuta* spicis masculis pluribus: femineis subsessilibus: capsulis obtusiusculis.

Michel. Gen. t. 32. f. 12.

In paludibus ac pratis subhumidis vulgarissima.

772. CAREX *vesicaria* spicis masculis pluribus: femineis pedunculatis: capsulis inflatis acuminatis.

Mich. Gen. t. 32. f. 6. 7.

Norv. *Skiane*.

In udibus nemorosis.

773. * CAREX *hirta* spicis remotis: masculis pluribus femineis subpedunculatis erectis: capsulis hirtis.

In fabulosis.

T E T R A N D R I A.

332. BETULA. 1052.

774. BETULA *alba* foliis ovatis acuminatis ferratis.

P. Fl. Oec. 36.

Dan. *Birk*.

In ripa lacus *Fuursoë* raro; in paludosis lacus *Lyngby* frequens; ad *Birckeröd* copiosissima.



775. *BETULA Abnus* pedunculis ramosis. *P. Fl. Oec.* 38.
 Dan. *Ell.*
 Ubique ad paludosa & ripas.

333. URTICA. 1054.

776. *URTICA urens* foliis oppositis ovalibus.
 Dan. *Nælde.*
 Norv. *Brande - Hatte.*
 In cultis ubique.
777. *URTICA dioica* foliis oppositis cordatis : racemis
 geminis.
 Dan. *Nælde.*
Pharmac. URTICÆ MAJORIS radix, herba, semina.
 In maceris.

PENTANDRIA.

334. XANTHIUM. 1056.

778. *XANTHIUM strumarium* caule inermi : foliis cor-
 datis trinervatis.
 Dan. *Gaase - Skreppe.*
 In fœtibus passim.

POLYANDRIA.

335. CERATOPHYLLUM. 1065.

779. *CERATOPHYLLUM demersum* foliis dichotome
 bigeminis : fructibus trispinosis.
 In fossis passim.

336. MYRIOPHYLLUM. 1066.

780. *MYRIOPHYLLUM spicatum* floribus masculis in-
 terupte spicatis.
 In amne.

781. MYRIOPHYLLUM *verticillatum* floribus omnibus
verticillatis.
Ibidem ad molendinum.

337. SAGITTARIA. 1067.

782. SAGITTARIA *sagittifolia* foliis sagittatis acutis;
Fl. Dan. t. 172.
Dan. Pül.-Urt.
In fossa ad molendinum novum.

338. POTERIUM. 1069.

783. POTERIUM *sanguisorba* inerme: caulibus suban-
gulosis.
In asperis incultis ad hortum.

339. QUERCUS. 1070.

784. QUERCUS *Robur* foliis deciduis oblongis, super-
ne latioribus: sinibus acutioribus: angulis obtu-
sis. *P. Fl. Oec. 119.*
Dan. Eeg.
Pharmac. QUERCUS cortex, folia, glandes, cupulæ.
Passim; frequens in sylva versus Tuguria *Razne-
buse.*

340. FAGUS. 1072.

785. FAGUS *sylvatica* foliis ovatis obsolete ferratis.
R. H. Oec. 69.
Dan. Bög.
Pharmac. FAGI fructus.
Vulgatissima, utilissima, gratissimaque *Danis Arbor.*

341. CARPINUS. 1073.

786. † CARPINUS *Betulus* squamis strobilorum planis.
P. Fl. Oec. 45.
Dan. Aunbög.

342. CORYLUS. 1074.

787. CORYLUS *Avellana* stipulis ovatis obtusis.
P. Fl. Oec. 57.
Dan. Hassel.
In pratis passim.

SYNGENESIA.

343. BRYONIA. 1093.

788. BRYONIA *alba* foliis palmatis utrinque callosè
 scabris.
Dan. Hunde - Bær ; Galle - Bær.
Pharmac. BRYONIÆ radix, baccæ, femina.
In sepibus passim.



Classis XXII.

D I O E C I A.

DIANDRIA.

344. SALIX. *Masc. Amenti squamæ. Cor. o. Glandula* baseos nectarifera. *Stam. 2 - 5.*
Fœm. Amenti squamæ. Cor. o. Stylus 2 - fidus.
Caps. 1 - locularis, 2 - valvis. Sem. papposa.

TRIANDRIA.

345. EMPETRUM. *Masc. Cal. 3 - partitus. Cor. 3 - petala. Stam. longa.*
Fœm. Cal. 3 - partitus. Cor. 3 - petala. Styli 9. Bacca 9 - sperma.
 † Valeriana dioïca.
Fœm. Cal. o. Cor. 2 - petala minor. Styl. 1.
 † Carex dioïca.
Fœm. Amentum imbricatum. Cal. 1 - phyllus. Stigm. 3.
Sem. triquetrum.

TETRANDRIA.

347. HIPPOPHAË. *Masc. Cal. 2 - partitus. Cor. o.*
Fœm. Cal. 2 - fidus. Cor. o. Styl. 1. Bacca 1 - sperma.
 346. VISCUM. *Masc. Cal. 4 - partitus. Cor. o. Filamenta o. Antheræ calyci adnatæ.*
Fœm. Cal. 4 - phyllus, superus. Styl. o. Stigma obtusum. Cor. o. Bacca 1 - sperma, infera. Sem. cordatum.
 † Urtica dioïca.
Fœm. Styl. 1. Stigma villosum.

348. MYRI-

348. MYRICA. *Masc.* *Amenti* squama lunata. *Cor.* 0.
Fem. *Amenti* squama lunata. *Cor.* 0. *Styli* 2. *Bacca*
 1 - sperma.

PENTANDRIA.

349. HUMULUS. *Masc.* *Cal.* 5 - phyllus. *Cor.* 0.
Fem. *Cal.* 1 - phyllus, oblique patens, integer. *Cor.*
 0. *Styli* 2. *Sem.* 1. intra calycem planum.
 † Rhamnus catharticus.
 † Salix pentandra.

HEXANDRIA.

- † RUMEX acetosa. 2. *Fem.* *Cal.* 3 - phyllus. *Cor.* 3 - petala.
 † RUMEX acetosella. 5. *Styl.* 3. *Sem.* 1. triquetrum.

OCTANDRIA.

350. POPULUS. *Masc.* *Amenti* *Cal.* lamina lacera.
Cor. turbinata, obliqua, integra *Stamina* 8-16.
Fem. *Amenti* *Cal.* & *Cor.* maris. *Stigma* 4 - fidum.
Caps. 2 - locularis. *Sem.* multa papposa.

ENNEANDRIA.

351. MERCURIALIS. *Masc.* *Cal.* 3 - partitus. *Cor.* 0.
Stam. 9. seu 12. *Antheræ* globosæ didymæ.
Fem. *Cal.* 3 - partitus. *Cor.* 0. *Styli* 2. *Caps.* dicoc-
 ca, 2 - locularis, 1 - sperma.
 352. HYDROCHARIS. *Masc.* *Spatba* 2 - phylla.
Cal. 3 - fidus. *Cor.* 3 - petala. *Filamenta* 3. inte-
 riora styliifera.
Fem. *Cal.* 3 - fidus. *Cor.* 3 - petala. *Styli* 6. *Caps.* 6 -
 locularis, polysperma, infera.

D E C A N D R I A.

† LYCHNIS *dioica*.*Fem. Caps. 3 - locul. oblonga. Petala 5 - unguiculata.**Cal. 1 - phyllus, oblongus, lævis. Styli 5.*


I C O S A N D R I A.

† RUBUS *chamemorus*.*Fem. Styli plures. Cal. 5 - fidus. Cor. 5 - petala.*

M O N A D E L P H I A.

353. JUNIPERUS. *Masc. Calyx Amenti squamæ.*
*Cor. o. Stam. 3.**Fem. Cal. 3 - partitus. Cor. 3 - petala. Styli 3. Bacca*
*3 - sperma, tribus tuberculis calycis inæqualis.*354. TAXUS. *Masc. Cal. 3 - phyllus gemmæ. Cor. o.*
*Stamina multa. Antheræ peltatæ 8 - fidæ.**Fem. Cal. 3 - phyllus gemmæ. Cor. o. Styl. o. Sem. 1,*
calyculo baccato, integerrimo.

S Y N G E N E S I A.

† GNAPHALIUM *dioicum*. v. p. 183.

DIANDRIA.

344. S A L I X. 1098.

* *Foliis glabris serratis.*

789. * *SALIX pentandra* foliis serratis glabris: flosculis pentandris.

Dan. *Piil*, nomen generis.

In insula paludosa lacus *Lyngbye* frequens.

β) *SALIX octandra* foliis serratis glabris: flosculis octandris.
In prato humido ad radices promontorii insulæ *Mone* vulgo *Klint*.

Obs. Speciem vix dicam, convenit enim omnino pentandrae flosculis exceptis, qui octandri, raro decandri.

790. *SALIX fragilis* foliis serratis glabris ovato-lanceolatis: petiolis dentato-glandulosis.

Dan. *Skjör - Piil*.

Ad ripas & pagos.

791. *SALIX purpurea* foliis serratis glabris lanceolatis: inferioribus oppositis.

Dan. *Sey - Piil*.

Passim.

* * *Foliis integerrimis, villis.*

792. *SALIX arenaria* foliis integris ovatis acutis: supra subvillosis: subtus tomentosis. *Fl. Dan. t. 197.*

In paludosis lacus *Lyngbye*.

793. *SALIX incubacea* foliis integerrimis lanceolatis: subtus villosis nitidis: stipulis ovatis acutis.

In insula palustri lacus *Lyngbye*.

794. * *SALIX fusca* foliis integris ovatis: subtus villosis nitidis.

In sterili subhumido Sylvæ *Næfset*.

795. * *SALIX Rosmarinifolia* foliis integerrimis lanceolato-linearibus strictis sessilibus subtus tomentosis.

In paludoso lacus *Lyngbye*.

*** Foliis *subserratis*, *villosis*.

796. *SALIX caprea* foliis ovatis rugosis: subtus tomentosis, undatis: superne denticulatis.

Dan. *Viddie*.

In sicco passim.

797. † *SALIX viminalis* foliis subintegerrimis lanceolato-linearibus longissimis acutis subtus sericeis: ramis virgatis.

In pago *Væren* legi raram in *Sialandia* arborem, unde propagari facile possit.

798. † *SALIX cinerea* foliis subserratis oblongo-ovatis: subtus subvillosis: stipulis dimidiato-cordatis.

799. *SALIX alba* foliis lanceolatis acuminatis serratis utrinque pubescentibus: ferraturis infimis glandulosis. *P. Fl. Oec.* 129.

Pharmac. SALICIS folia.

Passim.

TRIANDRIA.

345. EMPETRUM. 1100.

800. *EMPETRUM nigrum* procumbens.

Dan. *Kragebær*.

In ericetis citra *Friedensburgum*, ni fallor; ac in ericetis *Cimbria* copiose legi.

TETRANDRIA.

346. VISCUM. 1105.

801. † *VISCUM album* foliis lanceolatis obtusis: caule dichotomo: spicis axillaribus.

Dan. *Fugle-Liim*.

Pharmac. Visci AUCUPARII lignum.

347. HIPPOPHAË. 1106.

802. † HIPPOPHAË *Rhamnoïdes* foliis lanceolatis.

P. Fl. Oec. 82.

Dan. *Haftorn; Tidse.*

Rarum in patria fruticem legi in præruptis cretaceis
promontorii insulæ *Monæ*, ubi copiose crescit,
ac ad sæpes adhibetur.

348. MYRICA. 1107.

803. MYRICA *Gale* foliis lanceolatis subferratis: caule
suffruticoso. *P. Fl. Oec.* 102.

Dan. *Pors.*

Pharmac. MYRTI BRABANTICÆ folia.

In paludibus raro.

PENTANDRIA.

349. HUMULUS. 1116.

804. HUMULUS *Lupulus*. *P. Fl. Oec.* 83.

Dan. *Humle.*

Pharmac. LUPULI strobili.

In nemore.

OCTANDRIA.

350. POPULUS. 1123.

805. POPULUS *alba* foliis subrotundis dentato-angu-
latis, subtus tomentosis.

Dan. *Poppel - Piil.*

In nemore & ad hortum veterem.

806. POPULUS *tremula* foliis subrotundis, dentato-
angulatis, utrinque glabris. *P. Fl. Oec.* 112.

Dan. *Bevre - Æsk; Æspe - Træ.*

Rarior.

807. † *POPULUS nigra* foliis deltoïdibus acuminatis,
ferratis.

P. Fl. Oec. III.

Dan *Poppel-Træ; Poppel-Piil.*

Pharmac. POPULI oculi.

ENNEANDRIA.

351. MERCURIALIS. 1125.

808. *MERCURIALIS perennis* caule simplicissimo foliis
scabris. *P. Fl. Oec. 100.*

Dan. *Bingel - Urt.*

In nemore ac in septo *Deylighbeden* vulgaris.

352. HYDROCHARIS. 1126.

809. *HYDROCHARIS morsus ranae.*

In aqua pigra nemoris ad hortum.

MONADELPHIA.

353. JUNIPERUS. 1134.

810. † *JUNIPERUS communis* foliis ternis patentibus
mucronatis bacca longioribus. *P. Fl. Oec. 87.*

Dan. *Enebær - Træ.*

Pharmac. JUNIPERI baccæ, lignum.

In præcipitiis promontorii insulæ *Mona* ac in alne-
tis paludosis frequenter, uti quoque in campis mon-
tosis *Fridensburgum* versus legi.

354. TAXUS. 1135.

811. † *TAXUS baccata* foliis approximatis.

Classis XXIII.

POLYGAMIA.

MONOECIA.

355. *HOLCUS. Hermaphrod. Cal. Gluma 2 - flora.*
Cor. Gluma aristata. Stam. 3. Styli 2. Sem. 1.
Masc. Cal. Gluma 2 - valvis. Cor. o. Stam. 3.
356. *PARIETARIA. Hermaphr. Cal. 4 - fidus. Cor. o.*
Stam. 4. Styl. 1. Sem. 1. superum, elongatum.
Fem. Cal. 4 - fidus. Cor. o. Stam. o. Styl. 1. Sem. 1.
superum, elongatum.
357. *ATRIPLEX. Hermaphr. Cal. 5 - phyllus. Cor. o.*
Stam. 5. Styl. 2 - partitus. Sem. 1, depressum.
Fem. Cal. 2 - phyllus. Cor. o. Stam. o. Styl. 2 - par-
titus. Sem. 1. compressum.
358. *ACER. Hermaphr. Cal. 5 - fidus. Cor. 5 - petala.*
Stam. 8. Pist. 1. Caps. 2. f. 3. monospermæ, ala
terminatæ.
Masc. Cal. 5 - fidus. Cor. 5 - petala. Stamina 8.
 † *AVENA elatior.*
Hermaphr. Cal. 2 - valvis. Cor. mutica. Stam. 3.
Styli 2.
Masc. Cor. aristata. Stam. 3.

DIOECIA.

359. *FRAXINUS. Hermaphr. Cal. o - seu 4 - parti-*
tus. Cor. o - seu 4 - petala. Stam. 2. Pist. 1.
Sem. 1. lanceolatum.
Fem. Pist. 1. lanceolatum.
 † *Gnaphalium dioicum.*
Hermaphr. Cal. imbricatus squamis. Cor. tubulosa.
Stam. 5. Styl. 1.
Fem. Pistillum filiforme, bifidum.

MONOE CIA.

355. HOLCUS. 1146.

812. *HOLCUS lanatus* glumis bifloris villosis: hermaphrodito mutico: masculo arista recurva.

In pascuis vulgaris.

813. * *HOLCUS mollis* glumis bifloris nudiusculis: fasciculo hermaphrodito mutico: masculo arista geniculata.

In pratis arenosis rarius, ac in horto *Friedensburger*.

814. * *HOLCUS odoratus* glumis trifloris muticis glabris: hermaphrodito diandro.

In graminosis subhumidis.

356. PARIETARIA. 1152.

815. † *PARIETARIA officinalis* foliis lanceolato-ovatis alternis.

Dan. *St. Peders-Urt*; *Dag og Nat*.

357. ATRIPLEX. 1153.

816. † *ATRIPLEX Portulacoides* caule fruticoso: foliis obovatis.

Dan. *Strand-Portulac*.

817. *ATRIPLEX patula* caule herbaceo patulo: foliis subdeltoideo-lanceolatis: calycibus feminum disco dentatis.

In ruderatis.

818. *ATRIPLEX littoralis* caule herbaceo erecto: foliis omnibus linearibus integerrimis.

Dan. *Strand-Melde*.

In littore maris.

819. † *ATRIPLEX pedunculata* caule herbaceo divaricato : foliis lanceolato - obtusis integris : calycibus femineis pedunculatis.

In maritimis.

358. A C E R. 1155.

820. † *ACER campestre* foliis lobatis obtusis emarginatis.

In sylvis *Fionia*, insulæ *Mona* ac Australis *Sialundia* haud infrequens.

821. † *ACER pseudo-platanus* foliis quinquelobis inæqualiter serratis : floribus racemosis.

Pauli Fl. Oecon. p. 1.

Dan. *Løn*, *Valdbirk*; *Naur*.

D I O E C I A.

359. F R A X I N U S. 1160.

822. *FRAAXINUS excelsior* foliis serratis : floribus apetalis. *P. Fl. Oec. 72.*

Dan. *Æsk*.

Pharmac. FRAXINI lignum, cortex, folia, semina.

Passim juxta vias.



Classis XXIV.

CRYPTOGAMIA.

FILICES.

* Fructificationes *spicatae*.

360. EQUISETUM. *Spica* sparsa. *Fructific.* peltatae basi dehiscences, multivalves.
361. OPHIOGLOSSUM. *Spica* articulata, disticha: articulis transversim dehiscens.
362. OSMUNDA. *Spica* racemosa. *Fructif.* globosae, 2 - valves.

* * Fructificationes *frondosae in pagina inferiore*.

363. ACROSTICHUM. *Fructific.* discum totum frondis tegentes.
366. POLYPODIUM. *Fructific.* in punctis subrotundis, sparsis per discum.
365. ASPLENIUM. *Fructific.* in lineolis disci frondis, sparsis.
364. PTERIS. *Fructific.* in lineis marginalibus.

* * * Fructificationes *radicales*.

367. PILULARIA. Flores *masculi* ad latus frondis. *Fructificatio* feminea ad radicem globosa, 4 - locularis.
368. ISOËTES. Fl. *masculi* anthera intra basin frondis. Fl. *feminei*. *Capsula* 2 - locularis intra basin frondis.

M U S C I.

* *Acalyptrati.*

69. LYCOPODIUM. *Anthera* 2 - valvis , *sessilis*.
Calyptra nulla.
 170. SPHAGNUM. *Anthera* operculata, ore lævi.
Calyptra nulla.

* * * *Calyptrati diclini.*

174. POLYTRICHUM. *Anthera* operculata, apophysi minimæ insidens. *Calyptra* villosa. *Stella feminea* in distincto individuo.
 175. MNIMUM. *Anthera* operculata. *Calyptra* lævis.
Fem. capitulum nudum, pulverulentum, remotum.

* * * *Calyptrati monoclini.*

176. BRYUM. *Anthera* operculata. *Calyptra* lævis.
Pedunculus e tuberculo terminalis.
 177. HYPNUM. *Anthera* operculata. *Calyptra* lævis.
Pedunculus lateralis e perichætio.
 171. PHASCUM. *Anthera* operculata sessilis, ore ciliato. *Calyptra* lævis.
 173. BUXBAUMIA. *Anthera* operculata, obovata, pedunculata, altero latere membranacea. *Calyptra* lævis.
 172. FONTINALIS. *Anthera* operculata, sessilis. *Calyptra* perichætio inclusa.

A L G Æ.

* *Terrestres.*

179. MARCHANTIA. *Masc.* *Cal.* peltatus, subtus tectus corollis monopetalis. *Antheræ* multifidæ.
Fem. *Calyx* sessilis, campanulatus, polyspermus.
Stamina 12-24.

378. JUNGERMANNIA. *Stamina* 12-24. *Pistillum* unicum. *Capsula* globosa, 4-valvis, pedunculata, polysperma.

Observatio. Globulus, qui in JUNGERMANNIIS tempore fructificationis conspicitur pedunculo suffultus, revera pericarpium est, quod maturo semine rumpitur in quatuor valvulas *lit. a.* ac effundit pollinem filamentis capillaribus plerumque insidentem, verum harum plantularum semen Septembri enim vel Octobri, si vagina sive calyx *lit. b* intra quem pedunculus globulum sustinens postea oritur, caute removeatur, in cavitate ejus conspiciere licet filamenta minima fusca *lit. c* duodecim ad viginti quatuor. Hæc nutant centrum versus, quod magis elevatum apparet, ac in quo aliud filamentum distinguitur; procedente tempore e centro enascitur globus valde conspicuus *lit. d*, sub quo adhuc hærent illa filamenta jam exarida, ac a summitate affidet filamentum aliud erectum. Hoc, postquam dissipare filamenta procumbentia fusca, restat interdum ad rupturam globuli usque. Quid illa, nisi stamina? quid hoc, nisi pistillum? ergo quid globus, nisi capsula? quid polvis, nisi semen? Capsula hæc seminifera primo vere tempestate pluvia subito extollitur pedunculo pellucido ad altitudinem pro diversis speciebus 3 linearum ad 4 pollicum, in quatuor finditur partes, *lit. a* ac semina sequentium anni plantulas spargit.

Stamina hæc atque pistilla in JUNGERMANNIA *lanceolata*, *complanata*, *tamarisci*, *platyphylla* ac *epiphylla* sapienter vidi, nec in cæteris dari dubito. vid. *Figuram* 5. & 6.

380. BLASIA. *Masc. Cal.* cylindricus granis repletus. *Fem. Cal.* nudus: fructu subrotundo, foliis immixto, polyspermo.

381. RICCIA. *Masc.* nudus. *Anthera* subulata, truncata. *Fem.* nudus. *Caps.* globosa, 1-locularis, polysperma.
382. LICHEN. *Masc. Receptaculum* subrotundum planiusculum, nitidum.

Fem. Farina foliis adspersa.

389. BYSSUS. *Lanugo* aut pulvis simplicissimus.

* * *Aquaticæ.*

384. TREMELLA. *Alga* gelatinosa: *Fructificationem* vix manifestæ.

386. ULVA. *Alga* membranacea. *Fructif.* in membrana diaphana.
385. FUCUS. *Alga* coriacea. *Masc.* *Vesicula* villis intertextæ.
Fem. *Vesiculæ* adspersæ granis immerfis, apice prominulis. *Sem.* folitaria.
387. CONFERVA. *Alga* capillaris. *Tubercula* inæqualia in fibris capillaribus longissimis.
388. SPONGIA. *Alga* pilis intricatis intertexta.
383. CHARA, *Alga*, caulis teres ramosus, siccitate contortus. *Cal.* 2 - phyllus. *Anthera* globosa, sessilis. *Stigmata* 3. *Sem.* 1. subrotundum.

F U N G I.

* *Pileati.*

390. AGARICUS. *Fungus* horizontalis, subtus lævissimus.
391. BOLETUS. *Fungus* horizontalis, subtus porosus.
392. HYDNUM. *Fungus* horizontalis, subtus echinatus.
393. PHALLUS. *Fungus* supra reticulatus, subtus lævis.

* * *Sessiles.*

394. CLATHRUS. *Fungus* subrotundus cancellatus.
395. ELVELA. *Fungus* turbinatus lævis.
396. PEZIZA. *Fungus* campanulatus.
397. CLAVARIA. *Fungus* oblongus, lævis.
398. LYCOPERDON. *Fungus* subrotundus, seminibus farinaceis repletus.
399. MUCOR. *Fungus* vesiculis stipitatis.

FILICES.

360. E Q U I S E T U M. 1169.

823. E Q U I S E T U M *sylvaticum* caule spicato: frondibus compositis.

Dan. *Heste - Rumpe*, nomen generis.

In pratis sylvaticis.

824. E Q U I S E T U M *arvense* scapo fructificante nudo, sterili frondoso. *P. Fl. Oec. 64.*

Dan. *Stude - Knæ*.

Pharmac. E Q U I S E T I herba.

In arvis.

825. * E Q U I S E T U M *palustre* caule angulato: frondibus simplicibus.

In paludosis.

826. E Q U I S E T U M *fluviale* caule striato: frondibus subsimplicibus. *P. Fl. Oec. 65.*

Ad amnem.

827. E Q U I S E T U M *hyemale* caule nudo scabro: basi subramoso.

Dan. *Skov - Græs*.

Ad amnem.

828. E Q U I S E T U M *limosum* caule subnudo lævi.

In paludibus sylvæ.

361. O P H I O G L O S S U M. 1171.

829. † O P H I O G L O S S U M *vulgatum*, fronde ovata.

Fl. Dan. t. 147.

Dan. *Slange - Tunge*.

362. OSMUNDA. 1172.

* Scapis *insidentibus cauli ad basin frondis.*

130. † OSMUNDA *Lunaria* scapo caulino solitario : fronde pinnata solitaria. *Fl. Dan. t. 18.*

Dan. *Maane - Rude.*

In præcipitiis promontorii insulæ *Monæ* copiose legi.

* * *Fronde ipsa fructificationes ferente.*

131. † OSMUNDA *regalis* frondibus bipinnatis, apice racemiferis. *Fl. Dan. t. 217.*

Dan. *Engel - Bregne.*

In fossa ad viam inter *Zellam & Hannoveram* copiosa.

*** Frondibus *aliis foliaceis : aliis fructificantibus.*

32. † OSMUNDA *Struthiopteris* frondibus pinnatis : pinnis pinnatifidis : scapo fructificante disticho.

Fl. Dan. t. 169.

33. OSMUNDA *Spicant* frondibus lanceolatis pinnatifidis : laciniis confluentibus integerrimis parallelis.

Fl. Dan. t. 99.

Dan. *Milt - Urt.*

In declivi ad amnem in sylva *Nasset* frequens.

363. ACROSTICHUM. 1173.

14. * ACROSTICHUM *Thelypteris* fronde pinnata : pinnis pinnatifidis integerrimis.

In sylva.

364. PTERIS. 1174.

15. PTERIS *aquilina* frondibus supradecompositis : foliolis pinnatis : pinnis lanceolatis : infimis pinnatifidis : superioribus minoribus. *P. Fl. Oss. 117.*

Dan. *Bregne.*

In sylvis & maceriis vulgaris.

365. ASPLENIUM. 1178.

* Fronde *simplici*.

836. † ASPLENIUM *Scolopendrium* frondibus simplicibus cordato-lingulatis integerrimis: stipitibus hirsutis.

Dan. *Hiorte - Tunge*.

Pharmac. SCOLOPENDRII herba.

Ad *Dronninggaard*, ubi *Kylling* raram hanc plantam invenit, quam diligenter sæpius quæsi, at frustra.

* * Fronde *pinnata*.

837. ASPLENIUM *Trichomanes* frondibus pinnatis: pinnis subrotundis crenatis. *Fl. Dan. t. 119.*

Pharmac. ADIANTHI RUBRI herba.

In macerie nemoris lacum *Lyngbye* spectante.

838. † ASPLENIUM *Ruta muraria* frondibus alternatim decompositis: foliolis cuneiformibus crenulatis. *Fl. Dan. t. 190.*

Dan. *Muur - Rude*; *Steen - Rude*.

Pharmac. RUTÆ MURARIÆ herba.

366. POLYPODIUM. 1179.

* * Fronde *pinnatifida*, lobis *coadunatis*.

839. POLYPODIUM *vulgare* frondibus pinnatifidis: pinnis oblongis subserratis obtusis: radice squamata.

Dan. *Steenfød - Rød*; *Engelsød*.

Pharmac. POLYPODII radix.

In præruptis viarum ac præcipitiis frequens.

* * Fronde

* * Fronde *subbipinnata*, *pinnis basi confluentibus*.

840. * POLYPODIUM *Phegopteris* frondibus subbipinnatis: foliolis infimis reflexis: paribus pinnula quadrangulari coadunatis.

Morif. t. 4. f. 17.

In umbrosis.

841. * POLYPODIUM - - - frondibus subbipinnatis: foliolis ovato-oblongis: pinnis ovatis, incis: lacinii apice acute ferratis. vid. *Fig. II.*

In paludosis.

Pinnæ inferiores non confluunt. Incisuræ adeo profundæ, ut pinnatifidæ dicendæ; hi lobuli apice habent denticulum unicum, duas vel tres acutissimas, lateribus nullum. Variat foliolis alternis & oppositis; omnibus floriferis aut folis superioribus; colore saturate ac pallescente viridi.

* * * Fronde *bipinnata*.

842. POLYPODIUM *Filix mas* frondibus bipinnatis: pinnis obtusis crenulatis: stipite paleaceo.

Pharmac. FILICIS radix.

In sylvis ac maceris.

843. POLYPODIUM *Filix femina* frondibus bipinnatis: pinnulis lanceolatis pinnatifidis acutis.

In sylvis.

844. * POLYPODIUM - - - frondibus bipinnatis: foliolis lanceolatis: pinnulis ovatis, lacinia-
tis, obtuse ferratis. vid. *Fig. III.*

In nemore.

Lacinia infima cujusque pinnulæ exterior opposita paulo longior. Stipes flexuosus.

* * * * Fronde *subtripinnata*.

845. * POLYPODIUM - - - frondibus subtripinnatis: pinnis oblongis, subpinnatis: pinnulis utrinque ferratis. vid. *Fig. IV.*

In sylva.

N

Foliola triangularia acuminata. Pinnæ distantes, nec basi confluentes, harum vero pinnulæ basi parum coalitæ, ita, ut frons omnino tripinnata appareat. Pinnulæ in apice & lateribus denticulatæ at minus acute. Folia omnia florifera.

***** Fronde *supradecomposita*.

846. POLYPODIUM *Dryopteris* frondibus *supradecompositis*: foliolis ternis bipinnatis.

Dan. *Enge-Bregne*.

In nemore vulgare.

Tria POLYPODIA No. 841. 844. 845. nullo modo cum *Linneanis* descriptionibus aut aliorum satis juste conciliare potui, hinc peritis descriptiones ac figuræ foliorum traduntur.

367. PILULARIA. 1183.

847. PILULARIA *globulifera*.

Vaill. Bot. Paris. t. 15. f. 6. Dill. Musc. t. 79. f. 1. Fl. Dan. t. 223.

In paludoso ad pagum *Lyugbye* frequens.

Amicissimus *Kanig*, rei herbariæ studiosus indefessus, pascis abhinc annis locum hunc rarissimæ plantæ natalem et textit.

368. ISOËTES. 1184.

848. * ISOËTES *lacustris*.

Dill. Musc. t. 80. f. 2. Fl. Dan. t. 191.

In aquosis sylvæ dictæ *Mutrenalis*.



M U S C I.

369. LYCOPODIUM. 1185.

849. LYCOPODIUM *clavatum* foliis sparsis filamentosis
spicis teretibus pedunculatis geminis.

Fl. Dan. t. 126. Dill. t. 58. f. 1. P. Fl. Oec. 96.

Dan. *Jordmusc*; *Ulve-fod*.

Pharmac. MUSCI CLAVATI herba, femina LYCO-
PODII.

In muscosis sylvæ paa Næsset, ved Huulsoë og
Fuursoë.

850. * LYCOPODIUM *inundatum* foliis sparsis inter-
gerrimis: spica terminali foliosa.

Vaill. Bot. Paris. t. 16. f. 11. Dill. t. 62. f. 7.

In littore Huulsoë.

370. SPHAGNUM. 1187.

851. SPHAGNUM *palustre* ramis deflexis.

Vaill. Bot. Paris. t. 23. f. 3. Dill. Musc. t. 32. f. 1.

Dan. *Rødmoos*.

In paludibus ubique.

371. PHASCUM. 1189.

852. * PHASCUM *acaulon*, acaule, anthera sessili: foliis ovatis acutis conniventibus.

Vaill. Bot. Paris. t. 27. f. 2. Dill. Musc. t. 32. f. 12.

In vivario lateri viæ arenoso, rarum.

853. * PHASCUM *subulatum* acaule, anthera sessili:
foliis subulato - setaceis patulis.

Vaill. Bot. Paris. t. 29. f. 4. Dill. Musc. t. 32. f. 10.

In vivario obtegit latera viarum copiosissime; par-
tius in sylva circa Kollekolle.

854. * PHASCUM *Halleri* acaule, anthera sessili: foliis interioribus mucronatis.

Hall. Helv. Tab. 3. f. 3. Dill. Musc. t. 32. f. 13. Act. Stockholm. 1764. t. . f. .

In vivario in ripa tramites obtegit, in angustis viarum ad lacus *Huul* & *Silleröd* copiosum.

Obs. In hac specie stamen unicum videre licet interiori operculi affixum, parumper extra marginem porrectum, ac calyptram manifestam, quæ acu facillime ab operculo separatur. Vix ac ne vix ratio inveniri potest, cur celeb. *Linneus*, cui observationem ac cespitem plantularum misi, quique in *Actis Soc. Reg.* inferendam curavit, hanc tamen in numerum plantarum in nova *Specierum* editione haud adsciverit. Donetur igitur pulcherrimus advena civitate, ac ornetur nomine Viri in Botanicis ac Anatomicis nunquam non colendi.

372. BUXBAUMIA. 1188.

855. * BUXBAUMIA *aphylla*.

Dill. Musc. t. 68. f. 5. Fl. Dan. t. 44.

Rarissimam plantulam bis legi initio Novembr. 1762. in ripa declivi Nemoris Austrum spectante, ac ultimo Octobris 1764. in opposita ripa. Postea amicissimus *Kanig*, industrius herbarum sectator, in sylva *Friedensburgensi* quædam specimina offendit.

373. FONTINALIS. 1190.

856. FONTINALIS *antipyretica* foliis complicato - carinatis trifariis acutis: antheris lateralibus.

Vaill. Paris. t. 33. f. 5. Dill. Musc. t. 33. f. 1.

Dan. Vand. - Moos.

In amne ad palos.

857. † FONTINALIS *minor* foliis ovato - lanceolatis trifariis acutis, passim geminis: antheris terminalibus. *Dill. Musc. t. 33. f. 2.*

374. POLYTRICHUM. 1192.

858. *POLYTRICHUM commune* caule simplici, anthera parallelepipedæ.

Dill. Musc. t. 54. f. 1. 2. 3. Vaill. Bot. Paris. t. 23. f. 7. 8.

Dan. *Jomfrue-Haar*; *Guld-Haar*.

Pharmac. ADIANTHI AUREI herba.

Ubique. Variat locis uliginosis, pascuis & sterilibus.

859. * *POLYTRICHUM urnigerum* caule ramosissimo; pedunculis axillaribus.

Vaill. Bot. Paris. t. 28. f. 13. Dill. Musc. t. 55. f. 5.

In unica vivarii fossa.

375. MNIMUM. 1193.

860. * *MNIUM pellucidum* caule simplici, foliis ovatis.

Dill. Musc. t. 31. f. 2. Vaill. Bot. Par. t. 24. f. 7.

In pascuis humidis umbrosis.

861. * *MNIUM fontanum* caule simplici geniculis inflexo.

Vaill. Bot. Paris. t. 24. f. 10. Dill. Musc. t. 44. f. 2.

In paludibus passim.

862. * *MNIUM palustre* caule dichotomo, foliis subulatis.

Vaill. Bot. Paris. t. 24. f. 1. Dill. Musc. t. 31. f. 3.

In paludibus.

863. * *MNIUM androgynum* caule ramoso androgynum.

Vaill. Bot. Paris. t. 29. f. 6. Dill. Musc. t. 31. f. 1.

In sylva *Prun-Skov*.

864. * *MNIUM hygrometricum* acaule, anthera nutante: calyptra reflexa tetragona.

Vaill. Bot. Par. t. 26. f. 16. Dill. Musc. t. 52. f. 75.

In præustis vivarii.

865. * *MNIUM purpureum* caule dichotomo, axillis pedunculiferis: anthera erecta: foliis carinatis.
Dill. Musc. t. 49. f. 51.
 In pratis passim.
866. * *MNIUM setaceum* antheris erectis, operculis filiformibus longitudine antherarum.
Dill. Musc. t. 48. f. 44.
 In sepibus.
867. * *MNIUM cirratum* foliis arefactione revolutis.
Vaill. Bot. Paris. t. 24. f. 8. Dill. Musc. t. 48. f. 42.
 In sylva.
868. * *MNIUM annotinum* foliis ovatis acutis pellucidis, pedunculis subradicalibus: anthera nutante.
Dill. Musc. t. 50. f. 68.
 In aquosis.
869. * *MNIUM hornum* antheris pendulis: pedunculo curvato: surculo simplici, foliolis margine scabris.
Vaill. Bot. Paris. t. 24. f. 4. 5. Dill. Musc. t. 51. f. 71.
 Ad radices alnorum frequens.
870. * *MNIUM crudum* antheris pendulis, calyptra recurvata: foliis pellucidis.
Dill. Musc. t. 51. f. 70.
 Ad lacum *Fuursoë*.
871. *MNIUM pyriforme* antheris pendulis turbinatis: stipite filiformi: floribus femineis setiferis.
Dill. Musc. t. 50. f. 60.
 In nemore passim.
872. *MNIUM serpyllifolium* pedunculis aggregatis: foliis patentibus pellucidis.
- a) *MNIUM punctatum* pedunculis aggregatis: foliis obovatis integerrimis obtusis punctatis.
Dill. Musc. t. 53. f. 81. Vaill. Bot. Paris. t. 26. f. 5.
- β) *MNIUM cuspidatum* pedunculis aggregatis: foliis alternis acutis serratis.
Vaill. Bot. Paris. t. 26. f. 18. Dill. Musc. t. 53. f. 79.
- γ) *MNIUM proliferum* pedunculis aggregatis: foliis roseo-congestis, lanceolatis, acutis. *Dill. Musc. t. 52. f. 77.*

72) *MNIUM undulatum* pedunculis aggregatis: foliis oblongis undulatis.

Vaill. Bot. Paris. t. 24. f. 3. Dill. Musc. t. 52. f. 76.

In humidis ubique.

73. * *MNIUM polytrichoides* calyptra villosa.

Dill. Musc. t. 55. f. 6. 7. Vaill. Bot. Paris. t. 26. f. 15. t. 29. f. 11.

In præruptis fossarum.

74. * *MNIUM trichomanis* foliis distichis integerremis. *Dill. Musc. t. 31. f. 5.*

In udis.

75. * *MNIUM fissum* foliis distichis bifidis.

Dill. Musc. t. 31. f. 6.

In declivibus sylvestribus frequens.

76. * *MNIUM jungermannia* foliis distichis, pinnis subtus auriculatis. *Dill. Musc. t. 69. f. 1.*

In udis.

Feminas solas vidi, mense Octobri, Mares ultimo Aprilis florentes.

376. B R Y U M. 1194.

* *Antheris sessilibus.*

77. * *BRYUM apocarpum* antheris sessilibus terminalibus: calyptra minima.

Vaill. Bot. Paris. t. 21. f. 15. t. 27. f. 8. Dill. Musc. t. 32. f. 4. 5.

In arboribus & saxis frequens.

78. * *BRYUM striatum* antheris subsessilibus sparsis: calyptris striatis sursumve pilosis.

Dill. Musc. t. 55. f. 8. 9. 10. 11. Vaill. Bot. Paris. t. 25. f. 5. 6. t. 27. f. 9. 10.

In arboribus & saxis frequens.

* * *Antheris pedunculatis erectis.*

879. * *BRYUM pomiforme* antheris erectis sphaericis
Vaill. Bot. Paris. t. 24. f. 9. 12. Dill. Musc. t. 44. f. 1.

In præruptis montium ac viarum sylvestrium ubique.

880. * *BRYUM pyriforme* antheris erectis obovatis,
 calyptra subulata: furculis acaulibus, foliis ovatis
 muticis.

Vaill. Bot. Paris. t. 29. f. 3. Dill. Musc. t. 44. f. 6.

In cespitibus rarius.

881. * *BRYUM extinctorium* anthera erecta oblonga
 minore calyptris, laxis æqualibus.

Vaill. Bot. Paris. t. 26. f. 1. Dill. Musc. t. 45. f. 8. 9.

• In cespitibus aggerum passim.

882. * *BRYUM subulatum* antheris erectis subulatis:
 furculis acaulibus.

Vaill. Bot. Paris. t. 25. f. 8. Dill. Musc. t. 45. f. 10.

Ibidem.

883. * *BRYUM rurale* antheris erectiusculis: foliis
 liferis recurvis.

Vaill. Bot. Paris. t. 25. f. 3. Dill. Musc. t. 45. f. 12.

In tectis molendini.

884. * *BRYUM murale* antheris erectis: foliis piliferis
 rectiusculis, furculis simplicibus cespitosis.

Vaill. Bot. Paris. t. 24. f. 14. 15. & t. 25. f. 4.

Dill. Musc. t. 45. f. 14.

In muris.

885. * *BRYUM scoparium* antheris erectiusculis: pedun-
 culis aggregatis, foliis secundo recurvatis, caule
 declinato.

Vaill. Bot. Paris. t. 28. f. 12. Dill. Musc. t. 46. f. 6.

In sylvis frequens.

886. * *BRYUM undulatum* antheris erectiusculis: pedunculis subfolitariis, foliis lanceolatis carinatis undulatis patentibus ferratis.
Vaill. Bot. Paris. t. 26. f. 17. Dill. Musc. t. 46. f. 18.
887. * *BRYUM pellucidum* antheris erectiusculis: foliis recurvis acutis, caule hirsuto.
Dill. Musc. t. 46. f. 23. 24.
 In palude.
888. * *BRYUM flexuosum* antheris erectis: foliis fetaceis, pedunculis flexuosis. *Dill. Musc. t. 47. f. 33.*
 In sylva.
889. * *BRYUM heteromallum* antheris erectis: foliis fetaceis secundis.
Vaill. Bot. Paris. t. 27. f. 7. Dill. Musc. t. 47. f. 37.
 In arenosis viarum lateribus.
890. * *BRYUM truncatulum* antheris erectis subrotundis: operculo mucronato.
Vaill. Bot. Paris. t. 26. f. 2. Dill. Musc. t. 45. f. 7.
 In sepibus.
891. * *BRYUM viridulum* antheris erectis ovatis: foliis lanceolatis acuminatis imbricato-patulis.
Vaill. Bot. Paris. t. 29. f. 5. Dill. Musc. t. 48. f. 43.
 In agris.
892. * *BRYUM hypnoides* antheris erectis: caule erectiusculo: ramis lateralibus brevibus fertilibus,
Dill. Musc. t. 47. f. 27. 28.
 In saxis.
893. * *BRYUM paludosum* acaulon foliis fetaceis: antheris obtusissimis patulis. *Dill. Musc. t. 49. f. 53.*
 In humidis littoris elatioris.
894. *BRYUM verticillatum* pedunculis siccitate contortis: foliis pñiferis: furculis fastigiatis.
Dill. Musc. t. 47. f. 35.
 In declivi fossarum.

895. * *BRYUM squarrosum* antheris obliquis : foliis quinquefariam imbricatis recurvis : caule ferrugineo-tomentoso. *Buxb. Cent. 4. t. 65. f. 1.*
In paludibus apricis.
896. *BRYUM argenteum* antheris pendulis : furculis cylindricis imbricato-lævibus.
Dill. Musc. t. 50. f. 62. Vaill. Bot. Paris. t. 26. f. 3.
Dan. Flöyel - Moos.
In faxis & tectis passim.
897. * *BRYUM pulvinatum* antheris subrotundis : pedunculis reflexis, foliis piliferis.
Vaill. Bot. Paris. t. 6. f. 21. Dill. Musc. t. 50. f. 65.
In faxis & muris,
898. * *BRYUM caespiticiu*m antheris pendulis : foliis lanceolatis acuminato-setaceis : pedunculis longissimis.
Vaill. Bot. Paris. t. 29. f. 7. Dill. Musc. t. 50. f. 66.
In tectis & glareosis.
899. * *BRYUM capillare* antheris pendulis : foliis ovatis fetiferis carinatis : pedunculis longissimis.
Dill. Musc. t. 50. f. 67.
In glareosis.
900. * *BRYUM carneu*m antheris pendulis subglobosis : foliis acutis alternis. *Dill. Musc. t. 50. f. 69.*
In humidis.

377. H Y P N U M. 1195.

* Frondibus pinnatis.

901. * *HYPNUM bryoïdes* fronde simplicissima pinnata lanceolata, apice pedunculifera.
Vaill. Bot. Paris. t. 24. f. 13. Dill. Musc. t. 34. f. 1.
In littore stagni umbroso.

12. * *HYPNUM taxifolium* fronde simplicissima pinnata lanceolata basi pedunculifera.

Vaill. Bot. Paris. t. 24. f. 11. Dill. Musc. t. 34. f. 2.

In umbrosis. In fossis horti *Friderichsbergensis* copiose.

13. * *HYPNUM denticulatum* fronde pinnata simplici: pinnis duplicatis basi pedunculifera.

Vaill. Bot. Paris. t. 29. f. 8. Dill. Musc. t. 34. f. 5.

In umbroso sylvæ.

14. *HYPNUM complanatum* fronde pinnata ramosa: foliolis imbricatis acutis complicatis compressis.

Vaill. Bot. Paris. t. 23. f. 4. Dill. Musc. t. 34. f. 7.

Dan. *Bregne-Moos.*

In truncis arborum frequens.

* * Surculis vagis.

15. * *HYPNUM lucens* furculis ramosis, frondibus subpinnatis: foliolis punctatis.

Dill. Musc. t. 34. f. 10.

In ripa *Paa-Nasset* & alibi.

16. * *HYPNUM triquetrum* ramis vagis recurvis: foliis ovatis recurvatis patulis.

Vaill. Bot. Paris. t. 28. f. 9. Dill. Musc. t. 38. f. 28.

In sylva passim.

* * * Surculis pennatis.

17. * *HYPNUM filicinum* furculis pennatis, ramulis distantibus, foliolis imbricatis incurvis acutis secundis.

Vaill. Paris. t. 29. f. 9. Dill. Musc. t. 36. f. 19.

In pratis subhumidis.

18. *HYPNUM proliferum* furculis proliferis planopinnatis: pedunculis aggregatis.

Vaill. Bot. Paris. t. 25. f. 1. Dill. Musc. t. 35. f. 14.

Dan. *Steenbregne-Moos.*

In sterilibus frequens.

909. * HYPNUM *delicatulum* furculis subproliferis planis - pinnatis cuspidatis: pedunculis aggregatis.

Dill. Musc. t. 83. f. 6.

In sterilibus siccis.

910. * HYPNUM *parietinum* furculis plano-pinnatis continuatis: pedunculis aggregatis.

Dill. Musc. t. 35. f. 13.

Ligna passim obvestit.

911. * HYPNUM *prælongum* furculis subbipinnatis ramulis remotis: foliolis ovatis: antheris cernuis.

Dill. Musc. t. 35. f. 15.

In truncis & faxis passim.

912. * HYPNUM *crista castrensis* furculis pinnatis: ramulis approximatis: apicibus recurvis.

Vaill. Bot. Paris. t. 27. f. 14. Dill. Musc. t. 36. f. 20.

913. * HYPNUM *abietinum* furculis pinnatis teretibus furculis remotis inæqualibus.

Dill. Musc. t. 35. f. 17. Vaill. Bot. Paris. t. 29. f. 12.

In sylvis passim.

914. * HYPNUM *plumosum* furculis pinnatis repentibus, ramis confertis: foliis imbricatis subulatis, antheris erectis.

Dill. Musc. t. 35. f. 16.

In terra in vivario.

* * * * Foliis reflexis.

915. * HYPNUM *cupressiforme* furculo subpinnato: foliis secundis recurvis apice subulatis.

Vaill. Bot. Paris. t. 27. f. 13. Dill. Musc. t. 37. f. 23.

In sylvis ad radices arborum & in faxis.

916. * HYPNUM *aduncum* furculo erectiusculo subramoso: foliis secundis recurvatis subulatis: ramis recurvatis.

Dill. Musc. t. 37. f. 26.

In uliginosis.

17. * *HYPNUM squarrosum* ramis vagis, foliis ovatis apice subulatis, undique reflexis.

Dill. Musc. t. 39. f. 38. 39.

In subhumidis horti.

18. * *HYPNUM palustre* furculis repentibus: ramis confertis erectis: foliolis ovatis secundis: antheris erectiusculis. *Dill. Musc. t. 37. f. 27.*

In udis littoris stagni *Fuursoë*.

***** Surculis fasciculatis.

19. * *HYPNUM dendroïdes* furculo erecto: ramis fasciculatis terminalibus simpliciusculis: antheris erectis.

Vaill. Bot. Paris. t. 26. f. 6. Dill. Musc. t. 40. f. 48.

In ripa hortensj.

20. * *HYPNUM alopecurum* furculo erecto: ramis fasciculatis terminalibus subdivisis: antheris subnulantibus.

Vaill. Bot. Paris. t. 23. f. 5. Dill. Musc. t. 41. f. 49.

In sylvaticis subhumidis.

***** Surculis teretiusculis.

21. * *HYPNUM purum* furculis pinnato-sparsis subulatis: foliis ovatis obtusis conniventibus.

Vaill. Bot. Paris. t. 28. f. 3. Dill. Musc. t. 40. f. 45.

In pascuis sylvaticis.

22. * *HYPNUM cuspidatum* furculis vagis: apice foliis convolutis acuminatis. *Dill. Musc. t. 39. f. 34.*

In paludosis.

***** Surculis confertis.

23. * *HYPNUM sericeum* furculo repente: ramis confertis erectis: foliis subulatis: antheris erectis.

Vaill. Bot. Paris. t. 27. f. 3. Dill. Musc. t. 42. f. 59.

In truncis.

924. * *HYPNUM velutinum* furculo repente: ramis confertis erectis: foliis subulatis: antheris subulatis. *Vaill. Bot. Paris. t. 26. f. 9.*

925. * *HYPNUM serpens* furculis repentibus: ramis filiformibus: foliis oblitteratis.

Vaill. Bot. Paris. t. 28. f. 2. 6. 7. 8. Dill. Musc. t. 42. f. 64.

In faxis.

926. * *HYPNUM sciuroïdes* furculis erectis ramosis in curvatis. *Dill. Musc. t. 41. f. 54.*

In truncis arborum passim; in nemore & alibi.

A L G Æ.

378. JUNGERMANNIA. 1196.

* Frondibus pinnatis secundis.

927. * *JUNGERMANNIA viticulosa*, frondibus simpliciter pinnatis: foliolis subulatis.

Dill. Musc. t. 69. f. 7. 8.

In ripa umbrosa.

Obs. Quam ego cum fructif. inveni, magis refert f. 2. *Dill.* quoad rotunditatem foliorum, eorumque distans, quam f. 7.

928. * *JUNGERMANNIA lanceolata* frondibus simpliciter pinnatis, lanceolatis, apice floriferis: foliolis integerrimis. *Dill. Musc. t. 70. f. 10.*

In ripa stagni *Humföe*.

929. * *JUNGERMANNIA bidentata* frondibus simpliciter pinnatis: apice floriferis: foliolis bidentatis.

Vaill. Bot. Paris. t. 19. f. 8. Dill. Musc. t. 70. f. 11.

In umbroso sylvæ.

930. * *JUNGERMANNIA bisuspidata* frondibus simpliciter pinnatis, medio floriferis: foliolis bidentatis. *Dill. Musc. t. 70. f. 12.*

Ad radices alni passim.

* *Frondebis pinnatis: pinnulis auriculatis superne tectis.*

31. * *JUNGERMANNIA nemorosa* frondibus supra bipinnatis, apice floriferis: foliolis ciliatis.

Dill. Musc. t. 71. f. 18.

In sylva, sed raro.

32. * *JUNGERMANNIA reptans* frondibus subtus bipinnatis apice reptantibus: foliolis quadridentatis.

Dill. Musc. t. 71. f. 24.

In semetis umbrosis passim.

*** *Frondebis imbricatis.*

33. * *JUNGERMANNIA complanata* furculis repentibus: foliolis inferne auriculatis duplicato-imbricatis: ramis æqualibus. *Dill. Musc. t. 72. f. 26.*

In cortice fagorum frequens.

34. * *JUNGERMANNIA dilatata* furculis repentibus: foliis inferne auriculatis duplicato-imbricatis: ramis apice latioribus.

Vaill. Bot. Paris. t. 19. f. 10. Dill. Musc. t. 72. f. 27.

In cortice arborum passim.

35. * *JUNGERMANNIA tamarisci* foliis imbricatis serie duplici: superioribus subrotundis convexis, obtusis, quadruplo majoribus.

Vaill. Bot. Paris. t. 23. f. 10. Dill. Musc. t. 72. f. 31.

In faxis.

36. * *JUNGERMANNIA platyphylla* furculis procumbentibus, subtus imbricatis: foliolis cordatis acutis.

Vaill. Bot. Paris. t. 19. f. 9. Dill. Musc. t. 72. f. 32.

In truncis fagorum frequens.

37. * *JUNGERMANNIA varia* furculis erectiusculis bifariam imbricatis: foliolis bipartitis.

Dill. Musc. t. 73. f. 36.

In humidiusculis.

**** Surculis undique imbricatis, foliolis sparsis.

938. * JUNGERMANNIA *trichophylla* furculis teretibus: foliolis capillaceis æqualibus.

Dill. Musc. t. 73. f. 37.

Inter alias, sed rara.

***** *Acaules*, frondibus simplicibus.

939. * JUNGERMANNIA *epiphylla* acaulis, foliolo frondi innato. *Dill. Musc. t. 74. f. 41.*

In ripis elatioribus æque ac depressis frequentissima

940. * JUNGERMANNIA *pinguis* acaulis fronde oblonga sinuata pingui. *Dill. Musc. t. 74. f. 42.*

Rarius in palude.

941. * JUNGERMANNIA *furcata* acaulis fronde linearum ramosa, extremitatibus furcatis obtusiusculis.

Vaill. Bot. Paris. t. 23. f. 11. Dill. Musc. t. 74. f. 45. 47.

In truncis ac saxis frequens. *f. Dill. 47. in paludosis*

379. MARCHANTIA. 1198.

942. MARCHANTIA *polymorpha* calyce communi deconfido.

Dill. Musc. t. 76. f. 6. Mich. Gen. t. 1. f. 1.

Dan. Lever - Urt.

Pharmac. HEPATICÆ FONTANÆ herba.

In luco; in palude prope amnem & alibi. Totali fossam apud Nyebroë implet.

- * β) MARCHANTIA *stellata*.

Act. Paris. 1713. t. 5. Mich. Gen. t. 1. f. 2. Dill. Musc. t. 77. f. 7.

In paludosis ad lacum *Bagsoerd* ac promontorii *Mou* frequens.

- γ) MARCHANTIA capitulo non dissecto.

Mich. Gen. t. 1. f. 5.

In præstis locis.

943. * M

43. * *MARCHANTIA conica* calyce communi quinqueloculari subovato. *Dill. Musc. t. 75. f. 1.*
In ripa elatiori stagni *Fuursøe* radices alni obvestit.

380. *BLASIA*. 1199.

44. * *BLASIA pusilla*.
Dill. Musc. t. 31. f. 7. Fl. Dan. t. 45.
In elatiori ripa lacus *Fuursøe* ac in fossa ad tugurium *Bakkehus* copiose.

381. *RICCIA*. 1200.

45. * *RICCIA minima* frondibus glabris bipartitis acutis. *Dill. Musc. t. 78. f. 11.*
In inundatis insulæ *Kaaberholm*.
46. * *RICCIA crystallina* frondibus superficie papillofis. *Dill. Musc. t. 78. f. 12. Vaill. Bot. Paris. t. 19. f. 2.*
In vado prope Palatium.
47. * *RICCIA fluitans* frondibus dichotomis lineatifiliformibus.
Vaill. Bot. Paris. t. 10. f. 3. Dill. Musc. t. 74. f. 47.
In vado sylvestri.
Obs. Plantam figuræ *Dill.* simillimam initio veris cum folliculo lateri inferiori insidente, globuloque nigro *JUNGERMANNII* invenisse memini, hinc varietatem furcatæ crederem; jam vero, cum nec ad manus sit, nec de mente *Linnei* satis certo esse liceat, qui dubium, quod de ejus genere in *Fl. Suec.* prodidit, in edit. alt. *Specier.* in medio reliquit, hic posteris dijudicandam appono.

382. *LICHEN*. 1202.

* *Leprosi tuberculati.*

748. * *LICHEN scriptus* leprosus albicans: lineolis nigris ramosis characteriformibus.
Dill. Musc. t. 18. f. 1.
In cortice arborum.

949. * *LICHEN geographicus* leprosus flavescens: L
neolis nigris mappam referens.

Dill. Musc. t. 18. f. 5.

In faxis.

950. * *LICHEN rugosus* leprosus albicans, lineolis sum
plicibus punctisque nigris confertis.

Dill. Musc. t. 18. f. 2.

In truncis fagorum.

951. * *LICHEN sanguinarius* leprosus cinereo-vire
scens: tuberculis atris.

Dill. Musc. t. 18. f. 3.

In cortice arborum ac lapidibus.

952. * *LICHEN fusco-ater* leprosus fuscus, tuberculis
atris.

In faxis.

953. * *LICHEN calcareus* leprosus candidus, tubercu
lis atris. *Dill. Musc. t. 18. f. 8.*

In lapidibus.

954. * *LICHEN fagineus* leprosus albus, tuberculis
albis farinaceis.

In cortice fagi.

955. * *LICHEN carpinus* leprosus cinereus, tubercu
lis albidis rugosis.

In truncis.

956. * *LICHEN ericetorum* leprosus candidus, tubercu
culis incarnatis. *Dill. Musc. t. 14. f. 1.*

In glareosis.

* * Leprosi *scutellati.*

957. * *LICHEN candelarius* crustaceus flavus, scutell
lis luteis.

In muris & parietibus.

958. * *LICHEN pallescens* crustaceus albicans, scu
tellis pallidis.

Dill. Musc. t. 18. f. 17.

In cortice arborum.

959. * LICHEN *tartareus* crustaceus ex albedo virefcens, scutellis flavescentibus: margine albo.

Dill. Musc. t. 18. f. 12.

In faxis.

960. * LICHEN *subfuscus* crustaceus albicans, scutellis subfufcis: margine cinereo subcrenato.

Dill. Musc. t. 18. f. 16.

* * * *Imbricati.*

961. * LICHEN *centrifugus* imbricatus, foliolis multifidis obsolete lævibus albidis centrifugis: scutellis rufo-fufcis. *Dill. Musc. t. 24. f. 75.*

In faxis.

962. * LICHEN *saxatilis* imbricatus, foliolis sinuatis scabris lacunosis: scutellis folio concoloribus.

Dill. Musc. t. 24. f. 83.

In faxis frequens ac in cortice fagi.

963. * LICHEN *omphalodes* imbricatus: foliolis multifidis glabris obtusis canis: punctis vagis eminentibus.

Vaill. Bot. Paris. t. 20. f. 10. Dill. Musc. t. 24. f. 8.

Pharmac. USNEA.

In maceriebus.

964. * LICHEN *olivaceus* imbricatus, foliolis lobatis nitidis lividis.

Vaill. Bot. Paris. t. 20. f. 8. Dill. Musc. t. 24. f. 78.

In lapidibus.

965. LICHEN *parietinus* imbricatus, foliis crispis fulvis, peltis concoloribus fulvis.

Dill. Musc. t. 24. f. 76.

Dan. *Gnul-Skorpe-Moos.*

In lignis.

966. * LICHEN *physodes* imbricatus: laciniis obtusis subinflatis. *Dill. Musc. t. 20. f. 49.*

In truncis ac faxis.

967. * LICHEN *stellaris* imbricatus, foliolis oblongis laciniatis angustis cinereis, scutellis pullis.

Dill. Musc. t. 24. f. 70.

In ramis arborum vulgaris.

**** *Foliacei.*

968. * LICHEN *ciliaris* foliaceus erectiusculus: laciniis linearibus ciliatis, scutellis pedunculatis crenatis.

Vaill. Bot. Paris. t. 20. f. 4. Dill. Musc. t. 20. f. 45.

In arboribus.

969. * LICHEN *Islandicus* foliaceus adscendens laciniatus, marginibus elevatis ciliatis.

970. LICHEN *pulmonarius* foliaceus laciniatus obtusus glaber: supra lacunosus, subtus tomentosus.

Dill. Musc. t. 29. f. 113.

Dan. *Lunge-Moos.*

Pharmac. PULMONARIÆ ARBOREÆ herba.

In fagis antiquis.

971. * LICHEN *furfuraceus* foliaceus decumbens furfuraceus: laciniis acutis: subtus lacunosus atris

Dill. Musc. t. 21. f. 52.

In arboribus.

972. * LICHEN *farinaceus* foliaceus erectus compressus ramosus farinaceus: lateribus verrucosis.

Vaill. Bot. Paris. t. 20. f. 14. 15. Dill. Musc. t. 23. f. 63.

In arboribus.

973. * LICHEN *calicaris* foliaceus erectus linearis ramosus lacunosus convexus mucronatus.

Vaill. Bot. Paris. t. 20. f. 6. Dill. Musc. t. 23. f. 62.

In antiquis fagis.

974. * LICHEN *fraxineus* foliaceus erectus oblongus lanceolatus sublaciniatus lacunosus glaber, scutellis subpedunculatis. *Dill. Musc. t. 22. f. 59.*

In arboribus.

975. * LICHEN *prunastri* foliaceus erectiusculus lacunosus: subtus tomentosus albus.

Vaill. Bot. Paris. t. 20. f. 11. 12. Dill. Musc. t. 21. f. 55.

In arboribus.

976. * LICHEN *juniperinus* foliaceus laciniatus crispus fulvus, peltis lividis. *Dill. Musc. t. 24. f. 76.*

In arboribus.

977. * LICHEN *caperatus* pallide viridis rugosus: margine undulatus. *Dill. Musc. t. 25. f. 97.*

In faxis.

***** *Coriacei.*

978. * LICHEN *resupinatus* coriaceus repens lobatus, peltis marginalibus posticis.

Dill. Musc. t. 28. f. 105.

In sylva.

979. * LICHEN *venosus* coriaceus repens ovatus planus: subtus venosus villosus, peltis marginalibus horizontalibus. *Dill. Musc. t. 28. f. 109.*

In sylva.

980. * LICHEN *apthosus* coriaceus repens lobatus obtusus planus, verrucis sparsis, pelta marginali adscendente. *Dill. Musc. t. 28. f. 106.*

Pharmac. MUSCUS CUMATILIS.

In sylva.

981. * LICHEN *arcticus* coriaceus repens lobatus obtusus planus levis: subtus avenius villosus.

In terra.

982. LICHEN *caninus* coriaceus repens lobatus obtusus planus: subtus venosus villosus, pelta marginali adscendente.

Vaill. Bot. Par. t. 21. f. 16. Dill. Musc. t. 27. f. 102.

Pharmac. MUSCI CANINI herba.

In sylvis ubique.

***** *Scyphiferi.*

983. * *LICHEN cocciferus* scyphifer simplex integerrimus, stipite cylindrico, tuberculis coccineis.
Dill. Musc. t. 14. f. 7. Vaill. Bot. Paris. t. 21. f. 4.
Pharmac. MUSCI PYXIDATI herba.
 In ericetis ad lacum *Bagsværd*.
984. *LICHEN pyxidatus* scyphifer simplex crenulatus, tuberculis fuscis.
Dill. Musc. t. 14. f. 6. Vaill. Bot. Paris. t. 21. f. 5.
 8. 9. 11.
 Dan. *Begger - Moos.*
 In sylvis ubique.
985. * *LICHEN fimbriatus* scyphifer simplex denticulatus, stipite cylindrico.
Dill. Musc. t. 14. f. 8. Vaill. Bot. Paris. t. 21. f. 6.
 In sylvis.
986. * *LICHEN digitatus* subscyphifer ramosissimus, ramis cylindricis: calycibus integris.
Dill. Musc. t. 15. f. 12.
 In sylvis sterilibus.
987. * *LICHEN cornutus* scyphifer simpliciusculus subventricosus, calycibus integris.
Dill. Musc. t. 15. f. 14.
 In ericetis.

***** *Fruticulosi.*

988. *LICHEN rangiferinus* fruticulosus perforatus ramosissimus, ramulis nutantibus.
Dill. Musc. t. 16. f. 29. 30. Fl. Dan. t. 180.
 Dan. *Rensdyr - Meos.*
 In Insula *Kaaberholm*.
989. * *LICHEN uncialis* fruticulosus perforatus: ramulis brevissimis acutis. *Dill. Musc. t. 16. f. 22.*
 In campo ericetoso versus *Dronninggaard*.

90. * *LICHEN subulatus* fruticulofus dichotomus, ramis simplicibus subulatis. *Dill. Musc. t. 16. f. 26.*
In sylvis.
91. * *LICHEN paschalis* fruticulofus solidus, tectus foliolis crustaceis. *Dill. Musc. t. 17. f. 33.*
In macerie.

***** *Filamentosi.*

92. *LICHEN barbatus* filamentofus pendulus subarticulatus, ramis patentissimis. *Dill. Musc. t. 12. f. 6.*
In fagis.
93. * *LICHEN chalybeiformis* filamentofus subramofus decumbens implicate - flexuosus.
Dill. Musc. t. 13. f. 10.
In trunco antiquo quercus.
94. *LICHEN hirtus* filamentofus ramosissimus erectus, tuberculis farinaceis sparsis. *Dill. Musc. t. 13. f. 12.*
In arboribus & sepibus.
95. † *LICHEN floridus* filamentofus ramosus erectus: scutellis radiatis. *Dill. Musc. t. 13. f. 13.*
In fagetis.

383. *CHARA*. 1203.

96. *CHARA vulgaris* caulibus lævibus: frondibus interdentatis. *Vaill. in Act. Paris. 1719. t. 3. f. 1.*
In lacu *Lyngbye* & in fossis.
97. *CHARA hispida* aculeis caulinis capillaribus confertis. *Vaill. in Act. Paris. 1719. t. 3. f. 3.*
El. Dan. t. 154.
In paludibus turfosis.

384. *TREMELLA*. 1204.

98. * *TREMELLA juniperina* sessilis membranacea, auriformis fulva. *Schæff. Fung. Icon. t. 168.*
In ramalibus sepium.

999. TREMELLA *Nostoc* plicata undulata.

Dill. Musc. t. 10. f. 14.

Dan. *Trold - Smør.*

In terra arenosa horti ac in ripa elatiori lacus *Fuursoë* frequens.

1000. TREMELLA *auricula* sessilis membranacea auriformis cinerea. *Mich. Gen. t. 66. f. 1.*

Pharmac. FUNGUS SAMBUCINUS; AURICULA JUDE.

In Sambuco arida.

1001. * TREMELLA *Lichenoides* foliis erectis planis margine crispo lacinulato. *Dill. Musc. t. 19. f. 31.*

In ripa elatiori *Fuursoë* inter muscos ac in saxo in vivario.

1002. * TREMELLA *verrucosa* tuberculosa solidam-gosa. *Dill. Musc. t. 10. f. 6.*

In lapidibus lacus *Fuursoë*.

1003. * TREMELLA *hemisphærica*, hemisphærica sparsa.

In ripa *Fuursoë*.

1004. * TREMELLA *purpurea* subglobosa sessilis sœtaria, glabra. *Dill. Musc. t. 18. f. 6.*

In ramis arborum exaridis.

385. F U C U S. 1205.

* *Dichotomi frondescentes.*

1005. † FUCUS *ferratus* fronde plana dichotoma serrata ad apices tuberculata. *Morif. t. 9. f. 1.*

1006. † FUCUS *vesiculosus* fronde dichotoma integra: caule medium folium transcurrente: vesiculis verrucosis terminalibus. *P. Fl. Qec. 73.*

Dan. *Strand - Klevet.*

1007. † FUCUS *ceranoides* fronde dichotoma plana integra: apicibus bifidis vesiculosis.

Morif. t. 8. f. 13.

1008. † FUCUS *inflatus* fronde bifida: laciniis ovato-lanceolatis inflatis, apice divisis.

* * Dichotomi caulescentes.

1009. FUCUS *nodosus* caule compresso dichotomo: medio ramorum in vesiculam dilatato.

Fl. Dan. t. 146.

In mari Balthico.

1010. FUCUS *filiquosus* caule tereti ramosissimo: pedunculis alternis: vesiculis oblongis acuminatis.

Fl. Dan. t. 106.

In mari Balthico.

* * * Fructificationibus non vesicariis.

1011. FUCUS *filum* filiformis simplex fragilis opacus.

Aët. Hafn. 1671. f. 8.

In mari Balthico; frequens ad Insulam *Thorsholm*.

1012. † FUCUS *fastigiatus* dichotomus ramosissimus teres, uniformis, fastigiatus. Moris. t. 9. f. 9.

In littore maris Balthici sub promontorio Insulae *Mona* legi.

386. U L V A. 1206.

1013. * ULVA *intestinalis* tubulosa simplex.

Dill. Musc. t. 9. f. 7.

In mari Balthico.

1014. * ULVA *granulata* sphaerica aggregata.

Dill. Musc. t. 10. f. 17.

In ripa amnis.

1015. † ULVA *labyrinthiformis* (Lactuca Fl. Succ.) palmata prolifera membranacea: ramentis inferne angustatis. Dill. Musc. t. 8. f. 1.

387. CONFERRA. 1207.

* Filamentis *simplicibus*, *æqualibus*, *geniculis destitutis*.

1016. * CONFERRA *rivularis* filamentis simplicissimis æqualibus longissimis. *Dill. Musc. t. 2. f. 1.*
In amne.

1017. * CONFERRA *fontinalis* filamentis simplicissimis æqualibus digito brevioribus. *Dill. Musc. t. 2. f. 3.*
In fontibus.

* * Filamentis *ramosis æqualibus*.

1018. * CONFERRA *bullosa* filamentis æqualibus ramosis, bullas aëreas includens.
Dill. Musc. t. 3. f. 11.
In aquis stagnantibus.

* * * Filamentis *anastomosantibus*.

1019. † CONFERRA *reticulata* filamentis reticulatim coadunatis.

* * * * Filamentis *geniculatis*.

1020. * CONFERRA *capillaris* filis geniculatis simplicibus: articulis alternatim compressis.
Dill. Musc. t. 5. f. 25.
In fossa sylvæ.

1021. † CONFERRA *corallinoides* filis geniculatis dichotomis. *Dill. Musc. t. 6. f. 36. 37.*
In mari Balthico sub promontorio Infulæ *Mona*.

1022. † CONFERRA *agagropila* filamentis geniculatis ramosissimis e centro confertissimis, globum constituentibus.

388. BYSSUS. 1208.

* *Filamentosa.*

023. * BYSSUS *flos aqua* filamentis plumosis nantibus.

In aquis.

024. * BYSSUS *phosphorea* lanuginosa violacea lignis adnascens. *Dill. Musc. t. 1. f. 6.*

In trabe antiqua.

025. * BYSSUS *velutina* filamentosa viridis.

Dill. Musc. t. 1. f. 14.

In terra in nemore prope hortum.

026. * BYSSUS *aurea* capillacea, pulverulenta: fructificationibus sparsis: filamentis simplicibus ramosisque. *Dill. Musc. t. 1. f. 16.*

In saxo e ripa altiori stagni *Fuurfsoë* eminente.

* * *Pulverulenta.*

027. * BYSSUS *saxatilis* pulverulenta cinerea rupes operiens.

In faxis.

028. * BYSSUS *candelaris* pulverulenta flava lignis adnascens.

Dill. Musc. t. 1. f. 4.

In tectis & parietibus ligneis.

029. * BYSSUS *botryoïdes* pulverulenta viridis.

Dill. Musc. t. 1. f. 5.

In terra diu humida.

030. * BYSSUS *incana* pulverulenta incana, farinæ instar strata. *Dill. Musc. t. 1. f. 3.*

Ad latera fossarum.

031. * BYSSUS *lactea* pulverulenta crustacea albissima,

Dill. Musc. t. 1. f. 2.

In muscis.

389. SPONGIA. 1072. *Edio. V. Gener. Plant.*

1032. * SPONGIA *fluviatilis* erecta fragilis: ramis retiusculis.

In amne.

Hoc Genus in nova *Specierum ac Generum* editione plane omisit *System. Sexualis Auctor*, rationem omissionis haud indicans; sine dubio quod ad ZOOPHYTA ergo ad animale regnum transferendum putat.

F U N G I.

390. A G A R I C U S. 1209.

* *Stipitati pileo orbiculato.*

1033. * AGARICUS *umbelliferus* stipitatus, pileo plicato membranaceo, lamellis basi latioribus.

Mich. Gen. t. 80. f. 11. Buxb. Cent. IV. t. 32. f. 3.

In foliis faginis semiputridis.

1034. * AGARICUS *androsaceus* stipitatus albus, pileo plicato membranaceo, stipite nigro.

Vuill. Bot. Paris. t. 11. f. 21. 22. 23.

1035. * AGARICUS *piperatus* stipitatus, pileo planiusculo lactescente, margine deflexo, lamellis innato - pallidis.

In sylvis.

1036. * AGARICUS *exstinctorius* stipitatus, pileo campaniformi albido lacero, lamellis niveis, stipite subbulbofo subulato nudo.

In fimetis.

1037. * AGARICUS *campestris* stipitatus, pileo convexo squamato albido, lamellis rufis.

Sterb. Fung. t. 1. f. 6. D.

Dan. *Skampion.*

In pratis frequens; passim in sylvis. Hunc lippis & tonforibus notum, cur Nostrates Floris suis excluderint, vix satis constat.

038. * *AGARICUS cinnamomens stipitatus*, pileo fordide flavo, lamellis luteo-rufis.

In luco.

039. *AGARICUS chantarellus stipitatus*, lamellis ramosis decurrentibus.

Vaill. Bot. Par. t. II. f. 14. 15.

In sylvestribus gramineis.

040. *AGARICUS Georgii stipitatus*, pileo flavo convexo, lamellis albis.

In sylvis raro.

041. * *AGARICUS clavus stipitatus*, pileo luteo convexo striato, lamellis stipiteque albis.

Vaill. Bot. Paris. t. II. f. 19. 20.

In luco raro.

042. * *AGARICUS campanulatus stipitatus*, pileo campanulato striato pellucido, lamellis adscendenti-
bus, stipite nudo. *Vaill. Bot. Paris. t. 12. f. 1. 2.*

In pratis.

043. * *AGARICUS fimetarius stipitatus*, pileo campanulato lacero, lamellis nigris lateraliter flexuosis, stipite fistuloso.

Sterb. Fung. t. 23. f. k. Buxb. Cent. IV. t. 27. f. 1.

Schaff. Fung. Bavar. t. 7. S. 8.

In fimetis, gramineis, quid quod in *Xylis* sabulosis frequens.

044. * *AGARICUS clypeatus stipitatus*, pileo hemisphærico fordido: umbilico prominente. lamellis albis, stipite longo cylindraceo albo.

In sylvis.

045. * *AGARICUS mammosus stipitatus*, pileo convexo acuminato griseo, lamellis convexis griseis crenatis, stipite nudo.

In sylvis.

046. * *AGARICUS equestris stipitatus*, pileo pallido disco luteo, lamellis sulphureis.

In arvis.

1047. * AGARICUS *violaceus* stipitatus, pilei marginē violaceo tomentoso, stipite cærulescente, lamina ferruginea.

Michx. Gen. t. 74. f. 1. Buxb. Cent. IV. t. 7. f. 2.
In nemore frequens.

1048. * AGARICUS *integer* stipitatus, lamellis omnibus magnitudine æqualibus.

Scheff. Fung. Bav. t. 15. ☞ 15.
In sylvis frequens.

1049. AGARICUS *muscarius* stipitatus, lamellis dimidiatis solitariis; stipite volvato: apice dilatato, basi ovato.

Sterb. Fung. t. 22. f. A. Pauli Fl. Oec. 8. Müll. Dissert. de Fungis.

Dan. Flue-Svamp.

In pratis sylvaticis.

* * Parasitici, acaules, dimidiati.

1050. * AGARICUS *quercinus* acaulis, lamellis labyrinthiformibus. Buxb. Cent. V. t. 4. f. 5.

In truncis, trabibus ac pallis quercinis.

1051. * AGARICUS *alneus* acaulis, lamellis bifidis, verulentis.

In arboribus raro.

1052. * AGARICUS *betulinus* coriaceus villosus, margine obtuso, lamellis anastomosantibus.

In truncis raro.

391. BOLETUS. 1210.

* Stipitati.

1053. * BOLETUS *perennis*, stipitatus perennis: pilei utrinque planiusculo.

In caudicibus putridis.

1054. * *BOLETUS bovinus* stipitatus, pileo glabro pulvinato, marginato: poris compositis acutis: porulis angulatis brevioribus.

Schaff. Fung. Bav. t. 134. § 135. Müll. Monograph. de Fungis. t. 1. § 2.

In gramineis sylvaticis.

1055. * *BOLETUS luteus* stipitatus pileo pulvinato subviscido: poris rotundatis convexis flavissimis: stipite albo.

In sylvis.

1056. * *BOLETUS subtomentosus* stipitatus pileo flavo subtomentoso: poris subangulatis fulvis planis: stipite flavo.

In sylvis.

* * *Parasitici acaules.*

057. * *BOLETUS fomentarius* acaulis pulvinatus inæqualis obtusus: poris teretibus æqualibus glaucis.

Schaff. Fung. Bav. t. 137.

In fagis.

058. * *BOLETUS versicolor* acaulis, fasciis dicoloribus: poris albis.

In truncis & palis ligneis passim.

059. * *BOLETUS suaveolens* acaulis, superne lævis, fasciatus.

In salice frequens.

060. * *BOLETUS igniarius* acaulis pulvinatus lævis: poris tenuissimis.

In antiquis fagis frequens.

392. H Y D N U M. 1211.

061. * *HYDNUM repandum* stipitatum pileo convexo lævi flexuoso.

Mich. Gen. t. 72. f. 3. Schaff. Fung. Icon. t. 141.

In nemore frequens.

1062. * *HYDNUM auriscalpium* stipitatum, pileo
mediato.

Mich. Gen. t. 72. f. 8. Schaff. Fung. Icon. t. 14.
In sylvis rarius.

1063. * *HYDNUM parasiticum* arcuato-rugosum to-
mentosum.

In truncis ac lignis.

393. PHALLUS. 1212.

1064. *PHALLUS impudicus* volvatus stipitatus: pileo
celluloso.

Mich. Gen. t. 83. Fl. Dan. t. 175. Schaff. Fung.
Bavar. t. 196 - 198.

In luco copiose, ut & in luco *Silleroënsi*.

1065. † *PHALLUS esculentus* pileo ovali: stipite nudo
rugoso.

Mich. Gen. t. 85. f. 1. 2. Fl. Dan. t. 53.

394. CLATHRUS. 1213.

1066. * *CLATHRUS demudatus* capitulo oblongo
vato. *Mich. Gen. t. 49. f. 1.*

In aseribus humentibus.

1067. *CLATHRUS nudus* stipitatus capitulo oblongo
axi longitudinali adnato.

Mich. Gen. t. 94. Fl. Dan. t. 216.

In muscosis cespitibus.

1068. * *CLATHRUS recutitus* stipitatus capitulo glo-
boso: glande ovali.

In fossa cespitum ad *Kollekolle*.

¶ Character genericus melius in *CLATHRUM cancellatum*
nunquam a me inventum, quam in ceteros quadrat.

395. HELVELLA. 1214.

1069. * HELVELLA *Mitra* pileo deflexo adnato lobato difformi.

Mich. Gen. t. 86. f. 7. Fl. Dan. t. 116. Schaff. Fung. Bav. t. 154. & 162.

In cespitibus putrescentibus frequens.

396. PEZIZA. 1215.

1070. PEZIZA *lentifera* campanulata lentifera.

Mich. Gen. t. 102. f. 1. 2. 3. 4. Fl. Dan. t. 105. Schaff. Icon. Fung. t. 179.

In cespitosis, ramalibusque putridis, ac in areis hor-
torum frequens.

1071. * PEZIZA *punctata* turbinata truncata: disco punctato.

In fimo equino arido.

1072. * PEZIZA *cornucopioides* infundibuliformis: disco patente sinuato punctato.

Vaill. Bot. Paris. t. 13. f. 2. 3. Schaff. Fung. Icon. t. 165.

In vivario frequens.

1073. * PEZIZA *acetabulum* cyathiformis, extus angulata venis ramosis. *Vaill. Bot. Paris. t. 13. f. 1.*

In sylvis rarior.

1074. * PEZIZA *cyathoides*, cyathiformis: margine obtuso erecto.

In luco.

1075. * PEZIZA *cupularis*, globoso-campanulata, margine crenato. *Vaill. Bot. Paris. t. 11. f. 1. 2. 3.*

In ambulacro magno vivarii.

1076. * PEZIZA *scutellata* plana: margine convexo piloso. *Vaill. Bot. Paris. t. 13. f. 13. 14.*

In locis putrescentibus.

1077. * PEZIZA *cochleata* turbinata cochleata.
Vaill. Bot. Paris. t. 11. f. 8. Schæff. Fung. Icon.
t. 148.
 Ad caudices fagorum in sylva.

397. CLAVARIA. 1216.

* *Indivisa.*

1078. * CLAVARIA *pistillaris* clavæformis simplicissima
Mich. Gen. t. 87. f. 1. Schæff. Fung. Icon. t. 171.
 In sylva.
1079. * CLAVARIA *militaris* clavata integerrima: ca-
 pite squamoso. *Vaill. Bot. Paris. t. 7. f. 4.*
 In nemore & in fossa ad *Dronninggaard*; raro.
1080. * CLAVARIA *ophioglossoides* clavata integerrima
 compressa obtusa. *Vaill. Bot. Paris. t. 7. f. 3.*
 In pratis, paludosis ac subhumidis frequens.

* * *Ramosa.*

1081. * CLAVARIA *digitata* ramosa lignea nigra.
 In caudicibus faginis.
1082. * CLAVARIA *Hypoxylon* ramoso-cornuta com-
 pressa.
 Intra corticem & lignum truncorum excisorum.
1083. * CLAVARIA *coralloïdes* ramis confertis ramo-
 sissimis inæqualibus.
Vaill. Bot. Paris. t. 8. f. 4. Schæff. Fung. Icon. t. 177.
 In vivario.
1084. * CLAVARIA *fastigiata* ramis confertis ramosis-
 simis fastigiatis obtusis luteis.
 In sylvis.
1085. * CLAVARIA *muscoïdes* ramis ramosis acumina-
 tis inæqualibus luteis. *Schæff. Fung. Icon. t. 173.*
 Inter muscos versus montem cochlearem haud infre-
 quens.

398. LYCOPERDON. 1217.

* *Solidum, subterraneum.*

1086. * LYCOPERDON *cervinum* globosum, solidiusculum, lacerum, centro farinifero.

Mich. Gen. t. 99. f. 4.

In colliculo ruderibus exstructo ad dextram Villæ semel inveni.

* * *Pulverulenta, radicata supra terram.*

1087. LYCOPERDON *Bovista* subrotundum: lacerato dehiscens. *Schæff. Fung. Icon. t. 186.*

Dan. Ulve-Fis.

Pharmac. CREPITUS LUPI.

In campis sterilibus frequens.

1088. * LYCOPERDON *aurantium* sphaeroidale basi rugosum stipitatum: laciniis obtuse emarginatis dehiscens. *Vaill. Bot. Paris. t. 16. f. 9. 10.*

Ad latera arenosa fossarum.

1089. * LYCOPERDON *carpobolus* volva multifida: fructu globofo ex seminibus combinatis.

Mich. Gen. t. 101.

In asseribus subhumidis locisque cespitosi copiose.

* * * *Parasitica in farinam fatiscentia.*

1090. * LYCOPERDON *epidendrum* cortice farinaque purpurea.

Buxb. Cent. V. t. 29. f. 2. Schæff. Fung. Icon. t. 193.]

In lignis antiquis truncisque Cerasi.

1091. * LYCOPERDON *epiphyllum* aggregatum, ore multifido lacero: pulvere fulvo.

In dorso foliorum Tussilaginis Farfaræ ac Ranunculi Ficariæ.

399. MUCOR. 1218.

* Perennes.

1092. * MUCOR *spharocephalus* perennis, stipite filiformi nigro: capitulo globoso cinereo.

Hall. Helv. t. 1. f. 3.

In sepimentis.

1093. * MUCOR *Lichenoïdes* perennis stipite subulato nigro: capitulo lentiformi cinereo.

Dill. Musc. t. 14. f. 3.

In asseribus abietinis.

1094. * MUCOR *Embolus* seta nigra, villo fusco.

Hall. Helv. t. 1. f. 1.

In lignis putridis.

1095. * MUCOR *furfuraceus* perennis viridis, foliis furfuraceis: stipite filiformi: capitulo globoso.

In vallis cespitosis passim.

* * Fugaces.

1096. * MUCOR *mucedo* stipitatus, capsula globosa.

Mich. Gen. t. 95. f. 1.

In plantis.

1097. * MUCOR *glaucus* stipitatus, capitulo subrotundo aggregato.

Mich. Gen. t. 91. f. 1.

In pomis corruptis.

1098. * MUCOR *crustaceus* stipitatus: spicis digitatis.

Mich. Gen. t. 91. f. 3.

In cibis antiquis.

1099. * MUCOR *Erysiphe* albus, capitulis fuscis sessilibus.

In foliis humuli.



Ut pateat, quodnam incrementum Flora Patriæ intra
duo lustra ceperit, quorumque auctoritate noviciæ
hæ cives factæ, quas quisque addiderit, Appendicis loco,
quousque mihi innotuerunt, adjiciam.

A D D I T Æ

a H O L M I O

Callitriche verna.
autumnalis.

Veronica peregrina.

Scirpus acicularis.
maritimus.

Phleum pratense.

Agrostis rubra.
canina.

Aira aquatica.
præcox.

Poa trivialis.
annua.

compressa.

Festuca elatior.

Bromus sterilis.
giganteus.

Avena pratensis.

Lolium perenne.

Potamogeton maritimum.

Ruppia maritima.

Datura Stramonium.

Chenopodium urbicum.
glaucum.
vulvaria.

Sison inundatum.

Drosera longifolia.

Allium oleraceum.

Tulipa sylvestris.

Ornithogalum minimum.

Juncus filiformis.

Triglochin palustre.

Elatine hydropiper.

Sedum sexangulare.

Cerastium vulgatum.

Mespilus Cotoneaster.

Ranunculus polyanthemus.

Sonchus arvensis.

Carex leporina.

Populus alba.

Tremella Nostoc.

Auricula.

ab Ill. A L E N N E

Salsola hirsuta.

Mentha sylvestris.

Hypochaeris glabra.

Atriplex pedunculata.

P 3

a F O R S K Æ H L I O

Bromus triflorus.

a K O E N I G I O

Pilularia.

ab O E D E R O

Agrostis capillaris.

Cyperus fuscus.

Aira flexuosa.

Bromus pinnatus.

Avena elatior.

Centunculus minimus.

Scheuchzeria palustris.

Dianthus prolifer.

Melampyrum sylvaticum.

Subularia aquatica.

Lepidium ruderale.

Cardamine hirsuta.

Geranium lucidum.

Medicago polymorpha.

Hypochæris radicata.

Orchis ustulata.

Nam quam pro Ophry an-
thropophora exhibuit Cla-
rissimus Oeder, revera
Orchis ustulata est.

Asplenium trichomanes.

Chara hispida.

Clathrus nudus.

Potamogeton gramineus.

a M E

Trecentæ jam asterisco notatæ.

P L A N T Æ.

Quas centum abhinc annis *Kyllingius* in agro *Fri-
drichsdalensi* legerat, ex quibus *Lathraea Squama-
ria* & *Anemone Ranunculoïdes* jam in vicinas re-
giones demigrarunt.

Hippuris vulgaris.

Poa compressa.

Arundo epigeios.

Dipsacus Fullonum.

Gentiana Centaurium.
Selinum palustre.
Angelica Archangelica.
Linum Radiola.
Drosera longifolia.
Drosera rotundifolia.
Acorus calamus.
Paris quadrifolia.
Monotropa hypopitys.

Pyrola rotundifolia.
Prunus Padus.
Actæa spicata.
Nymphæa alba.
Anemone Ranunculoides.
Ballota nigra.
Lathræa Squamaria.
Ceratophyllum demersum.

P L A N T Æ,

Quæ quidem in viridario *Kyllingii* ac *Holmii* Prodromo Floræ Danicæ enumerantur, ob distantiam tamen locorum vere natalium ægre indigenæ censendæ, hinc Flora nostra Danica exclusæ.

Schoenus mucronatus.
Panicum glaucum.
Lonicera alpigena.
Atropa Belladonna.
Lychnis Chalcedonica.
Potentilla opaca.

Hesperis matronalis.
Amaranthus albus.
Atriplex hortensis.
Lycopodium alpinum.
Fucus saccharinus.



FAUNÆ FRIDRICHSDALINÆ
 NOVICIA,
 QUÆ ANNI SPATIO IN AGRO
 FRIDRICHSDALENSI LEGI.
 INSECTA.
 COLEOPTERA.

SCARABÆUS

subterraneus.
 erraticus.
 conspurcatus.
 hemipterus.
 brunneus.
 quadrimaculatus.
 aquaticus.
 rufipes.
 parallelepipedus.

DERMESTES

pertinax.
 violaceus.
 Scrophulariæ.
 musæorum.

SILPHA

littoralis.
 quadripunctata.
 aquatica.

COCCINELLA

unipunctata.
 19 - punctata.
 22 - punctata.

COCCINELLA

bipustulata.
 decempustulata.

CHRYSOMELA

Tanaceti.
 vulgatissima.
 pallida.
 marginella.
 Helxines.
 exoleta.
 quadripunctata.
 Coryli.
 Nymphææ.
 Caprea.
 Cerasi.
 Cyanella.

CURCULIO

Alliariæ.
 Quercus.
 Polygoni.
 Betulæ.
 Populi.
 cervinus.

ERAMBYX

Coriarius.
Textor.

ANTHARIS

biguttata.
minima.

LATER

ruficollis.
livens.
marginatus.
murinus.
minutus.

CICINDELA

hybrida.

BYTISCUS

fuscipes.
luridus.
striatus.
erythrocephalus.
scarabæoides.
minutus.

CARABUS

violaceus.
melanocephalus.

CARABUS

multipunctatus.
quadrinaculatus.

TENEBRIO

culinaris.
Fosfor.
gibbus.
Caraboides.

STAPHYLINUS

riparius.
lignorum.
biguttatus.
chrysomelinus.
Boleti.
subterraneus.

GRYLLUS

bipunctatus.
subulatus.
verrucivorus.
apricarius.
biguttulus.
rufus.
grossus.
pedestris.

HEMIPTERA.

CICADA

cornuta.
nervosa.
leucophthalma.
leucocephala.
lateralis.

CICADA

striata.
flavicollis.
vittata.
Ulmi.
Rosæ.
P 5

CIMEX

bidens.
 hæmorrhoidalis.
 coeruleus.
 biguttatus.
 annulatus.
 ater.
 umbratilis.
 nemorum.
 Rolandri.
 lævigatus.

CIMEX

striatus.
 dolabratus.
 sylvestris.
 Coryli.

APHIS

Pastinacæ.
 Cirsii.

THRIPS

minutissima.

LEPIDOPTERA.

PAPILIO

Pl. Betulæ.
 Pl. Rubi.

SPHINX

Elpenor.
 Porcellus.

PHALÆNA

B. Camelina.
 N. Humuli.
 N. Hec̃ta.
 N. Gothica.
 N. Typica.
 N. pallens.
 G. vernaria.
 G. strigilata.
 G. rectangulata.
 G. cratægata.
 G. lemnata.
 G. remutata.
 Tor. fasciana.

PHALÆNA

Tor. Branderiana.
 Tor. Xylosteana.
 Tor. Rosana.
 Tor. Oporana.
 Tor. Solandriana.
 Tor. Modeciana.
 Tor. Heraclea.
 Tin. colonella.
 Tin. pinetella.
 Tin. culmella.
 Tin. trigonella.
 Tin. pruniella.
 Tin. Petiverella.
 Tin. Swammerda.
 mella.
 Tin. gemmella.
 Tin. picarella.
 Tin. Xylostella.
 Tin. Jungiella.
 Tin. vestianella.

ALÆNA

Tin. granella.
Tin. bractœella.
Tin. argentella.
Tin. tessella.

PHALÆNA

Tin. bicostella.
Tin. cinerella.
Tin. fuscella.
Tin. porrectella.

NEUROPTERA.

EMERA

culiciformis.
horaria.
diptera.

PHRYGANEÆ

bimaculata.
umbrosa.

HEMEROBIUS

sexpunctatus.

HYMENOPTERA.

THREDO

femorata.
Padi.
Salicis.
atra.
ovata.
septentrionalis.
duodecimpunctata.

ICHNEUMON

fossorius.
peregrinator.
incubitor.
compunctor.
delusor.
tirillator.
Refinellæ.
prærogator.
enervator.
pugillator.
ramidulus.
glaucopterus.
circumflexus.
cinctus.
Puparum.

NEUMON

extensorius.
culpatorius.
infractorius.
faturatorius.
molitorius.
pisorius.
luctatorius.
vaginatorius.
lituratorius.
deliratorius.

SPHEX

figulus.
fusca.

SPHEX

pectinipes.
variegata.
gibba.
fossoria.

CHRYSIS

cyanea.

VESPA

cribraria.
spinipes.

APIS

tumulorum.
retusa.

APIS

rufa.
truncorum.
fuocincta.
coeruleescens.
cunicularia.
floriformis.
conica.
acervorum.
subterranea.

FORMICA

fusca.
obsoleta.
cespitem.

DIPTERA.

TIPULA

contaminata.
Marci.
brevicornis.
florilega.
juniperina.

MUSCA

plebeja.
microleon.
tringaria.
intricaria.
transfuga.
segniis.
femorata.
canicularis.

MUSCA

cellaris.
meteorica.
polita.
pubera.
faltuum.
germinationis.
cerasi.

CONOPS

irritans.

ASILUS

germanicus.
crabroniformis.

A P T E R A.

ISMA
faccharina.

PODURA
nivalis.
arborea.

URA
aquatica.

TERMES
fatidicum.

V E R M E S.

I N T E S T I N A.

ARIS
vermicularis.

MBRICUS
terrestris.

CIOLA
hepatica.

HIRUDO
fanguifuga.
medicinalis.
stagnalis.
complanata.

M O L L U S C A.

MAX
ater.
rufus.
maximus.

LIMAX
agrestis.
flavus.

T E S T A C E A.

LLINA
cornea.

STILUS
anatinus.

ELIX
planorbis.

HELIX
complanata.
vortex.
cornea.
nemoralis.
stagnalis.
putris.
auricularia.

NERITA
fluviatilis.

PATELLA
lacustris.

L I T H O P H Y T A.

TUBIPORA
Urceus.

Z O O P H Y T A.

ISIS
anastatica.

VOLVOX
globator.

HYDRA
Polypus.
socialis.



INDEX GENERUM.

acer	pag. 183. 185.	Antirrhinum	106. 113. 114.
chillea	142. 156.	Aphanes	23. 27.
conitum	95. 99.	Apium	35. 54.
corus	58. 63.	Aquilegia	96. 99.
costichum	186. 191.	Arabis	116. 121.
ctæa	95. 97.	Arbutus	74. 77.
cloxa	68. 72.	Arctium	140. 147.
gopodium	35. 54.	Arenaria	75. 81.
thufa	34. 53.	Arnica	141. 154.
aricus	189. 220. 221. 222.	Artemisia	140. 150. 151.
rimonia	85. 86.	Arundo	9. 20.
rostemma	76. 82.	Arum	160. 164.
roftis	8. 14. 15.	Asarum	85. 86.
ra	8. 15. 16.	Asclepias	32. 46.
uga	104. 107.	Asparagus	58. 62.
chemilla	23. 27.	Asperugo	30. 38.
ifma	59. 66.	Asperula	22. 25.
lium	58. 60. 61.	Asplenium	186. 192.
opecurus	8. 14.	Aster	142. 153.
line	35. 55.	Astragalus	128. 134.
thæa	123. 125.	Athamanta	34. 51.
agallis	30. 40.	Atriplex	183. 184. 185.
ichufa	30. 37. 38.	Avena	9. 19. 20. 183.
idromeda	74. 77.		
lemone	96. 100.	B.	
gelica	34. 51.	Ballota	105. 110.
themis	142. 156.	Bellis	141. 155.
thericum	58. 62.	Betonica	104. 107.
thoxanthum	3. 6.	Betula	166. 172. 173.
thyllis	127. 131.	Bidens	140. 149.

INDEX GENERUM.

Blasia	188. 209.	Chenopodium	33. 46. 47.
Blitum	1. 2.	Chrysanthemum	141. 116.
Boletus	189. 222. 223.	Chrysosplenium	74. 7.
Brassica	116. 121.	Cichorium	139. 14.
Briza	8. 17.	Cicuta	35. 52.
Bromus	9. 18. 19.	Cineraria	141. 154.
Bryonia	167. 175.	Circæa	3. 4.
Bryum	187. 199. 200.	Cistus	95. 98.
	201. 202.	Clathrus	189. 224.
Bunias	116. 122.	Clavaria	189. 226.
Bupleurum	34. 50.	Clinopodium	105. 110.
Butomus	73	Cnicus	140. 149.
Buxbaumia	187. 196.	Cochlearia	115. 118.
Byssus	188. 219.	Comarum	89. 94.
		Conserva	189. 218.
C.		Conium	34. 50.
Calla	160. 164.	Convallaria	58. 62. 63.
Caltha	96. 103.	Convolvulus	31. 42.
Callitriche	1. 2.	Conyza	141. 152.
Campanula	31. 41. 42.	Coreopsis	142. 115.
Cardamine	116. 119.	Cornus	23. 57.
Carduus	140. 148.	Coronilla	128. 130.
Carex	165. 169. 170. 171.	Corrigiola	35. 55.
	172.	Corylus	167. 155.
Carlina	140. 149.	Crambe	116. 122.
Carpinus	167. 175.	Cratægus	88. 92.
Carum	35. 54.	Crepis	139. 146.
Caucalis	33. 50.	Cucubalus	75. 80.
Centaurea	142. 157.	Cuscuta	23. 27.
Centunculus	22. 27.	Cynoglossum	30. 38.
Cerastium	75. 83.	Cynolurus	9. 17.
Geratophyllum	167. 173.	Cyperus	7. 11.
Chærophylum	34. 53.	Cypripedium	160. 164.
Chara	189. 215.		
Chelidonium	95. 97.	D.	
		Dactylis	8. 17.
			Daphn.

INDEX GENERUM.

Naphne	68. 71.	Fragaria	89. 93.
Natura	31. 43.	Fraxinus	183. 185.
Naucus	33. 50.	Fucus	189. 216. 217.
Nelphinium	95. 99.	Fumaria	127. 129.
Nentaria	116. 119.		
Nianthus	75. 79.		
Nipfacus	22. 24.		
Nraba	115. 117.		
Nrosera	36. 56.		

E.

Echium	36. 39.
Elatine	68. 72.
Elvela	189.
Elymus	9. 21.
Empetrum	176. 180.
Epilobium	68. 69.
Equisetum	186. 190.
Erica	68. 70. 71.
Erigeron	141. 152.
Eriophorum	7. 12.
Ervum	128. 133.
Eryngium	49. 33.
Erysimum	116. 120.
Euonymus	32. 45.
Eupatorium	140. 149.
Euphorbia	85. 87.
Euphrasia	105. 112.

F.

Fagus	167. 174.
Festuca	8. 17. 18.
Filago	142. 157. 158.
Fontinalis	187. 196.

G.

Galeopsis	104. 109.
Galium	22. 25. 26.
Genista	127. 130.
Gentiana	32. 48. 49.
Geranium	123. 124. 125.
Geum	89. 94.
Glaux	32. 46.
Glecoma	104. 108.
Gnaphalium	141. 151. 178.

H.

Hedera	32. 45.
Hedysarum	128. 133.
Helvella	225.
Heracleum	33. 51.
Herniaria	33. 46.
Hieracium	139. 145. 146.
Hippophaë	176. 181.
Hippuris	1. 2.
Holcus	183. 184.
Hordeum	9. 21.
Hottonia	31. 40.
Humulus	177. 181.
Hyacinthus	58. 63.
Hydnum	189. 223. 224.
Hydrocharis	177. 182.
Hydrocotyle	33. 49.
Hyoscyamus	31. 43.

Q

INDEX GENERUM.

Hyoferis	140. 146.	Lichen	188. 209. 210
Hypericum	137. 138.		211. 212. 213. 214. 215
Hypnum	187. 202. 203.	Ligusticum	34. 7.
	204. 205. 206.	Lilium	58. 6.
Hypochæris	139. 146.	Limofella	106. 114.
	147.	Linum	36. 56.
		Lithospermum	30. 37.
		Lolium	9. 20. 21.
		Lonicera	32. 42.
		Lotus	128. 135. 136.
		Lunaria	115. 119.
		Lychnis	76. 82. 83. 178.
		Lycoperdon	189. 227.
		Lycopodium	187. 195.
		Lycopsis	30. 39.
		Lycopus	3. 6.
		Lyfimachja	30. 40.
		Lythrum	85. 86.

I.

Jasione	142. 158.
Iberis	115. 118.
Ilex	23. 28.
Impatiens	142. 159.
Imperatoria	34. 53.
Inula	141. 154.
Iris	7. 10.
Ifatis	116. 122.
Ifoetes	186. 194.
Juncus	58. 63. 64.
Jungermannia	188. 206.
	207. 208.
Juniperus	178. 182.

L.

Lactuca	139. 144.
Lamium	104. 108. 109.
Lapsana	139. 147.
Lathræa	105. 113.
Lathyrus	128. 131. 132.
Ledum	74 77.
Lemna	165. 168.
Leontodon	139. 144. 145.
Leonurus	104. 110.
Lepidium	115. 117.

M.

Malva	123. 126.
Marchantia	187. 208. 209.
Marrubium	105. 110.
Matricaria	141. 155. 156.
Medicago	128. 136.
Melampyrum	105. 112.
Melica	8. 16.
Mentha	104. 107. 108.
Menyanthes	31. 39.
Mercurialis	177. 182.
Mespilus	88. 91.
Milium	8. 14.
Mnium	187. 197. 198. 199.
Monotropa	74 77.
Montia	9. 24.

INDEX GENERUM.

Mucor	189. 228.	Paris	68. 72.
Myagrum	115. 117.	Parinassia	36. 56.
Myosotis	30. 37.	Pastinaca	35. 54.
Myosurus	36. 57.	Pedicularis	106. 113.
Myrica	177. 181.	Peplis	59. 64.
Myriophyllum	166. 173.	Peziza	189. 225. 226.
	174.	Phalaris	8. 13.
		Phallus	189. 224.
		Phascum	187. 195. 196.
		Phellandrium	34. 52.
		Phleum	8. 13. 14.
		Physalis	31. 44.
		Phyteuma	31. 42.
		Picris	139. 143.
		Pilularia	186. 194.
		Pimpinella	35. 54.
		Pinguicula	3. 5.
		Pisum	128. 131.
		Plantago	22. 26.
		Poa	8. 16. 17.
		Polygala	127. 129.
		Polygonum	68. 71. 72.
		Polypodium	186. 192.
			193. 194.
		Polytrichum	187. 197.
		Populus	177. 181. 182.
		Potamogeton	23. 28. 29.
		Potentilla	89. 93.
		Poterium	167. 174.
		Prenanthes	139. 144.
		Primula	31. 39.
		Prunella	105. 111.
		Prunus	88. 90.
		Pteris	186. 191.
		Pulmonaria	30. 38.
		Pyrola	74. 77. 78.

N.

Narcissus	58. 60.
Nardus	7. 13.
Nepeta	104. 107.
Nymphæa	95. 98.

O.

Oenanthe	33. 52.
Oenothera	68. 69.
Ononis	127. 130.
Onopordon	140. 149.
Ophioglossum	186. 190.
Ophrys	160. 162. 163.
Orchis	160. 161. 162.
Origanum	105. 110.
Ornithogalum	58. 61. 62.
Ornithopus	128. 133.
Orobus	128. 131.
Osmunda	186. 191.
Oxalis	76. 82.

P.

Panicum	7. 13.
Papaver	95. 97. 98.
Parietaria	183. 184.

INDEX GENERUM.

Pyrus	88. 91.	Schoenus	7. 10. 11.
		Scirpus	7. 11. 12.
Q.		Scleranthus	74. 78. 79.
		Scorzonera	139. 143.
Quercus	167. 174.	Scrophularia	106. 114.
		Scutellaria	105. 111.
R.		Sedum	75. 81. 82.
		Selinum	34. 50.
Ranunculus	96. 101. 102.	Sempervivum	85. 87.
Raphanus	116. 122.	Senecio	141. 152. 153.
Refeda	85. 86.	Serapias	160. 163.
Rhamnus	32. 44.	Serratula	140. 147. 148.
Rhinanthus	105. 111.	Sherardia	22. 24.
Ribes	32. 45.	Silene	75. 80.
Riccia	188. 209.	Sinapis	116. 121.
Rosa	88. 91. 92.	Sifon	34. 52.
Rubus	89. 92. 178.	Sifymbrium	116. 119. 120.
Rumex	59. 65. 177.	Sium	52. 34.
Ruppia	23. 29.	Solanum	31. 44.
		Solidago	141. 153.
S.		Sonchus	139. 143. 144.
		Sorbus	88. 90.
Sagina	23. 29.	Sparganium	165. 169.
Sagittaria	166. 174.	Spartium	127. 130.
Salicornia	1. 2.	Spergula	75. 84.
Salix	176. 179. 180.	Sphagnum	187. 195.
Salsola	33. 48.	Spiræa	88. 91.
Sambucus	35. 55.	Spongia	189. 220.
Samolus	31. 42.	Stachys	104. 109.
Sanicula	33. 49.	Statice	36. 56.
Saponaria	75. 79.	Stellaria	75. 80. 81.
Satyrrium	160. 162.	Stratiotes	96. 99.
Saxifraga	74. 78.	Subularia	115. 117.
Scabiosa	22. 24.	Symphytum	30. 38.
Scandix	34. 53.		
Scheuchzeria	59. 66.		

INDEX GENERUM.

T.

Tanacetum	141. 150.
Taxus	178. 182.
Teucrium	104. 107.
Thalictrum	96. 101.
Thlaspi	115. 118.
Thymus	105. 111.
Tilia	95. 98.
Tordylium	33. 50.
Tormentilla	89. 94.
Tragopogon	139. 143.
Tremella	188. 215. 216.
Trientalis	67.
Trifolium	128. 134. 135.
Triglochin	59. 66.
Triticum	9. 21.
Trollius	96. 103.
Tulipa	58. 61.
Turritis	116. 121.
Tussilago	141. 152.
Typha	165. 168.

V.

Vaccinium	68. 70.
-----------	---------

Valeriana	7. 10.
Verbascum	31. 43.
Verbena	3. 6.
Veronica	3. 4. 5.
Viburnum	35. 55.
Vicia	128. 132. 133.
Viola	142. 158. 159.
Viscum	176. 180.
Ulmus	33. 48.
Ulva	188. 217.
Urtica	166. 173.
Utricularia	3. 6.

X.

Xanthium	166. 173.
----------	-----------

Z.

Zannichellia	165. 168.
Zostera	160. 164.



NOMINA DANICA.

A.

Ager - Urt. pag. 155.

Aakande 98.

Abrod, vild 150

Adel - Klefver 100.

Arenpriis 4.

Æsk 185.

Æspe - Træ 181.

Agermaane 86.

Ager - Kaal 121. 122.

Akkeley 99

Alm 48

Amur 134.

Andemad 168.

Angelik 51.

Arfve 40.

Arum 164.

Aspargis 62.

Attik 55.

Aunbög 175.

Axelbar 90.

B.

Baldrian 10.

Basilikum, vild 111.

Beeg - Urt 83.

Beenbræk 62.

Beenved 45.

Besver - Gras 17.

Begger - Moos 214.

Benedict - Urt 94.

Beske - Klefver 39.

Betonie 107.

Bettler - Luus 10.

Bevæ - Æsk 181.

Beyler 25.

Bierg - Lakrits 134.

Bierg - Nellike 56.

Bierg - Roed 54.

Bierg - Mandel 91.

Biergmands - Træ 49.

Bingel - Urt 182.

Bink 172.

Bitter Pile - Urt 71.

Bitter Söde 4.

Biörne - Kløj 51.

Biörne - Lap 51.

Blaa - Ber 70.

Blaa - May 39.

Blaa - Urt 157.

Blaatoppet - Græs 15.

Blaatroid - Urt 152.

Blod - Blomster 138.

Blod - Roed 94.

Bokar 25.

Bornholms Rosner 90.

Brænde - Hætte 173.

Brambar 92.

N O M I N A D A N I C A.

<i>Frue - Bær</i>	92.	H.
<i>Fugle - Bær</i>	55.	
<i>Fugle - Græs</i>	55.	<i>Haar - Væxt</i>
<i>Fugle - Liim</i>	180.	<i>Hage - Bær</i>
<i>Fugle - Melk</i>	61. 87.	<i>Haf - Torn</i>
<i>Følle - Fod</i>	152.	<i>Hage - Torn</i>

G.

<i>Gaase - Skrappe</i>	173.	<i>Hampe - Nelde</i>	139
<i>Gaase - Urt</i>	93. 155. 156.	<i>Hane - Fod</i>	101. 102
<i>Gæslinge - Blomster</i>	117.	<i>Hane - Kamm</i>	111. 129
<i>Galle - Bær</i>	175.	<i>Hane - Kamm - Græs</i>	17
<i>Giedde - Skiæg</i>	91	<i>Hare - Fod</i>	151
<i>Gienem - Væxt</i>	50.	<i>Hare - Kaal</i>	147
<i>Gier</i>	117.	<i>Hare - Klever</i>	135
<i>Giøge - Mad</i>	82.	<i>Hassel</i>	175
<i>Glas - Urt</i>	2.	<i>Havn - Torn</i>	98
<i>Gonsalve - Rød</i>	38.	<i>Havre - Græs</i>	14
<i>Gotberaad</i>	66.	<i>Heede - Nödder</i>	15
<i>Graabone</i>	150.	<i>Helle - Knop</i>	8
<i>Græsløg</i>	61.	<i>Heste - Hov</i>	15
<i>Græs - Nellike</i>	82.	<i>Heste - Mynthe</i>	15
<i>Grine ved Middag</i>	40.	<i>Heste - Rumpe</i>	15
<i>Guds - Haand</i>	162. bis.	<i>Heyre</i>	18
<i>Guldblomme</i>	154.	<i>Heyre - Græs</i>	18
<i>Guld - Haar</i>	197.	<i>Hielme</i>	20.
<i>Guld - Sæl - Oeye</i>	145.	<i>Hierte - Span</i>	115
<i>Guul - Arfve</i>	40.	<i>Himmel - Dug</i>	16
<i>Guul - Klever</i>	136.	<i>Hierte - Tunge</i>	19
<i>Guul - Penge - Blad</i>	40.	<i>Hofmod</i>	83
<i>Guul Skorpe - Moos</i>	211.	<i>Horse - Tænder</i>	43
<i>Gylden Vind - Urt</i>	153.	<i>Hvas - Græs</i>	17
<i>Gøgs - Urt</i>	161.	<i>Hvas - Sive - Græs</i>	18
		<i>Hvid - Klever</i>	134
		<i>Hvidløg - Urt</i>	12
		<i>Hvid Regn - Fan</i>	15
		<i>Hvid - Roed</i>	6
		<i>Hvid - Sippe</i>	10

N O M I N A D A N I C A.

<i>Hvid - Torn</i>	90.
<i>Hvide - Ved</i>	100.
<i>Hvine - Græs</i>	14.
<i>Humle</i>	181.
<i>Humle - Klover</i>	135.
<i>Hunde - Bær</i>	44. 175.
- Græs	21.
- Noffer	161.
- Tunge	37. 38.
- Urt	156.
<i>Hundrede - Hoved</i>	49.
<i>Hundrede - Knuder</i>	72.
<i>Huul - Røed</i>	129.
<i>Huus - Løg</i>	87.
<i>Hyld</i>	55.
<i>Hypericon</i>	138.
<i>Hynde - Taske</i>	118.
<i>Hyven - Torn</i>	92.
<i>Høgs - Urt</i>	145. 146.
<i>Hønse - Bær</i>	27.
<i>Hønse - Græs</i>	72.
<i>Hør - Silke</i>	27.

J.

<i>Jacobs - Urt</i>	153.
<i>Jern - Urt</i>	6.
<i>Jomfrue - Haar</i>	197.
<i>Jomfrue - Kær</i>	27.
<i>Jord - Bær</i>	93.
- Humle	138.
- Must	195.
- Rødder	132.
- Røg	129.
- Vedbende	108.
<i>Juttel - Trøst</i>	148.

K.

<i>Kuars - Blomster</i>	129.
<i>Kaars - Urt</i>	152.
<i>Kabbeleje</i>	103.
<i>Kafle - Græs</i>	14.
<i>Kalmus</i>	63.
<i>Kameel - Blomster</i>	155.
<i>Karde - Tidsel</i>	24.
<i>Kat - Ost</i>	126.
<i>Katte - Klover</i>	131.
- Urt	86. 157.
- Oeyne	109.
<i>Kiar - Uld</i>	12.
<i>Kiærligheds - Græs</i>	17.
<i>Kierlinge - Kruus</i>	98.
<i>Kierlinge - Tænder</i>	136.
<i>Klammer - Urt</i>	25.
<i>Klinte</i>	82.
<i>Klittetag</i>	20.
<i>Klokker</i>	41. 42.
<i>Kna - Græs</i>	81. 84.
<i>Knævel</i>	78.
<i>Knæpi - Panden</i>	101.
<i>Knop - Urt</i>	157. bis.
<i>Knoppe - Græs</i>	21.
<i>Koe - Bield</i>	100.
<i>Koe - Blomme</i>	103.
<i>Koe - Driver</i>	39.
<i>Koe - Huede</i>	112.
<i>Konge - Lys</i>	43.
<i>Konning</i>	110.
<i>Kor - Bær</i>	92.
<i>Korsbær - Torn</i>	44.
<i>Kørn - Blomster</i>	157.
<i>Kørn - Rose</i>	97.

Q.

N O M I N A D A N I C A .

Kraft - Urt	81.	Manne - Gras	18.
Krage - Bar	180.	Mare - Torn	28.
Krage - Kloer	136.	Mari - Gras	162.
Krage - Taer	102.	Marie Sengebalm	25.
Krage - Torn	130.	Marrube	112.
Kroſing - Bar	70.	Marts - Violer	158.
Kruſed - Skreppe	65.	Meelbar - Riis	77.
Kulſukker - Rod	38.	Milt - Urt	191.
Kummen	54.	Multe - Bar	92.
Körvel	53.	Munke - Hatte	99.

L.

Ledmye	4.	Muse - Rumpa	57.
Lemmike	4.	Muse - Urt	151. 158.
Lever - Blomſter	56.	Muur - Rude	192.
Lever - Urt	208.	Muus - Erter	131. 132.
Lilie - Konval	62.	Muus - Oere	145.
Lind	98.	Myntke	107.
Lunge - Moos	212.	Myre - Sif	11.
Lunge - Urt	38.	Myſſike	25.
Luus - Urt	113.	Mød - Urt	91.
Lyng	70.		
Lyſe - Sif, Glat	63.		
Lyſe - Sif, Skarp	64.		
Løbe - Roſe	42.		
Løg	60.		
Løgs - Urt	107.		
Løn	185.		
Løve - Fod	27.		
Løve - Tund	144.		

M.

Maane - Rude	191.		
Mal - Urt	150.		

N.

Nalde	173. bis.
Naur	185.
Nellike - Roed	94.

O.

Onde - Urter	155.
Orme - Gras	162.
Orme - Urt	114.
Oxe - Tunge	37. 39. bis.
Oxe - Oere	100.
Oxe - Oeye	155.

N O M I N A D A N I C A.

P.

astke - Lilia	60.
eders - Urt	184.
enge - Blad	40.
enge - Urt	118.
terfilie	54.
il	179.
il - Urt	174.
ile - Urt	71.
indsvin - Knoppe	169.
iricum	138.
isse - Urt	46.
appel - Piil	181. 182.
appel - Træ	182.
ors	181.
ræste - Kone	144.

Q.

duæse - Urt	54.
-------------	-----

R.

Rave - Rumpe	14. 112.
Rams	61. 161.
Rapuntzel	41.
Ravne - Fod	26.
Reinfand	150.
Rensdyr - Moos	214.
Revrumpe - Græs	14. 112.
Rhabur	65.
Ribs	45.
Ridderfjore	99.
Roberts - Urt	124.
Rod - arfve	40.

Rød Hane - Kamm	129.
Rød Klever	135.
Rød - Moos	40.
Rød - Skreppe	65.
Rollike	156.
Røn	90.
Rør	20.

S.

Salomons Segl	63.
Salt - Urt	2.
Sanikel	49.
St. Ellens - Roed	154.
St. Hans - Urt	81.
St. Jacobs - Urt	153.
St. Peders - Urt	184.
St. Stephens - Urt	4.
Seene - Græs	20.
Senip	121.
Sey - Piil	179.
Sey - Urt	110.
Sif	11.
Sif - Græs	64.
Skab - Urt	24.
Skampion	220.
Skærntyde	50. 52. 53.
Skaw - Græs	190.
Skjæne	172.
Skierbugs - Urt	118.
Skjör - Piil	179.
Skordium	107.
Skorpions - Urt	37.
Skov - Æble	91.
Skov - Melde	2.

N O M I N A D A N I C A .

<i>Skov - Merke</i>	25.	<i>Strand - Kaal</i>	12
<i>Skov - Tidsel</i>	28.	<i>- Karfe</i>	112
<i>Slaaen - Torn</i>	90.	<i>- Klever</i>	28
<i>Slange - Hoved</i>	39.	<i>- Krage - Taer</i>	184
<i>Slange - Myrder</i>	143.	<i>- Melde</i>	184
<i>Slange - Tunge</i>	190.	<i>- Portulac</i>	65
<i>Slange - Urt</i>	71.	<i>- Skreppe</i>	66
<i>Smaa - Klokke</i>	41.	<i>- Siv - Gras</i>	113
<i>Smal - Blad</i>	146.	<i>- Stierne - Urt</i>	26
<i>Smör - Urt</i>	102. 103.	<i>- Veybred</i>	49
<i>Snerle</i>	72.	<i>Strandmands - Troe</i>	190
<i>Snerre</i>	26.	<i>Stude - Knæ</i>	97
<i>Snerre</i>	41.	<i>Svale - Urt</i>	46
<i>Soe - Bønne</i>	43.	<i>Svale - Roed</i>	10
<i>Soel - Bär</i>	45.	<i>Sverd - Lilie</i>	144
<i>Soel - Dug</i>	56.	<i>Svine - Dild</i>	144
<i>Sommer - Hyld</i>	55.	<i>Svine - Tidsel</i>	44
<i>Sov - Bar</i>	44.	<i>Svine - Urt</i>	22
<i>Spids Strand - Gras</i>	21.	<i>Svingel</i>	135
<i>Spring - Urt</i>	159.	<i>Sue - Kop</i>	135
<i>Squalder - Kaal</i>	54.	<i>Sækker - Bröd</i>	14
<i>Steenbreck</i>	37. 78.	<i>Suppe - Gras</i>	65
<i>Steenbregne - Moos</i>	203.	<i>Suur - Kaal</i>	82
<i>'Steen - Karfe</i>	118.	<i>Suur - Klever</i>	163
<i>- Klever</i>	134. 135.	<i>Sygs - Knap</i>	61
<i>- Klinte</i>	37.	<i>Syrer</i>	93
<i>- Rude</i>	192.	<i>Salv - Urt</i>	
<i>Steenföd - Roed</i>	192.		
<i>Stinkende Lactuc</i>	144.		
<i>Stolt Henrik</i>	46.	<i>T.</i>	
<i>Stor - Skreppe</i>	147.	<i>Tand - Urt</i>	78. 113
<i>Stor - Snerle</i>	41	<i>Tang</i>	164
<i>Storke - Næb</i>	124.	<i>Tiere - Urt</i>	80. 83
<i>Storm - Hat</i>	99.	<i>Tidse</i>	181
<i>Strand - Gras</i>	164.	<i>Tidsel</i>	148. 161
<i>- Isop</i>	46.	<i>Toblac</i>	154
		<i>Torden - Urt</i>	

N O M I N A D A N I C A.

Armentil	94.	Vau	86.
Ariske-Mund	113.	Vedbende	45.
Arst	110.	Vey - Bred	26.
Arane - Bær	70.	Vey - Græs	72.
Arane - Hals	124.	Vibe - Fæt	6.
Arold - Smør	216.	Viddie	180.
Arold - Urt	152.	Vikker	131. 132.
Arind - Dyder	155.	Vild Abrod	150.
Arind - Fryd	155.	Basilicum	111.
Arind - Gylden	48.	Cornel - Træ	27.
Artte - Bær	70.	Cypres	150.
Arste - Træ	44.	Erter	132.
		Gule-Rödder	50. 51.
		Havre	19.
		Hör	113. 117.
		Kaal	121. bis.
Ark - Urt	63.	Kirse - Bær	90.
Arldbirk	185.	Klokker	42.
Arldmue	97.	Korvel	53. bis.
Armosøster	161.	Lakrits	134.
Arnd - Aloë	99.	Løg	60.
- Arfve	45.	Meyran	110.
- Græs	16.	Nellikær	79.
- Hyld	55.	Nysseroed	163.
- Ingfer	164.	Pastinakel	50. 54.
- Karse	119. bis.	Peber - Roed	118.
- Krage - Taer	49.	Peterfilie	53.
- Marru	6.	Pimpinelle	54.
- Merke	54.	Rosmarin	77.
- Moos	196.	Senip	120.
- Pastinak	52. bis.	Thymian	111.
- Peber	71.	Vinter - Grøn	67. 77.
- Peberoed	120.	Vinter - Karse	120.
- Poley	108.	Volverley	154.
- Portulac	64.	Vor - Frues - Særk	41.
- Røllike	6. 40.	Vor - Frues Sengehaln	111.
- Stenbrek	52.		

NOMINA DANICA.

Vorte - Urt
Vrie - Torn

87. *Ulve - Fod*
44. *Ulve - Melk*
Urue


U.

Oe.

Uldborrer
Ulf - Bar
Ulve - Fiis

38.
72. *Oeyen - Rod*
227. *Oeyen - Trost*

80. 112



NOMINA PHARMACEUTICA.

A.

A bsinthium vulgare	150.
Acacia nostras	90.
acetosa nostras	65.
acetosella	82.
corus palustris	10.
dianthum aureum	197.
dianthum rubrum	192.
grimonia	86.
lechmilla	27.
likekengi	44.
liaria	120.
angelica fativa	51.
angelica sylvestris	51.
merina	93.
nitirrhinum	114.
onium	54.
quilegia	99.
histolochia fabacea	129.
moracia	118.
nica	154.
temisia	150.
arum	86.
paragus	62.
triplex sylvestris	47.
ricula Judæ	216.
ricula muris	145.

B.

rdana	147.
-------	------

Beccabunga	4.
Bellis minor	155.
Bellis pratensis	155.
Betonica	107.
Bonus Henricus	46.
Bryonia	175.
Buglossa	37.
Bursa pastoris	118.

C.

Calamus vulgaris	63.
Caprifolium	42.
Cardiaca	110.
Carvum	54.
Caryophyllata	94.
Centaurium minus	48.
Centumnodia	72.
Cerefolium	53.
Chamæmorus	92.
Chamomilla nostras	155.
Chelidonium majus	97.
Chelidonium minus	101.
Cichorium	147.
Cicuta	50. 52.
Cicutaria	53.
Cochlearia	118.
Consolida major	38.
Consolida media	107.
Consolida regalis	99.
Crepitus lupi	227.

NOMINA PHARMACEUTICA.

Cuscuta	27.	Geranium robertian.	124.
Cyanus	157.	Gnaphalium	158.
Cynoglossa	38.	Gramen	2.

D.

Daucus sylvestris

E.

Ebulus

Enula

Epithymum

Equisetum

Eryngium

Erysimum

Esula

Eupatorium

Euphrasia

F.

Fagus

Farfara

Fibrinum trifolium

Filix

Fragaria

Frangula

Fraxinus

Fumaria

Fungus Sambucinus

G.

Genista

Gentianella

H.

50. Hedera 45.

Hedera terrestris 108.

Hepatica alba 56.

Hepatica fontana 208.

55. Hepatica nobilis 100.

154. Herniaria 46.

27. Hirundinaria 46.

190. Hyoscyamus 43.

49. Hypericum 138.

120.

87. I.

149.

112. Juniperus 154.

L.

174. Lamium album 108.

152. Lapathum acutum 66.

39. Laureola 71.

193. Lilium Convallium 62.

93. Linaria 113.

44. Lithospermum 37.

185. Lupulus 188.

129. Lycopodium 197.

216.

M.

Malva vulgaris 12.

130. Marrubium album 111.

49. Matricaria 15.

Mat.

NOMINA PHARMACEUTICA.

Matrifylva	25.	Polypodium	192.
Melilotus	134.	Populus	182.
Mentha sylvestris	108.	Primula veris	39.
Millefolium	156.	Prunella	111.
Morsus diaboli	24.	Ptarmica	156.
Muscus caninus	213.	Pulegium	108.
Muscus cumatilis	213.	Pulmonaria	38.
Muscus clavatus	195.	Pulmonaria arborea	212.
Muscus pyxidatus	214.	Pulsatilla	100.
Nyrtillus	70.	Pyrola	77.
Nyrtus brabantica	181.		

N.

Nasturtium	117.
Nasturtium aquaticum	119.
Nymphaea	40.
Nymphaea alba	98.

O.

Ononis	130.
Origanum vulgare	110.
Orobanchus	131.

P.

Papaver	97.98.
Parietaria	184.
Paspalum	72.
Pastinaca	54.
Pentaphragmum	93.
Pernicaria	71.
Petalites	152.
Pimpinella	54.
Polygonum	26.

Q.

Quercus	174.
---------	------

R.

Ranunculus albus	100.
Ribes	45.
Ribes nigrum	45.
Rosa sylvestris	91.
Rosmarinus sylvestris	77.
Ros solis	56.
Rubus Idæus	92.
Ruta muraria	192.

S.


Sal alcali	2.
Salix	180.
Sambucus	55.
Sanicula	49.
Saponaria	79.
Satyrion	161.
Saxifraga alba	78.
Saxifraga rub	91.

R

NOMINA PHARMACEUTICA.

Scabiosa	24.	Thlaspi	117.
Scolopendrium	192.	Tilia	8.
Scordium	107.	Tormentilla	8.
Scrophularia	114.	Trifolium album	134.
Sedum minus	82.	Trifolium aquaticum	39.
Sempervivum	87.	Tussilago	152.
Serpyllum	111.		
Sigillum Salomonis	63.		
Sinapis	121.	V.	
Solanum	44.		
Sophia	120.	Valeriana	10.
Sorbus aucuparia	90.	Verbascum	43.
Spina cervina	44.	Verbena	6.
Sporocarpium citrina	151.	Veronica	4.
Succisa	24.	Vincetoxicum	46.
Symphytum	38.	Viola	158.
		Virga aurea	153.
		Viscum	180.
		Ulmaria	9.
		Ulmus	48.
Tanacetum	150.	Urtica major	173.
Taraxacum	144.	Urtica	111.

T.



EXPLICATIO LITERARUM

TAB. I. FIG. I.

Iconographiæ Agri Fridrichsdalensis.

- | | |
|--|---|
| <p>A. Palatium, vulgo <i>Fri-
drichsdals- Slot.</i></p> <p>B. Hortus major.</p> <p>C. Amnis.</p> <p>D. Sylva <i>Hare-Skov.</i></p> <p>E. Sylva <i>Nasset.</i></p> <p>F. Sylva <i>Frue-Skov.</i></p> <p>G. Nemus <i>Spare-Skiud.</i></p> <p>H. Arvum <i>Teilvang.</i></p> <p>I. Arvum <i>Ladegaards-
vang.</i></p> <p>K. Arvum <i>Deyligheden.</i></p> <p>L. Agri.</p> <p>M. Pagus <i>Bagsværd.</i></p> <p>N. Oppidum <i>Langby.</i></p> <p>O. Lucus <i>Brankebakke.</i></p> <p>P. Lucus <i>Bremerholm.</i></p> <p>Q. Pagus <i>Virum.</i></p> | <p>a. Hortus minor.</p> <p>b. Molendinum.</p> <p>c. Edes privatorum.</p> <p>d. Diverforium.</p> <p>e. Paludes.</p> <p>f. Palus major nemoris.</p> <p>g. Palus minor.</p> <p>h. Paludosa.</p> <p>i. Alheta.</p> <p>k. Palus turfosa.</p> <p>l. Tuguria, vulgo <i>Kulbuse.</i></p> <p>m. Mons cochlearis.</p> <p>n. Piscinæ.</p> <p>o. Domiilium hortulani.</p> <p>p. Hesperides.</p> |
|--|---|



EXPLICATIO TABULÆ II.


(Fig. II. III. IV. vid. pag. 193.)

FIG. V. VI.

- a. *Jungermannia* Capsula quadrivalvis.
- b. Ejus fructificatio, quemadmodum autumnno calyci suo adhuc involuta conspicitur.
- b. * Vagina hæc sive Calyx dissectus ejusque lacinie reflexæ.
- c. 1. Filamenta versus centrum conniventia.
- c. 2. Eadem lente aucta.
- c. 3. Eadem ex calyce exempta.
- c. 4. Stamen microscopio auctum.
- d. Capsula successu temporis prodiens cum suo stylo, & staminibus subtus adhuc hærentibus exaridis.
- d. * Eadem lente vitrea aucta.
- e. Pedunculus verno tempore subito elongatus.

Typis JOH. HENR. HEITZII, Univ. Typogr.

Indigeros



743. *Sparganium natans*. Tab. 260.
 748. *Carex leporina*. 294.
 752. ——— *muricata*. 284.
 755. ——— *canescens*. 285.
 767. ——— *panicea*. 261.
 796. ——— *Salix caprea*. 245.
 802. *Hippophaë rhamnoides*. 265.
 852. *Phascum acaule*. 249.
 853. ——— *subulatum*. 249.
 858. *Polytrichum commune*. 295.
 860. *Mnium pellucidum*. 300.
 861. ——— *fontanum*. 298.
 863. ——— *androgynum*. 299.
 943. *Marchantia conica*. 274.
 947. *Riccia fluitans*. 275.
 993. *Lichen chalybeiformis*. 262.
 1039. *Agaricus Cantharellus*. 264.
 1071. *Peziza punctata*. 288.



AUCTORES

quorum Icones in hoc Opere citantur.

Act. Paris. (Hist. de l'Acad. Royale des Sciences

Acta Hafniensia 1671.

MORISON *Historia Plantarum.*

STEERBECK *Fungorum Theatrum.*

MICHELII *Nova Genera Plantarum.*

VAILLANT *Botanicon Parisiense.*

DILLENII *Historia Muscorum.*

HALLERI *Enumeratio Stirpium Helveticarum.*

SCHEUCHZERI *Agrostographia.*

BUXBAUMII *Centuriæ.*

OEDERI *Flora Danica.*

Acta Holmiensia.

SCHÆFFERI *Fungorum Bavaric. Icones.*

MULLERI *Monograph. de Boletto bovino.*

Accedit

PAULI *Flora Oeconomica lingua vernacula.*






Fig. II.



Fig. III.



